



5. (a) The diagram shows where two metal spheres, S_1 and S_2 , touch the surface of water (in a tray).



S_2 remains stationary while S_1 is made to vibrate up and down, so waves spread outwards from it across the water. The wavelength of the waves is 20 mm. A displacement-time graph is given for point P.



- (i) Calculate the wave speed. [2]

.....

.....


.....

- (ii) S_1 and S_2 are now made to vibrate **in phase**, to act as wave sources of equal amplitude. Carefully sketch, on the lower two grids **opposite**, displacement-time graphs for point P, due to:


I. waves from S_2 ; [1]

II. the superposition of waves from S_1 and S_2 . [1]





- (iii) A student suggests that, with S_1 and S_2 vibrating as before, the displacement at Q (see diagram below) will be zero all the time.



Evaluate whether or not she is correct, explaining your reasoning.

[3]

.....

.....

.....

.....

.....



- (b) Light of wavelength 590 nm is incident normally on a plane diffraction grating. The centres of the grating's slits are 2000 nm apart.

- (i) Calculate the angle to the normal at which **third** order beams emerge. [2]


.....

.....

.....

.....

- (ii) The diagram shows light emerging **at this angle** from two adjacent slits in the grating.



- State the value of x . [1]

.....

