



Surname	Centre Number	Candidate Number
First name(s)		0



GCSE

3410U20-1



S24-3410U20-1

FRIDAY, 17 MAY 2024 – MORNING

CHEMISTRY – Unit 2:
Chemical Bonding, Application of Chemical Reactions
and Organic Chemistry
FOUNDATION TIER

1 hour 45 minutes

ADDITIONAL MATERIALS

In addition to this examination paper you will need a calculator and a ruler.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	6	
2.	6	
3.	5	
4.	9	
5.	8	
6.	6	
7.	11	
8.	9	
9.	7	
10.	7	
11.	6	
Total	80	

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

Question 8(a) is a quality of extended response (QER) question where your writing skills will be assessed.

The Periodic Table is printed on the back cover of this paper and the formulae for some common ions on the inside of the back cover.






JUN243410U20101

Answer **all** questions.

1. (a) A student carried out an experiment to prepare magnesium chloride crystals.

Diagrams **A**, **B** and **C** show the stages of the experiment she carried out. The stages are **not** in the correct order.



- (i) I. Give the **letter** that shows the **first** stage of the experiment. [1]

Letter

- II. Give the **letter** of the stage that shows filtration. [1]

Letter

- (ii) The gas formed in the reaction pops with a lighted splint.

Underline the name of this gas. [1]

oxygen

hydrogen

carbon dioxide

- (iii) Magnesium chloride contains Mg^{2+} and Cl^- ions.

Tick (✓) the box next to the formula of magnesium chloride. [1]

MgCl_2 ☐



Mg_2Cl ☐

MgCl ☐


Mg_2Cl_2 ☐

- (b) The word equations below show reactions of acids.

Underline the substance in each box that correctly completes the equation. [2]



2. (a) The diagram shows a cell used in the extraction of aluminium from aluminium oxide.



- (i) Underline the correct word in the brackets to complete each sentence. [3]

The process of extracting aluminium from aluminium oxide is called (**corrosion** / **electrolysis** / **cracking**).

Aluminium ions, Al^{3+} , move towards the negative electrode because (**opposite** / **similar** / **neutral**) charges attract.

Aluminium leaves the cell as a (**gas** / **solid** / **liquid**).

- (ii) Balance the **symbol** equation that shows the overall reaction. [1]



(b) Tick (✓) the box next to the **two** properties of aluminium that makes it suitable for making aeroplane wings. [2]

low density

☐

resists corrosion

☐

good thermal conductor

☐

non-toxic

☐

shiny

☐

6



3. (a) When a mixture of zinc and copper(II) oxide is heated it ignites and burns brightly.
When silver is heated with copper(II) oxide no reaction takes place.

- (i) List silver, copper and zinc in order of reactivity. [1]


Most reactive

.....

Least reactive

- (ii) The boxes below show the word equation and an incomplete symbol equation for the reaction taking place between zinc and copper(II) oxide.

Complete the symbol equation. [2]



- (b) A **solid** precipitate of silver chloride is formed when silver nitrate solution is added to sodium chloride solution.

- (i) Complete the equation for this reaction by writing the **state symbol** for the silver chloride formed in the brackets. [1]



- (ii) Underline the correct word(s) in the bracket to complete the sentence. [1]

The reaction shows that silver chloride is (**soluble** / **insoluble** / **a mixture**) in water.



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**




4. (a) Flying Saucers sweets contain sherbet. Sherbet is a mixture of two solid substances – citric acid and sodium hydrogencarbonate.

When put in your mouth, the mixture fizzes and goes cold as it dissolves in your saliva.



A student investigated the temperature change during the reaction between citric acid and a solution of sodium hydrogencarbonate.

Some sodium hydrogencarbonate solution was added to a polystyrene cup and the temperature of the solution was recorded. Citric acid was added to the solution and the temperature was recorded every 10 seconds for 60 seconds.



The results are shown in the table below.


Time (s)	0	10	20	30	40	50	60
Temperature (°C)	21.0	19.0	17.8	17.2	17.0	17.1	17.3

(i) Complete the grid below by plotting the results for 10, 20 and 30 seconds.

The results for 0, 40, 50 and 60 seconds have been plotted for you.

Draw a suitable line using all the results.

[3]



(ii) Use the graph to answer parts I and II.

I. Circle the maximum **change** in temperature during the reaction. [1]

3.7 °C 4.0 °C 21.0 °C 17.3 °C

II. This is described as an endothermic reaction. [1]

Underline the correct word(s) in the brackets to complete the sentence.

In an endothermic reaction the temperature
(**increases** / **stays the same** / **decreases**).

III. The temperature of the contents of the cup was recorded two hours later.

Underline the most likely temperature reading. [1]

4.0 °C 18.0 °C 17.3 °C 21.0 °C

(iii) All the temperature readings were higher than expected due to heat being taken in from the surroundings.

Tick (✓) the box next to the change that would **not** result in the temperature readings being closer to their expected values. [1]

put bubble wrap around the cup

☐

put a lid on the cup

☐

use a cup made from copper

☐

(b) Calculate the relative formula mass (M_r) of sodium hydrogencarbonate, NaHCO_3 . [2]

$$A_r(\text{H}) = 1$$

$$A_r(\text{C}) = 12$$

$$A_r(\text{O}) = 16$$

$$A_r(\text{Na}) = 23$$

$$M_r = \dots\dots\dots$$



5. (a) The table shows information about the Haber process and the contact process.

	Haber process	Contact process
Raw materials	nitrogen comes from the air hydrogen is made from natural gas	sulfur is made from impurities in natural gas oxygen comes from the air
Conditions	iron catalyst 450 °C 200 atm	vanadium(V) oxide catalyst 450 °C 1 atm
Equation	$\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$	$2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3$

Use the information in the table and your own knowledge to answer this question.


State whether the statements below are **true** or **false**.

[3]

	True or false?
Both processes use a catalyst that is a metallic element	
Both processes are carried out at the same temperature and pressure	
Both processes are reversible reactions	
Both processes use air as a raw material	



- (b) The equation below shows the bonds that are broken and the bonds that are formed when ammonia is produced.



- (i) The energy released when one N—H bond forms is 391 kJ.

Give the total number of N—H bonds in two molecules of ammonia and show that the amount of energy released when all the bonds are formed is 2346 kJ. [1]

- (ii) The energy released when all the bonds in the **product** are formed is 2346 kJ.

The energy needed to break all the bonds in the **reactants** is 2253 kJ.

- I. Give the term that describes a reaction where more energy is released in forming bonds than is needed to break the bonds in the reactants. [1]

.....

- II. Calculate the overall energy change that takes place during this reaction. [1]

Overall energy change = kJ



- (c) Ammonia is used to make ammonium sulfate and ammonium carbonate.
- Tests can be carried out to identify the negative ion in each of these compounds.
- Draw a line from each ion to the appropriate test and observation. [2]

Ion	Test and observation
carbonate ion, CO_3^{2-}	add dilute hydrochloric acid; gas formed turns limewater milky
	carry out a flame test; flame turns brick red
	add sodium hydroxide solution; pungent smelling gas is formed
sulfate ion, SO_4^{2-}	carry out a flame test; flame turns apple green
	add barium chloride solution; white precipitate is formed




BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**




6. (a) The diagram shows the electronic changes that occur when calcium reacts with oxygen to form calcium oxide. The $\bullet$ and $\times$ symbols represent outer shell electrons.




- (i) Complete the right-hand side of the diagram by putting the **charge** of the calcium ion in the box and the **electronic structure** of the oxide ion in the brackets. [2]
- (ii) Complete and balance the symbol equation for the reaction between calcium and oxygen. [2]



- (b) The diagrams below show the outer shell electrons in an atom of hydrogen and an atom of oxygen.



Complete the following diagram to show the outer shell electrons in a molecule of water. [2]



7. (a) Crude oil can be separated into simpler mixtures, called fractions. Fractions contain hydrocarbon compounds called alkanes.

The table shows information about some of the fractions obtained from crude oil.

Fraction	Boiling point range (°C)	Number of carbon atoms in the alkanes
refinery gases	–160 to 20	1–4
petrol	20 to 240	4–12
naphtha	100 to 250	7–14
kerosene	200 to 280	11–15
diesel oil	280 to 350	15–19

Use only the information in the table to answer parts (i)–(iii).

- (i) Complete the sentence below by underlining the correct word(s). [1]

As the number of carbon atoms in an alkane increases,
the boiling point (**increases** / **decreases** / **stays the same**).

- (ii) Hexane has a boiling point of 69 °C. [1]
Give the name of the fraction that contains hexane.






.....

- (iii) One alkane is found in the kerosene and diesel oil fractions. [1]
Give the number of carbon atoms in this alkane.

.....



(b) Diagrams **A–E** show the structural formulae of some carbon compounds.

**A****B****C****D****E**

(i) Circle the molecular formula for compound **A**. [1]



(ii) Give the **letter** of the compound that is unsaturated. [1]

.....

(iii) Explain why compound **E** is **not** a hydrocarbon. [1]

.....


(iv) Give the **letter** of the compound that belongs to the family of carbon compounds with the general formula C_nH_{2n}. [1]

.....



Examiner
only

- (c) The chart shows the stages used in the laboratory to make ethanol from glucose solution.



- (i) Give the number of the stage in which distillation is used. [1]

Stage

- (ii) The formula of ethanol is $\text{C}_2\text{H}_5\text{OH}$.

Complete the structure of ethanol by adding all the atoms and bonds present. [1]



- (iii) Ethanol is the alcohol found in alcoholic drinks.

The following information appears on a bottle of wine.

Volume	750 millilitres
Alcohol by volume	13.5%

The formula below can be used to calculate the mass of ethanol in an alcoholic drink.

$$\text{mass of ethanol (g)} = 8 \times \text{volume (litres)} \times \text{alcohol by volume (\%)}$$

Use the formula to find the mass of ethanol in a bottle of wine.

[2]

$$1 \text{ litre} = 1000 \text{ millilitres}$$

Mass = g

11



8. (a) The photographs show three different methods of fighting fires.
Each method removes a different part of the fire triangle.



Method A



Method B



Method C



Examiner
only

Use the fire triangle to explain how the different methods shown in the photographs help to extinguish fires or prevent them from spreading. [6 QER]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



(b)

Fighting Fires

There are five main types of fire extinguisher – water, foam, dry powder, carbon dioxide and wet chemical. Modern fire extinguishers have a red body with a coloured band at the top. Each colour represents a different type of extinguisher, used on different types or ‘classes’ of fire.



There is no one extinguisher type that works on all classes of fire.

The chart below shows which type of extinguisher should be used on each class of fire.

TYPE FIRE EXTINGUISHER	CLASS A Flammable solids, for example paper	CLASS B Flammable liquids, for example ethanol	CLASS C Flammable gases, for example methane	CLASS D Flammable metals, for example magnesium	CLASS E Electrical equipment, for example computer	CLASS F Cooking oil, for example deep fat fryer
WATER	✓	✗	✗	✗	✗	✗
FOAM	✓	✓	✗	✗	✗	✗
DRY POWDER	✓	✓	✓	✓	✓	✗
CO ₂	✗	✓	✗	✗	✓	✗
WET CHEMICAL	✓	✗	✗	✗	✗	✓



Use only the information in the chart to answer parts (i)–(iii).

- (i) Underline the type of extinguisher used to put out a fire involving burning cooking oil. [1]

water foam dry powder carbon dioxide wet chemical


- (ii) Underline the type of extinguisher that would be the most useful in a school chemistry laboratory. [1]

water foam dry powder carbon dioxide wet chemical

- (iii) Tick (✓) the box next to the fire that should be put out using a water extinguisher. [1]



chip pan fire

☐


burning plug and socket

☐


burning butane cylinder

☐


burning waste cardboard

☐



9. The list below shows part of the reactivity series.

magnesium
zinc
iron
nickel

A student investigated the temperature rise when four different metal powders were added to excess copper(II) sulfate solution.

The same mass of each metal was added to 50 cm³ samples of copper(II) sulfate solution.

The maximum temperature for each reaction was measured. The temperature rise was calculated in each case and used to find the energy given out.



The results are shown in the table below.

Metal	Temperature rise (°C)	Energy given out (J)
magnesium	40.5	8500
zinc	33.0	6900
iron	23.2	4900
nickel	19.0	4000



- (a) In one of the reactions, the initial temperature was 19.7 °C and the maximum temperature was 52.7 °C.

State which one of the four metals was used in this reaction.

[1]

.....

- (b) Tick (✓) the box next to the conclusion the student can draw from the results.

[1]

The higher the metal in the reactivity series, the greater the energy given out

☐

The lower the metal in the reactivity series, the greater the energy given out

☐

The energy given out is not related to the metal's position in the reactivity series

☐


- (c) The word equation below represents the reaction between iron and nickel(II) sulfate solution.



Iron(II) sulfate contains Fe^{2+} and SO_4^{2-} ions.

Complete the symbol equation for the reaction.

[2]



- (d) The experiment shows that a more reactive metal will replace a less reactive metal in its compounds.

Give the term used to describe this type of reaction.

[1]

.....



- (e) When the experiment was repeated using **titanium**, the temperature rise recorded was 35.4 °C.

Calculate the energy given out during the reaction between **titanium** and 50 cm³ of copper(II) sulfate solution. Give your answer to the nearest 100 J. [2]

$$\text{energy given out (J)} = \text{volume of solution} \times 4.2 \times \text{temperature rise}$$

Energy given out = J

7




BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



10. (a) The diagram shows the apparatus used by a student to investigate the electrolysis of water.



- (i) The student collected a sample of the oxygen formed in a test tube.

Give the test the student would carry out to show the presence of oxygen in the test tube. Include the observation the student would expect. [1]

.....

.....

- (ii) Use the formula of water, H_2O .

Give the volume of hydrogen that would form in the same time as 10 cm^3 of oxygen. [1]

..... cm^3



Examiner
only

(b) The table shows information about the electrolysis of three different electrolytes. The table is incomplete.

Electrolyte	Ions present in the electrolyte		Observations	
	Positive ion(s)	Negative ion(s)	At the negative (–) electrode	At the positive (+) electrode
molten lead(II) bromide	Pb^{2+}		grey metal A formed	orange gas formed
aqueous copper(II) chloride	 and H^{+}	Cl^{-} and OH^{-}	brown metal formed	green-yellow gas B formed
aqueous compound C	Zn^{2+} and H^{+}	I^{-} and OH^{-}	grey metal formed	brown solution formed

- (i) Complete the table by adding the **symbols** of the missing **ions**. [2]
- (ii) Name substances **A**, **B** and **C**. [3]


Metal **A**

Gas **B**


Compound **C**



11. The diagrams show the structure of diamond and graphite.



diamond



graphite

(a) Name the atom being represented by a • in both diagrams.

[1]

.....

(b) Name the type of **bonding** found in both diamond and graphite.

[1]

.....



Examiner
only

(c) The table shows some properties of graphite.

Properties of graphite
soft
high melting point
insoluble in water
conducts electricity

Use only properties from the table to answer this question.

Give **two** properties of graphite that are **different** from those of diamond. Give a use relating to each property. [4]

Property 1

Use

Property 2

Use

6

END OF PAPER



[illegible]

Examiner
only



FORMULAE FOR SOME COMMON IONS

POSITIVE IONS		NEGATIVE IONS	
Name	Formula	Name	Formula
aluminium	Al^{3+}	bromide	Br^-
ammonium	NH_4^+	carbonate	CO_3^{2-}
barium	Ba^{2+}	chloride	Cl^-
calcium	Ca^{2+}	fluoride	F^-
copper(II)	Cu^{2+}	hydroxide	OH^-
hydrogen	H^+	iodide	I^-
iron(II)	Fe^{2+}	nitrate	NO_3^-
iron(III)	Fe^{3+}	oxide	O^{2-}
lithium	Li^+	sulfate	SO_4^{2-}
magnesium	Mg^{2+}		
nickel	Ni^{2+}		
potassium	K^+		
silver	Ag^+		
sodium	Na^+		
zinc	Zn^{2+}		





THE PERIODIC TABLE

Group

1 2

3

4

5

6

7

0

1	H	Hydrogen	1
---	---	----------	---

7	Li	Lithium	3
9	Be	Beryllium	4
23	Na	Sodium	11
24	Mg	Magnesium	12
39	K	Potassium	19
40	Ca	Calcium	20
86	Rb	Rubidium	37
88	Sr	Strontium	38
133	Cs	Caesium	55
137	Ba	Barium	56
223	Fr	Francium	87
45	Sc	Scandium	21
48	Ti	Titanium	22
51	V	Vanadium	23
52	Cr	Chromium	24
55	Mn	Manganese	25
56	Fe	Iron	26
59	Co	Cobalt	27
59	Ni	Nickel	28
63.5	Cu	Copper	29
65	Zn	Zinc	30
89	Y	Yttrium	39
91	Zr	Zirconium	40
93	Nb	Niobium	41
96	Mo	Molybdenum	42
99	Tc	Technetium	43
101	Ru	Ruthenium	44
103	Rh	Rhodium	45
106	Pd	Palladium	46
108	Ag	Silver	47
112	Cd	Cadmium	48
115	In	Indium	49
119	Sn	Tin	50
122	Sb	Antimony	51
127	I	Iodine	53
131	Xe	Xenon	54
139	La	Lanthanum	57
179	Hf	Hafnium	72
181	Ta	Tantalum	73
184	W	Tungsten	74
186	Re	Rhenium	75
190	Os	Osmium	76
192	Ir	Iridium	77
195	Pt	Platinum	78
197	Au	Gold	79
201	Hg	Mercury	80
204	Tl	Thallium	81
207	Pb	Lead	82
209	Bi	Bismuth	83
210	Po	Polonium	84
210	At	Astatine	85
222	Rn	Radon	86
226	Ra	Radium	88
227	Ac	Actinium	89

11	B	Boron	5
12	C	Carbon	6
14	N	Nitrogen	7
16	O	Oxygen	8
19	F	Fluorine	9
20	Ne	Neon	10
27	Al	Aluminium	13
28	Si	Silicon	14
31	P	Phosphorus	15
32	S	Sulfur	16
35.5	Cl	Chlorine	17
40	Ar	Argon	18
70	Ga	Gallium	31
73	Ge	Germanium	32
75	As	Arsenic	33
79	Se	Selenium	34
80	Br	Bromine	35
84	Kr	Krypton	36
115	In	Indium	49
119	Sn	Tin	50
122	Sb	Antimony	51
127	I	Iodine	53
131	Xe	Xenon	54
204	Tl	Thallium	81
207	Pb	Lead	82
209	Bi	Bismuth	83
210	Po	Polonium	84
210	At	Astatine	85
222	Rn	Radon	86

Key

A_r	relative atomic mass
Symbol	
Name	
Z	atomic number