2. Section Kolpakowskianae Raamsd. ex Zonn.& Veldk.

	Sp. Pl. 1: 305 (1753). Type: <i>Tulipa kolpakowskiana</i> Regel (Trudy Imp. SPeterburgsk. Bot. Sada 5: 266 (1877).
1.	Small plants up to 20 cm tall; leaves approximate, exceeding the flower; ovary with long style
_	Relatively tall plants (up to 45 cm); leaves more or less widely spaced; ovary with sessile or subsessile stigma
2.	Perianth segments yellowish, without basal blotch, outer ones greenish on the back, the edges of the upper half often notched, two-thirds as broad as and slightly longer than the inner segments; tunics brown or grayish, papery
3.	Bulb tunics fibrous, splitting, black to tawny, prolonged to soil surface, inside densely covered with curly woolly hairs; stamens one-third to two-fifths the length of perianth filaments black to purple
	slightly prolonged (if prolonged up to soil surface, in this case lined with woolly hairs at the top and glabrous below); stamens two or three times shorter than the perianth, filaments yellow
4.	Upper leaves exceeding or at least reaching the flower; aboveground part of the plant one half to two-thirds as long as the underground part; perianth segments with a dark viole blotch on both faces
_	Upper leaves do not exceed the flower; aboveground part as long as or slightly longer than the underground part; perianth segments commonly with a dark basal blotch on the inner face, rarely on both faces
5. -	Upper part of stem and peduncle pubescent; leaves never lying on the soi surface
6.	Bulb tunics not prolonged, coriaceous; plants of relatively slender habit; leaves narrow lanceolate, do not exceed the flower; flower pendulous in the bud, perianth segments yellow, without blotch, elliptic, at the apex obtuse or sinuate, after anthesis turn brownish in the upper part; filaments yellow, narrow triangular (dilated at the base); ovary not bottle like; grows in piedmonts on slightly saline soils
_	Bulb tunics usually prolonged, thin coriaceous; plants of relatively stocky habit (sometimes with almost sessile flowers); leaves usually exceeding or at least reaching the flower perianth segments red, yellow, orange, pink or motley, acute or with a small tip at apex sometimes obtuse; grows in foothills and mountains on gravelly and stong slopes
7.	Bulb tunics tough, almost scaly, dark in colour; leaves approximate to more or less widely spaced; perianth segments usually red with a small black marginless basal blotch sometimes yellow-red or yellow, and pink colour; filaments commonly black in the lower part and purple in the upper part, sometimes entirely black or purple
_	Bulb tunics thin coriaceous, darkish brown in colour; leaves always almost verticillate perianth segments yellow, reddish-orange or light red9

yellow	oth segments of red or yellow. In the segments p					3. T.	korolk	owii
- I CHan	itii segilielits p	mik in coloui	•••••	• • • • • • • • • • • • • • • • • • • •	т. 2	i. Korotko	wii 1. 70	Beu
base v shape – The sl	petals unbent with almost padd); ovary is paddape of the persh	rallel margin le yellow rianth is diffe	s, but with rent, filam	a slightly i ents with th	inflated part	t in the mi rt at the ba	ddle (bo 10 ase; ova	ottle- ry is
brown - Perian pale	nth segments y hish in their ba hith reddish-ora brownish in	sal part; ova ange or light the basal	ry pale yello red, with b part ar	ow6 lack or dark ld red or	6. T. intern brown spo reddish	nedia var. ts; filamen in the	interme ts bicolo upper	edia ored: part
								_