



Free virions: NEW model vs CLASSICAL BMVR
Solid navy = NEW model (renewal / burst-aware) | Dashed orange + circles = CLASSICAL BMVR (constant release rate p)


(a) supercritical, $\mu = 0$
 $\beta = 1.0, \mu = 0.0, \delta = 0.1 \mid \gamma T = 0.04, c = 0.25 \mid R_0 = 1.76$




(b) supercritical, $\mu > 0$
 $\beta = 1.0, \mu = 0.2, \delta = 0.05 \mid \gamma T = 0.03, c = 0.2 \mid R_0 = 2.10$




(c) fast burst (large δ)
 $\beta = 1.0, \mu = 0.0, \delta = 0.5 \mid \gamma T = 0.25, c = 0.3 \mid R_0 = 2.5$



(d) subcritical $R_0 < 1$
 $\beta = 1.0, \mu = 0.0, \delta = 0.1 \mid \gamma T = 0.01, c = 0.4 \mid R_0 = 0.27$



(e) high intracellular death
 $\beta = 1.0, \mu = 0.9, \delta = 0.1 \mid \gamma T = 0.5, c = 0.25 \mid R_0 = 2.63$



(f) free-virion seed only ($I_0 = 0$)
 $\beta = 1.0, \mu = 0.0, \delta = 0.1 \mid \gamma T = 0.05, c = 0.2 \mid R_0 = 2.7$

