
Problem Set 4 for Machine Learning 15 Fall

Jingyuan Liu
AndrewId: jingyual
jingyual@andrew.cmu.edu

1 VC dimension

1.1 Show the VC dimension of linear classifier

To prove that the linear classifier h with x in R^n has the VC dimension of $n + 1$, we need to prove that $VCdim(h_n) \geq n + 1$, and then prove that $VCdim(h_n) \leq n + 1$.

(a). Prove $VCdim(h_n) \geq n + 1$

First of all, we could use the Mathematical Induction to prove that $VCdim(H) \geq n + 1$:

For $n = 1$, it is easy to get $VCdim(h_1) \geq 2$

For $n = 2$, we could also get $VCdim(h_2) \geq 3$, which could be proved using following figures:




Figure 1: The representation of $VCdim(h_2)$

For $n = i$, we assume that the $VCdim(h_i) \geq i + 1$. We could form a matrix of $(i+1) \times (i+1)$, with the rank of $i+1$, since the VCdim is $i+1$.

For $n = i+1$, we could form a matrix of $(i+2) \times (i+2)$, since the variable x have one more independent dimension. Therefore, $VCdim(h_{i+1}) \geq VCdim(h_i) + 1 \geq i + 2$.

Therefore, we have $VCdim(h_n) \geq n + 1$.

(b). Prove $VCdim(h_n) \leq n + 1$

Then we should prove that $VCdim(h_n) \leq n + 1$:

Suppose we could shatter $n+2$ points, then using the convex combinations of points in C , we could separate points into $S1$ and $S2$, that:

$$conv(s1) \cap conv(s2) \neq \emptyset \quad (1)$$

However, if $VCdim(h_n) = n + 2$, then we could split points to half space contains $s1$ and the complement of the half space contains $s2$. This implies that the both half space contains the convex hull of $s1$ and the complement of the half space the contains the convex hull of $s2$. Thus, we get:

$$conv(s1) \cap conv(s2) = \emptyset \quad (2)$$

These two observations are contradictory to each other. So we know that for the $n+2$ points here, we could not use the linear classifier to shatter all of them.

In conclusion, we prove that $VCdim(h_n) \geq n + 1$ and $VCdim(h_n) \leq n + 1$ for the linear classifier, then we have $VCdim(h_n) = n + 1$

1.2 Show the VC dimension of axis-aligned boxes

Similarly, for this case, to prove that the axis-aligned boxes classifier h with x in R^n has the VC dimension of $2n$, we need to prove that $VCdim(h_n) \geq 2n$, and then prove that $VCdim(h_n) \leq 2n$.

(a). Prove $VCdim(h_n) \geq 2n$

First of all, we need to prove that for any case, $VCdim(h_n) \geq 2n$, which means if the x has n dimensions, we can use the axis-aligned boxes classifiers to shatter $2n$ points.

Suppose we have n dimensions, then we can map all the x to a dimension i . In the dimension i , we can have x_{max}^i and x_{min}^i . Therefore, in this dimension i , we could always shatter at least 2 points via a split between the x_{max}^i and x_{min}^i .

Considering we have n dimensions, and the mapping of each dimension of the data is independent to the mapping of other dimension of the data. Therefore, we could at least shatter $2 \times n$ points via the axis-aligned boxes. Therefore, we have $VCdim(h_n) \geq 2n$.

(b). Prove $VCdim(h_n) \leq 2n$

Then we need to prove that $VCdim(h_n) \leq 2n$. Suppose we have $2n + 1$ points, we could always find the $2n$ “boundries”, which is the min value and max value for each dimension, and form an “area”. Then the $2n+1$ th points will be guaranteed to appear within the selected “area”.

For any $2n+1$ points, we could always transfer to the above mentioned scenario. In this scenario, we could not classify the $2n+1$ th point, because it is contained within the “area”, and no rules are guaranteed to rightly classify it.

Therefore, by mapping the data to each dimension and form an “area”, we could prove that $VCdim(h_n) \leq 2n$.

In conclusion, we prove that $VCdim(h_n) \geq 2n$ and $VCdim(h_n) \leq 2n$ for the axis-aligned boxes, then we have $VCdim(h_n) = 2n$

2 AdaBoost

3 Gaussian Mixture Model

3.1 Show the expectation

$$E(x) = \int p(x)dx = \sum_k p(x_k)x_k \quad (3)$$

$$= \sum_k \pi_k E_k(x) \quad (4)$$

We know that for each k , the implicit distribution of x is a Gaussian distribution, and $E_k(x) = \mu_k$, μ_k is the mean of the k th gaussian distribution, so we have:

$$E(x) = \sum_k \pi_k \mu_k \quad (5)$$

3.2 Show the covariance

$$\text{cov}(x) = E(xx^T) - E(x)E(x)^T \quad (6)$$

$$= \sum_k \pi_k E_k(xx^T) - E(x)E(x)^T \quad (7)$$

$$= \sum_k \pi_k (\Sigma_k + \mu_k \mu_k^T) - E(X)E(x)^T \quad (8)$$

4 K-Means

5 Collaboration

I dicussed with Zheng Chen with problem 1.2 on how to calculate the parameters, and he also informed me about 2.2.1(b).