

6.4: Values of the Trigonometric Functions

E. Kim


MTH 151

All notation and terminology is based on Swokowski, Cole. *Algebra and Trigonometry: with analytic geometry*. Classic 12th Edition.


Accompanying handout:

- ▶ Black-and-white: <http://www.uwlax.edu/faculty/ekim/resources/unit-circle.pdf>
- ▶ Color: <http://www.uwlax.edu/faculty/ekim/resources/unit-circle-color.pdf>


Goal



Goal



Goal



Why understand the unit circle?


For all special angles, we can compute sine, cosine, etc. by knowing these values only for 30° , 45° , and 90° .

Why understand the unit circle?


For all special angles, we can compute sine, cosine, etc. by knowing these values only for 30° , 45° , and 90° .

For the other special angles, just change the sign as appropriate.


The First Quadrant




The First Quadrant




The First Quadrant



The First Quadrant




The First Quadrant



What is a reference angle?


Take a nonquadrantal¹ angle θ .



¹Nonquadrantal means that θ is not a multiple of 90° .

What is a reference angle?

Take a nonquadrantal¹ angle θ .




θ_R , the reference angle

The acute angle made by

¹Nonquadrantal means that θ is not a multiple of 90° .

What is a reference angle?

Take a nonquadrantal¹ angle θ .



θ_R , the reference angle


The acute angle made by

► terminal side of θ

¹Nonquadrantal means that θ is not a multiple of 90° .

What is a reference angle?

Take a nonquadrantal¹ angle θ .




θ_R , the reference angle

The acute angle made by

- ▶ terminal side of θ
- ▶ x -axis

¹Nonquadrantal means that θ is not a multiple of 90° .


Formula for reference angle θ_R in every quadrant



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1


$$\theta_R = \theta$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1


$$\theta_R = \theta$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1


$$\theta_R = \theta$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$



Formula for reference angle θ_R in every quadrant


θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$



Formula for reference angle θ_R in every quadrant


θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$



Formula for reference angle θ_R in every quadrant


θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$



Formula for reference angle θ_R in every quadrant


θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2


$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3

$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2


$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3

$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2


$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3

$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2


$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3

$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3


$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$

θ in Quadrant 4

$$\theta_R = 2\pi - \theta$$

$$\theta_R = 360^\circ - \theta$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3


$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$

θ in Quadrant 4

$$\theta_R = 2\pi - \theta$$

$$\theta_R = 360^\circ - \theta$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3


$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$

θ in Quadrant 4

$$\theta_R = 2\pi - \theta$$

$$\theta_R = 360^\circ - \theta$$



Formula for reference angle θ_R in every quadrant

θ in Quadrant 1

$$\theta_R = \theta$$

θ in Quadrant 2

$$\theta_R = \pi - \theta$$

$$\theta_R = 180^\circ - \theta$$

θ in Quadrant 3


$$\theta_R = \theta - \pi$$

$$\theta_R = \theta - 180^\circ$$


θ in Quadrant 4

$$\theta_R = 2\pi - \theta$$


$$\theta_R = 360^\circ - \theta$$




Example: $\theta = 315^\circ$



Example: $\theta = 315^\circ$




Example: $\theta = 315^\circ$




$$\theta_R = 360^\circ - 315^\circ$$

Example: $\theta = 315^\circ$




$$\theta_R = 360^\circ - 315^\circ = 45^\circ$$


Example: $\theta = \frac{5\pi}{6}$




Example: $\theta = \frac{5\pi}{6}$




Example: $\theta = \frac{5\pi}{6}$




Example: $\theta = \frac{5\pi}{6}$




Example: $\theta = 4$. (Note this is four **radians**, not degrees!)



Example: $\theta = 4$. (Note this is four **radians**, not degrees!)



Example: $\theta = 4$. (Note this is four **radians**, not degrees!)



What if θ is not between 0° and 360° ?

What if θ is not between 0° and 360° ?

First, find the coterminal angle to θ between 0° and 360° .

What if θ is not between 0° and 360° ?

First, find the coterminal angle to θ between 0° and 360° .

Example: $\theta = -240^\circ$.

What if θ is not between 0° and 360° ?

First, find the coterminal angle to θ between 0° and 360° .

Example: $\theta = -240^\circ$.

$\theta = -240^\circ$ is coterminal to $-240^\circ + 360^\circ$

What if θ is not between 0° and 360° ?

First, find the coterminal angle to θ between 0° and 360° .

Example: $\theta = -240^\circ$.

$\theta = -240^\circ$ is coterminal to $-240^\circ + 360^\circ = 120^\circ$

What if θ is not between 0° and 360° ?


First, find the coterminal angle to θ between 0° and 360° .

Example: $\theta = -240^\circ$.


$\theta = -240^\circ$ is coterminal to $-240^\circ + 360^\circ = 120^\circ$

To find θ_R , use $\theta = 120^\circ$.


Example: $\theta = -240^\circ$, but using 120°



Example: $\theta = -240^\circ$, but using 120°




Example: $\theta = -240^\circ$, but using 120°



$$\theta_R = 180^\circ - 120^\circ$$

Example: $\theta = -240^\circ$, but using 120°




$$\theta_R = 180^\circ - 120^\circ = 60^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$

$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

Reference angles and exact values of sin, cos, and tan

$$\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 120^\circ$$


$$\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \theta = 60^\circ$$




$$\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 240^\circ$$

$$\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right), \theta = 300^\circ$$

If $\theta = \frac{5\pi}{6}$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$



If $\theta = \frac{5\pi}{6}$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$



If $\theta = \frac{5\pi}{6}$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = \frac{\pi}{6}$

If $\theta = \frac{5\pi}{6}$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$


Reference angle is $\theta_R = \frac{\pi}{6}$

$$\sin \frac{\pi}{6} = \frac{1}{2} \qquad \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2} \qquad \tan \frac{\pi}{6} = \frac{\sqrt{3}}{3}$$

If $\theta = \frac{5\pi}{6}$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = \frac{\pi}{6}$


$$\sin \frac{\pi}{6} = \frac{1}{2} \quad \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2} \quad \tan \frac{\pi}{6} = \frac{\sqrt{3}}{3}$$



If $\theta = \frac{5\pi}{6}$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = \frac{\pi}{6}$


$$\sin \frac{\pi}{6} = \frac{1}{2} \quad \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2} \quad \tan \frac{\pi}{6} = \frac{\sqrt{3}}{3}$$



If $\theta = \frac{5\pi}{6}$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$


Reference angle is $\theta_R = \frac{\pi}{6}$

$$\sin \frac{\pi}{6} = \frac{1}{2} \quad \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2} \quad \tan \frac{\pi}{6} = \frac{\sqrt{3}}{3}$$




$$\sin \frac{5\pi}{6} = +\frac{1}{2} \quad \cos \frac{5\pi}{6} = -\frac{\sqrt{3}}{2} \quad \tan \frac{5\pi}{6} = -\frac{\sqrt{3}}{3}$$

If $\theta = 315^\circ$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$



If $\theta = 315^\circ$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$



$$\theta_R = 360^\circ - 315^\circ = 45^\circ$$

If $\theta = 315^\circ$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = 45^\circ$

If $\theta = 315^\circ$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = 45^\circ$

$$\sin 45^\circ = \frac{\sqrt{2}}{2} \qquad \cos 45^\circ = \frac{\sqrt{2}}{2} \qquad \tan 45^\circ = 1$$


If $\theta = 315^\circ$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = 45^\circ$

$$\sin 45^\circ = \frac{\sqrt{2}}{2}$$

$$\cos 45^\circ = \frac{\sqrt{2}}{2}$$

$$\tan 45^\circ = 1$$




If $\theta = 315^\circ$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = 45^\circ$

$$\sin 45^\circ = \frac{\sqrt{2}}{2}$$

$$\cos 45^\circ = \frac{\sqrt{2}}{2}$$


$$\tan 45^\circ = 1$$



If $\theta = 315^\circ$, find $\sin \theta$, $\cos \theta$, and $\tan \theta$

Reference angle is $\theta_R = 45^\circ$

$$\sin 45^\circ = \frac{\sqrt{2}}{2} \quad \cos 45^\circ = \frac{\sqrt{2}}{2} \quad \tan 45^\circ = 1$$



$$\sin 315^\circ = -\frac{\sqrt{2}}{2} \quad \cos 315^\circ = +\frac{\sqrt{2}}{2} \quad \tan 315^\circ = -1$$

Summary of using reference angles

1. Find the reference angle θ_R for your angle θ .
2. Compute \sin , \cos , and \tan for the reference angle θ_R .
3. Adjust the sign based on the quadrant of terminal side of θ .

Finding angles with a calculator

Inverse trigonometric functions

Problem

If θ is an acute angle and $\sin \theta = 0.6635$, what is θ ?

Inverse trigonometric functions

Problem

If θ is an acute angle and $\sin \theta = 0.6635$, what is θ ?

$$\text{If } \sin \theta = k, \text{ then } \theta = \sin^{-1} k.$$

Inverse trigonometric functions

Problem

If θ is an acute angle and $\sin \theta = 0.6635$, what is θ ?

If $\sin \theta = k$, then $\theta = \sin^{-1} k$.

$$\theta = \sin^{-1}(0.6635)$$

Inverse trigonometric functions

Problem

If θ is an acute angle and $\sin \theta = 0.6635$, what is θ ?

If $\sin \theta = k$, then $\theta = \sin^{-1} k$.

$$\theta = \sin^{-1}(0.6635) \approx 41.57^\circ \approx 0.7255$$

Inverse trigonometric functions

Problem

If θ is an acute angle and $\sin \theta = 0.6635$, what is θ ?

If $\sin \theta = k$, then $\theta = \sin^{-1} k$.

$$\theta = \sin^{-1}(0.6635) \approx 41.57^\circ \approx 0.7255$$

Inverse trigonometric functions

Problem

If θ is an acute angle and $\sin \theta = 0.6635$, what is θ ?

$$\text{If } \sin \theta = k, \text{ then } \theta = \sin^{-1} k.$$

$$\theta = \sin^{-1}(0.6635) \approx 41.57^\circ \approx 0.7255$$

degrees radians

Inverse trigonometric functions

Problem

If θ is an acute angle and $\sin \theta = 0.6635$, what is θ ?

$$\text{If } \sin \theta = k, \text{ then } \theta = \sin^{-1} k.$$

$$\theta = \sin^{-1}(0.6635) \approx 41.57^\circ \approx 0.7255$$

degrees radians

$$\text{If } \cos \theta = k, \text{ then } \theta = \cos^{-1} k.$$

$$\text{If } \tan \theta = k, \text{ then } \theta = \tan^{-1} k.$$

What about csc, sec, and cot?

Use the reciprocal formulas:

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

What about csc, sec, and cot?

Use the reciprocal formulas:

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

Example: $\csc \theta = 2$

What about csc, sec, and cot?

Use the reciprocal formulas:

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

Example: $\csc \theta = 2$

Convert to

$$\frac{1}{\sin \theta} = 2$$

What about csc, sec, and cot?

Use the reciprocal formulas:

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

Example: $\csc \theta = 2$

Convert to

$$\frac{1}{\sin \theta} = 2$$

Take reciprocal of both sides

$$\sin \theta = \frac{1}{2}$$

What about csc, sec, and cot?

Use the reciprocal formulas:

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

Example: $\csc \theta = 2$

Convert to

$$\frac{1}{\sin \theta} = 2$$

Take reciprocal of both sides

$$\sin \theta = \frac{1}{2}$$

Use inverse sine function (also called **arcsin** or **asin**)

$$\theta = \sin^{-1} \left(\frac{1}{2} \right)$$

What about csc, sec, and cot?

Use the reciprocal formulas:

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

Example: $\csc \theta = 2$

Convert to

$$\frac{1}{\sin \theta} = 2$$

Take reciprocal of both sides

$$\sin \theta = \frac{1}{2}$$

Use inverse sine function (also called **arcsin** or **asin**)

$$\theta = \sin^{-1} \left(\frac{1}{2} \right) = 30^\circ$$

An example of each inverse trig function

$$\sin \theta = 0.5$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

An example of each inverse trig function

$$\sin \theta = 0.5$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\cos \theta = 0.5$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

An example of each inverse trig function

$$\sin \theta = 0.5$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\cos \theta = 0.5$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\tan \theta = 0.5$$

$$\theta = \tan^{-1}\left(\frac{1}{2}\right) \approx 26.57^\circ \approx 0.4636$$

An example of each inverse trig function

$$\sin \theta = 0.5$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\csc \theta = 2$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\cos \theta = 0.5$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\tan \theta = 0.5$$

$$\theta = \tan^{-1}\left(\frac{1}{2}\right) \approx 26.57^\circ \approx 0.4636$$

An example of each inverse trig function

$$\sin \theta = 0.5$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\csc \theta = 2$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\cos \theta = 0.5$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\sec \theta = 2$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\tan \theta = 0.5$$

$$\theta = \tan^{-1}\left(\frac{1}{2}\right) \approx 26.57^\circ \approx 0.4636$$

An example of each inverse trig function

$$\sin \theta = 0.5$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\csc \theta = 2$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\cos \theta = 0.5$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\sec \theta = 2$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\tan \theta = 0.5$$

$$\theta = \tan^{-1}\left(\frac{1}{2}\right) \approx 26.57^\circ \approx 0.4636$$

$$\cot \theta = 2$$

$$\theta = \tan^{-1}\left(\frac{1}{2}\right) \approx 26.57^\circ \approx 0.4636$$

An example of each inverse trig function

$$\sin \theta = 0.5$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\csc \theta = 2$$

$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = 30^\circ \approx 0.5236$$

$$\cos \theta = 0.5$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\sec \theta = 2$$

$$\theta = \cos^{-1}\left(\frac{1}{2}\right) = 60^\circ \approx 1.0472$$

$$\tan \theta = 0.5$$

$$\theta = \tan^{-1}\left(\frac{1}{2}\right) \approx 26.57^\circ \approx 0.4636$$

$$\cot \theta = 2$$

$$\theta = \tan^{-1}\left(\frac{1}{2}\right) \approx 26.57^\circ \approx 0.4636$$

Column on right copies column on left
because of reciprocal identities

Since $f(x) = \sin(x)$ is periodic, what is $\sin^{-1} k$ giving you?

Inverse Sine

If you put $\sin^{-1} k$ into your calculator, the answer will be an angle

- ▶ in $[-\frac{\pi}{2}, \frac{\pi}{2}]$ using radian mode
- ▶ in $[-90^\circ, 90^\circ]$ in degree mode

Since $f(x) = \sin(x)$ is periodic, what is $\sin^{-1} k$ giving you?

Inverse Sine

If you put $\sin^{-1} k$ into your calculator, the answer will be an angle

- ▶ in $[-\frac{\pi}{2}, \frac{\pi}{2}]$ using radian mode
- ▶ in $[-90^\circ, 90^\circ]$ in degree mode

Inverse Cosine

If you put $\cos^{-1} k$ into your calculator, the answer will be an angle

- ▶ in $[0, \pi]$ using radian mode
- ▶ in $[0, 180^\circ]$ in degree mode

Since $f(x) = \sin(x)$ is periodic, what is $\sin^{-1} k$ giving you?

Inverse Sine

If you put $\sin^{-1} k$ into your calculator, the answer will be an angle

- ▶ in $[-\frac{\pi}{2}, \frac{\pi}{2}]$ using radian mode
- ▶ in $[-90^\circ, 90^\circ]$ in degree mode

Inverse Cosine

If you put $\cos^{-1} k$ into your calculator, the answer will be an angle

- ▶ in $[0, \pi]$ using radian mode
- ▶ in $[0, 180^\circ]$ in degree mode

Inverse Tangent

If you put $\tan^{-1} k$ into your calculator, the answer will be an angle

- ▶ in $(-\frac{\pi}{2}, \frac{\pi}{2})$ using radian mode
- ▶ in $(-90^\circ, 90^\circ)$ in degree mode

Examples with negative values

$$\sin \theta = -0.5$$

$$\theta = \sin^{-1}\left(-\frac{1}{2}\right) = -30^\circ \approx -0.5236$$

Examples with negative values

$$\sin \theta = -0.5$$

$$\theta = \sin^{-1}\left(-\frac{1}{2}\right) = -30^\circ \approx -0.5236$$

$$\cos \theta = -0.5$$

$$\theta = \cos^{-1}\left(-\frac{1}{2}\right) = 120^\circ \approx 2.0944$$

Examples with negative values

$$\sin \theta = -0.5$$

$$\theta = \sin^{-1}\left(-\frac{1}{2}\right) = -30^\circ \approx -0.5236$$

$$\cos \theta = -0.5$$

$$\theta = \cos^{-1}\left(-\frac{1}{2}\right) = 120^\circ \approx 2.0944$$

$$\tan \theta = -0.5$$

$$\theta = \tan^{-1}\left(-\frac{1}{2}\right) \approx -26.57^\circ \approx -0.4636$$

Examples with negative values

$$\sin \theta = -0.5$$

$$\theta = \sin^{-1}\left(-\frac{1}{2}\right) = -30^\circ \approx -0.5236$$

$$\cos \theta = -0.5$$

$$\theta = \cos^{-1}\left(-\frac{1}{2}\right) = 120^\circ \approx 2.0944$$

$$\tan \theta = -0.5$$

$$\theta = \tan^{-1}\left(-\frac{1}{2}\right) \approx -26.57^\circ \approx -0.4636$$

If the calculator doesn't give you the angle θ you wanted...

...use reference angles to find the angle you want!

Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Putting $\tan^{-1}(-0.4623)$ in the calculator (in degree mode).

Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Putting $\tan^{-1}(-0.4623)$ in the calculator (in degree mode).

- Get $\approx -24.8^\circ$.

Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Putting $\tan^{-1}(-0.4623)$ in the calculator (in degree mode).

► Get $\approx -24.8^\circ$.


Problem: We wanted θ such that $0^\circ \leq \theta < 360^\circ$.

Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Putting $\tan^{-1}(-0.4623)$ in the calculator (in degree mode).

► Get $\approx -24.8^\circ$.

Problem: We wanted θ such that $0^\circ \leq \theta < 360^\circ$.




Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Putting $\tan^{-1}(-0.4623)$ in the calculator (in degree mode).

► Get $\approx -24.8^\circ$.

Problem: We wanted θ such that $0^\circ \leq \theta < 360^\circ$.




Solution: \tan is
 180° -periodic:

Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Putting $\tan^{-1}(-0.4623)$ in the calculator (in degree mode).

► Get $\approx -24.8^\circ$.

Problem: We wanted θ such that $0^\circ \leq \theta < 360^\circ$.



Solution: \tan is
 180° -periodic:


- Add 180° to -24.8°
 - $\theta = 155.2^\circ$

Find θ such that $\tan \theta = -0.4623$ and $0^\circ \leq \theta < 360^\circ$

Putting $\tan^{-1}(-0.4623)$ in the calculator (in degree mode).

► Get $\approx -24.8^\circ$.

Problem: We wanted θ such that $0^\circ \leq \theta < 360^\circ$.



Solution: \tan is
 180° -periodic:

- Add 180° to -24.8°
 - $\theta = 155.2^\circ$
- Add 180° to 155.2°
 - $\theta = 335.2^\circ$

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$


Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).

- Get ≈ 1.9651 .

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).


- ▶ Get ≈ 1.9651 .
- ▶ Since 1.9651 is between 0 and π , reference angle is $\approx \pi - 1.9651 \approx 1.1765$



Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).

- ▶ Get ≈ 1.9651 .
- ▶ Since 1.9651 is between 0 and π , reference angle is $\approx \pi - 1.9651 \approx 1.1765$




$\theta \approx 1.9651$ is a solution.

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).

- ▶ Get ≈ 1.9651 .
- ▶ Since 1.9651 is between 0 and π , reference angle is $\approx \pi - 1.9651 \approx 1.1765$




$\theta \approx 1.9651$ is a solution.

Find another θ with the same x value?

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).

- ▶ Get ≈ 1.9651 .
- ▶ Since 1.9651 is between 0 and π , reference angle is $\approx \pi - 1.9651 \approx 1.1765$




$\theta \approx 1.9651$ is a solution.

Find another θ with the same x value?

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).

- ▶ Get ≈ 1.9651 .
- ▶ Since 1.9651 is between 0 and π , reference angle is $\approx \pi - 1.9651 \approx 1.1765$




$\theta \approx 1.9651$ is a solution.

Find another θ with the same x value?

Find θ such that $\cos \theta = -0.3842$ and $0 \leq \theta < 2\pi$

Putting $\cos^{-1}(-0.3842)$ in the calculator (in radian mode).

- ▶ Get ≈ 1.9651 .
- ▶ Since 1.9651 is between 0 and π , reference angle is $\approx \pi - 1.9651 \approx 1.1765$




$\theta \approx 1.9651$ is a solution.

Find another θ with the same x value?

$\theta \approx \pi + 1.1765 \approx 4.3180$ is a second solution


Main idea

Use the symmetry in the circle with \pm to get \sin , \cos , \tan



Main idea

Use the symmetry in the circle with \pm to get \sin , \cos , \tan



The angles which have related x and y value have the same reference angle!