

IPMH: Integration of stepped care for perinatal mood and anxiety disorders among women attending MCH clinics

Data and Safety Monitoring Board (DSMB): Closed Report

13 May 2026



John Kinuthia, MBChD, MMed, MPH, Kenyatta National Hospital

Keshet Ronen, PhD, MPH, University of Washington

Amritha Bhat, MBBS, MD, MPH, University of Washington

Report prepared by: David Owaga & Yuwei Wang

Date of data freeze: 11 May 2026


Contents

1. Study Summary	4
2. Study Enrollment	5
2.1 Enrollment Progress - Overall	6
2.2 Enrollment Progress - Per Facility	7
3. Women Living with HIV Cohort	8
4. Ineligibility	9
5. Baseline Demographics	10
5.1 Participant Baseline Demographics	10
5.2 WLWH Cohort Baseline Demographics	11
6. Study Terminations	11
7. Referrals	11
7.1 Psychosocial Referrals	11
7.2 Intervention Treatment Referrals	12
8. Primary and secondary study endpoints at 14 weeks postpartum	13
9. Adverse Events	14
9.1 Serious Adverse Event Summary	14
9.2 Adverse Event Summary	15
9.2.1 List of Adverse Events: Intervention Arm	16
9.2.2 List of Adverse Events: Control Arm	18


1. Study Summary

Title	IPMH: Integration of stepped care for perinatal mood and anxiety disorders among women attending MCH clinics
Primary Objective	Integrate IPMH, a stepped care intervention for screening and treatment of perinatal mood and anxiety disorders (PMADs), and its implementation strategies, into maternal and child health clinics and pragmatically evaluate its clinical, service delivery, and implementation outcomes
Study Design	A 2-arm, non-blinded cluster randomized controlled trial (RCT) comparing the effect of IPMH and associated implementation strategies on depression, anxiety, quality of life, and adverse perinatal outcomes vs. control (standard of care) among Kenyan women
Study Population	Pregnant Kenyan women
Inclusion Criteria	-Receiving antenatal care at study facility ->=20 weeks gestation -Age>=14 -screen positive for PMAD symptoms (PHQ-2>=3 and/or GAD-2>=3)
Target Sample Size	2970, including 405 women living with HIV (WLWH)
Actual Enrollment	2383
Participants Completed Follow-up	966
Duration	Follow-up through 6 months postpartum
Date Study Open	September 19, 2023
Date of First Complete Enrollment	February 17, 2025
Estimated End of Study	July 31, 2028
Endpoints	Primary: Depression (Patient Health Questionnaire-9 [PHQ-9] score) and anxiety (Generalized Anxiety Disorder-7 [GAD-7] score) Secondary: -Problem Management Plus (PM+) Skill Acquisition (Reducing Tension Checklist score) -Quality of Life (World Health Organization Quality of Life [WHOQOL-BREF] score) -Any adverse pregnancy outcome

2. Study Enrollment




2.1 Enrollment Progress - Overall




2.2 Enrollment Progress - Per Facility

All 20 facilities have surpassed the 50% enrollment milestone, each enrolling at least 74 perinatal women. Progress has continued steadily, with 20 facilities surpassing the 75% milestone (≥ 111 enrollments), demonstrating strong momentum toward full recruitment.



3. Women Living with HIV Cohort

Overall 2383 participants have been enrolled into the study, of which 361 women are living with HIV (WLHIV), accounting to 15.1% of total enrollments. Of these, 215 (59.6%) have completed their 6 weeks, 170 (47.1%) have completed 14 weeks, and 118 (32.7%) have completed their 6 months postpartum follow-up visits.



4. Ineligibility

Women were ineligible if their gestational age at eligibility assessment was <20 weeks, if they endorsed thoughts/actions of self-harm, and if they endorsed symptoms of potential psychosis (e.g., hearing voices that others cannot hear, holding unusual beliefs, or feeling watched/followed), or if they endorsed experiencing memory problems.

Table 2: Summary of Study Ineligibility by Arm

Characteristic	Control N = 422	Intervention N = 419	Overall N = 841
Reasons for Ineligibility			
Gestation <20 Weeks	362 (86%)	409 (98%)	771 (92%)
Self harm	14 (3.3%)	7 (1.7%)	21 (2.5%)
Hearing voices that others cannot hear	15 (3.6%)	0 (0%)	15 (1.8%)
Memory problem	11 (2.6%)	2 (0.5%)	13 (1.5%)
Holding unusual beliefs	7 (1.7%)	0 (0%)	7 (0.8%)
Self harm and memory problem	7 (1.7%)	0 (0%)	7 (0.8%)
Feels watched/followed	5 (1.2%)	0 (0%)	5 (0.6%)
Seeing things that others cannot see	1 (0.2%)	1 (0.2%)	2 (0.2%)

¹ n (%)

5. Baseline Demographics

5.1 Participant Baseline Demographics

Demographic characteristics were balanced across study arms, supporting baseline comparability.

Table 3: Basic Demographic Summary

Characteristic	N	Control N = 1,201	Intervention N = 1,182	Overall N = 2,383
Age (Years)	2,381	26.0 [IQR: 21.0, 31.0]	25.0 [IQR: 21.0, 30.0]	26.0 [IQR: 21.0, 31.0]
Do you currently have a partner (Yes)	2,381	1,037 (86.4%)	983 (83.2%)	2,020 (84.8%)
Currently Married (Yes)	2,373	938 (78.6%)	882 (74.7%)	1,820 (76.7%)
Marriage (Monogamous)	1,820	831 (88.6%)	770 (87.3%)	1,601 (88.0%)
Shares residence with partner (Yes)	2,020	879 (84.8%)	785 (79.9%)	1,664 (82.4%)
Currently in School	2,381	96 (8.0%)	118 (10.0%)	214 (9.0%)
Completed years in School	2,381	11.0 [IQR: 8.0, 12.0]	12.0 [IQR: 10.0, 15.0]	12.0 [IQR: 9.0, 14.0]
Do you have regular employment (Yes)	2,381	150 (12.5%)	122 (10.3%)	272 (11.4%)
Number of people in Household	2,381	4.0 [IQR: 3.0, 5.0]	4.0 [IQR: 3.0, 5.0]	4.0 [IQR: 3.0, 5.0]
Number of people sleep in same house	2,381	3.0 [IQR: 3.0, 5.0]	3.0 [IQR: 2.0, 5.0]	3.0 [IQR: 2.0, 5.0]
Rooms in the house most often sleep in	2,381	2.0 [IQR: 1.0, 3.0]	2.0 [IQR: 1.0, 3.0]	2.0 [IQR: 1.0, 3.0]
Time to the Clinic (Minutes)	2,381	30.0 [IQR: 20.0, 40.0]	30.0 [IQR: 15.0, 40.0]	30.0 [IQR: 20.0, 40.0]
Have you been pregnant before (Yes)	2,371	901 (75.1%)	825 (70.5%)	1,726 (72.8%)

¹ Median [IQR: Q1, Q3]; n (%)

5.2 WLWH Cohort Baseline Demographics

Demographic characteristics were balanced across study arms, supporting baseline comparability.

Table 4: WLWH Demographic Summary

Characteristic	N	Control N = 204	Intervention N = 157	Overall N = 361
Age (Years)	361	30.0 [IQR: 25.0, 34.0]	30.0 [IQR: 25.0, 33.0]	30.0 [IQR: 25.0, 33.0]
Do you currently have a partner (Yes)	361	183 (90%)	130 (83%)	313 (87%)
Currently Married (Yes)	357	169 (85%)	121 (77%)	290 (81%)
Marriage (Monogamous)	290	146 (86%)	91 (75%)	237 (82%)
Shares residence with partner (Yes)	313	161 (88%)	103 (79%)	264 (84%)
Currently in School	361	9 (4.4%)	12 (7.6%)	21 (5.8%)
Completed years in School	361	8.0 [IQR: 8.0, 11.0]	11.0 [IQR: 8.0, 13.0]	10.0 [IQR: 8.0, 12.0]
Do you have regular employment (Yes)	361	10 (4.9%)	22 (14%)	32 (8.9%)
Number of people in Household	361	5.0 [IQR: 3.0, 6.0]	4.0 [IQR: 3.0, 5.0]	5.0 [IQR: 3.0, 5.0]
Number of people sleep in same house	361	4.0 [IQR: 3.0, 5.0]	4.0 [IQR: 3.0, 5.0]	4.0 [IQR: 3.0, 5.0]
Rooms in the house most often sleep in	361	2.0 [IQR: 1.0, 3.0]	2.0 [IQR: 2.0, 3.0]	2.0 [IQR: 1.0, 3.0]
Time to the Clinic (Minutes)	361	30.0 [IQR: 20.0, 60.0]	30.0 [IQR: 20.0, 50.0]	30.0 [IQR: 20.0, 60.0]
Have you been pregnant before (Yes)	351	190 (93%)	133 (90%)	323 (92%)

¹ Median [IQR: Q1, Q3]; n (%)

6. Study Terminations

To date, no early terminations have been recorded aside from maternal deaths (see adverse event section). A total of 966 participants have completed follow-up and exited the study at the scheduled endpoint, including 502 from the control arm and 464 from the intervention arm. The first termination occurred on 2025-08-18.

7. Referrals

7.1 Psychosocial Referrals

During enrollment and follow-up visits participants are asked if they would like to be referred for psychosocial support if they screen positive for depression, anxiety and intimate partner violence (IPV). Out of 6280 total study visits, 982 (15.6%) participants screened positive for referable conditions, of which, 897 (91.3%) have accepted referrals. These referrals reflect routine care and services available within the facility, not the implementation treatment

Referral Uptake by Study Arm

Condition	Control			Intervention		
	Screened +	Accepted	Uptake (%)	Screened +	Accepted	Uptake (%)
IPV Score $\geq$ 10	70	62	88.6	87	69	79.3
Anxiety Score $\geq$ 10	152	140	92.1	185	174	94.1
Depression Score $\geq$ 10	227	209	92.1	239	221	92.5
Self-harm	9	9	100.0	13	13	100.0

7.2 Intervention Treatment Referrals

During this study period, 21138 women attended MCH in the study clinics and were screened for PMAD symptoms, of whom 3313 (15.7%) screened positive for PMAD. Among these, 215(6.5%) had moderate symptoms in either PHQ9 or GAD7 and were referred to PM+, and 41 (1.2%) had severe symptoms and were referred to a psychiatrist.

Table 5: Stepped Care Intervention Table

Step	Intervention	Eligibility / Trigger	Delivered By	Duration/Frequency	Participants (n)
Step 1	PMAD Screening	All ANC Participants	Lay Health Worker	1 session (30-45 mins)	3313
Step 2	Problem Management Plus (PM+)	Moderate symptoms (10-14 Scores in either GAD-7 or PHQ-9)	Trained HC Worker (Nurse)	5 sessions (weekly)	215
Step 3	Telepsych / Specialist Referral	Severe symptoms ($\geq$ 15 in PHQ-9 or GAD7) or no response to PM+	Psychiatrist	As recommended	41

8. Primary and secondary study endpoints at 14 weeks postpartum

Table 6: Primary and Secondary Outcomes Summary by Arm

Characteristic	N	Control N = 668	Intervention N = 628	Overall N = 1,296
Patient Health Questionnaire-9 score	1,296	3.0 (0.0, 5.0)	3.0 (1.0, 6.0)	3.0 (0.0, 5.0)
Generalized Anxiety Disorder-7 score	1,296	2.0 (0.0, 5.0)	3.0 (0.0, 5.0)	3.0 (0.0, 5.0)
Quality of Life (WHOQOL BREF score)	1,275	62.5 (62.5, 75.0)	75.0 (62.5, 75.0)	75.0 (62.5, 75.0)
Any adverse pregnancy outcome	1,296	160 (24.0%)	173 (27.5%)	333 (25.7%)
Reducing Tensions Checklist score	1,296	17.0 (13.0, 22.0)	16.5 (9.0, 22.0)	17.0 (12.0, 22.0)

¹ Median (Q1, Q3); n (%)

9. Adverse Events

9.1 Serious Adverse Event Summary

Table 7: Summary of Reported Serious Adverse Events by Arm

Characteristic	Control N = 29	Intervention N = 25	Overall N = 54
Maternal Death	1 (1.9%)	1 (1.9%)	2 (3.7%)
Infant Death	18 (33.3%)	17 (31.5%)	35 (64.8%)
Infants Death Cause			
Respiratory complications	9 (25.7%)	9 (25.7%)	18 (51.4%)
Fever/Vomiting	1 (2.9%)	3 (8.6%)	4 (11.4%)
Other / Unknown	3 (8.6%)	0 (0.0%)	3 (8.6%)
Birth asphyxia / Intrapartum hypoxia	2 (5.7%)	0 (0.0%)	2 (5.7%)
Birth asphyxia secondary to prolonged labor	1 (2.9%)	1 (2.9%)	2 (5.7%)
Prematurity	0 (0.0%)	2 (5.7%)	2 (5.7%)
Respiratory failure / Aspiration	1 (2.9%)	1 (2.9%)	2 (5.7%)
Congenital malformations / Birth defects	1 (2.9%)	0 (0.0%)	1 (2.9%)
Still birth	0 (0.0%)	1 (2.9%)	1 (2.9%)
Maternal Death Cause			
Gastrointestinal bleeding	1 (50.0%)	0 (0.0%)	1 (50.0%)
Intraoperative complication during CS	0 (0.0%)	1 (50.0%)	1 (50.0%)
Stillbirth	9 (16.7%)	6 (11.1%)	15 (27.8%)
New/prolonged hospitalization	1 (1.9%)	1 (1.9%)	2 (3.7%)

¹ n (%)

9.2 Adverse Event Summary

Table 8: Summary of Adverse Events by Arm

Characteristic	Control N = 22	Intervention N = 12	Overall N = 34
Adverse Event			
<i>Persistent or significant psychosocial distress</i>	12 (55%)	2 (17%)	14 (41%)
<i>Self-harm/Suicidal behavior</i>	3 (14%)	6 (50%)	9 (26%)
<i>Experienced violence or abuse</i>	4 (18%)	3 (25%)	7 (21%)
<i>Kicked out of home</i>	3 (14%)	0 (0%)	3 (8.8%)
<i>Injury</i>	0 (0%)	1 (8.3%)	1 (2.9%)

¹ Counts reflect events, not unique participants. Persistent or significant psychosocial distress is defined as PHQ-9 score ≥ 10 at any follow-up visit.

9.2.1 List of Adverse Events: Intervention Arm

Table 9: List of Study Adverse Events for Arm X

PTID	Study Site	Adverse Event reported	Age of participant	Days from Onset to Report	Days from enrolment to AE report
21198187	Airport Health Centre (Kisumu)	Maternal Death (SAE)	34	25	43
21031003	Uyawi Sub County Hospital	Infant Death (SAE)	17	11	12
21031015	Uyawi Sub County Hospital	Infant Death (SAE)	30	0	15
21041543	Got Agulu Sub-District Hospital	Infant Death (SAE)	21	20	55
21041553	Got Agulu Sub-District Hospital	Infant Death (SAE)	23	12	34
21052094	Ukwala Sub County Hospital	Infant Death (SAE)	33	40	120
21052106	Ukwala Sub County Hospital	Infant Death (SAE)	27	41	147
21062515	Madiany Sub County Hospital	Infant Death (SAE)	26	44	85
21073090	Kabondo Sub County Hospital	Infant Death (SAE)	33	8	45
21073127	Kabondo Sub County Hospital	Infant Death (SAE)	32	57	78
21094059	Miriu Health Centre	Infant Death (SAE)	19	50	91
21145531	Gita Sub County Hospital	Infant Death (SAE)	20	48	112
21156092	Akala Health Centre	Infant Death (SAE)	24	11	49
21156107	Akala Health Centre	Infant Death (SAE)	23	36	69
21166542	Usigu Health Centre	Infant Death (SAE)	35	38	168
21166559	Usigu Health Centre	Infant Death (SAE)	31	57	197
21177015	Ramula Health Centre	Infant Death (SAE)	18	0	168
21187505	Simenya Health Centre	Infant Death (SAE)	24	13	63
21187522	Simenya Health Centre	Infant Death (SAE)	23	14	75
21187627	Simenya Health Centre	Infant Death (SAE)	21	18	70
21187630	Simenya Health Centre	Infant Death (SAE)	24	20	24
21187637	Simenya Health Centre	Infant Death (SAE)	23	20	63
21198037	Airport Health Centre (Kisumu)	Infant Death (SAE)	18	8	39
21198045	Airport Health Centre (Kisumu)	Infant Death (SAE)	22	11	50
21198138	Airport Health Centre (Kisumu)	Infant Death (SAE)	26	16	0
21208555	Nyalenda Health Centre	Infant Death (SAE)	21	20	39
21208581	Nyalenda Health Centre	Infant Death (SAE)	41	56	85
21208621	Nyalenda Health Centre	Infant Death (SAE)	25	33	36
21229609	Ndiru Level 4 Hospital	Infant Death (SAE)	16	1	35
21229621	Ndiru Level 4 Hospital	Infant Death (SAE)	15	49	106

21020522	Sigomere Sub County Hospital	Stillbirth (SAE)	20	4	28
21020575	Sigomere Sub County Hospital	Stillbirth (SAE)	26	29	128
21041565	Got Agulu Sub-District Hospital	Stillbirth (SAE)	26	46	98
21052061	Ukwala Sub County Hospital	Stillbirth (SAE)	35	6	33
21052089	Ukwala Sub County Hospital	Stillbirth (SAE)	32	2	43
21073011	Kabondo Sub County Hospital	Stillbirth (SAE)	26	2	47
21145514	Gita Sub County Hospital	Stillbirth (SAE)	39	8	132
21166529	Usigu Health Centre	Stillbirth (SAE)	27	2	9
21177121	Ramula Health Centre	Stillbirth (SAE)	22	5	6
21177126	Ramula Health Centre	Stillbirth (SAE)	22	7	0
21187649	Simenya Health Centre	Stillbirth (SAE)	20	12	35
21198003	Airport Health Centre (Kisumu)	Stillbirth (SAE)	28	89	98
21208607	Nyalenda Health Centre	Stillbirth (SAE)	31	25	130
21219053	Mirogi Health Centre	Stillbirth (SAE)	24	36	49
21020565	Sigomere Sub County Hospital	New/prolonged hospitalization (SAE)	31	4	148
21073028	Kabondo Sub County Hospital	New/prolonged hospitalization (SAE)	23	0	51
21041564	Got Agulu Sub-District Hospital	Self-harm/Suicidal behavior	29	28	132
21229560	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	15	7	97
21229601	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	17	12	86
21229611	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	27	7	157
21229642	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	23	7	95
21041564	Got Agulu Sub-District Hospital	Persistent or significant psychosocial distress	29	28	132
21052106	Ukwala Sub County Hospital	Persistent or significant psychosocial distress	27	73	147
21156108	Akala Health Centre	Persistent or significant psychosocial distress	38	17	151
21156116	Akala Health Centre	Persistent or significant psychosocial distress	37	10	110
21229529	Ndiru Level 4 Hospital	Persistent or significant psychosocial distress	21	38	57
21156116	Akala Health Centre	Experienced violence or abuse	37	10	110
21229529	Ndiru Level 4 Hospital	Experienced violence or abuse	21	38	57
21156116	Akala Health Centre	Kicked out of home	37	10	110
21010120	Rwambwa Sub-county Hospital	Injury	35	52	54

9.2.2 List of Adverse Events: Control Arm

Table 10: List of Study Adverse Events for Arm Y

PTID	Study Site	Adverse Event reported	Age of participant	Days from Onset to Report	Days from enrolment to AE report
21198187	Airport Health Centre (Kisumu)	Maternal Death (SAE)	34	25	43
21031003	Uyawi Sub County Hospital	Infant Death (SAE)	17	11	12
21031015	Uyawi Sub County Hospital	Infant Death (SAE)	30	0	15
21041543	Got Agulu Sub-District Hospital	Infant Death (SAE)	21	20	55
21041553	Got Agulu Sub-District Hospital	Infant Death (SAE)	23	12	34
21052094	Ukwala Sub County Hospital	Infant Death (SAE)	33	40	120
21052106	Ukwala Sub County Hospital	Infant Death (SAE)	27	41	147
21062515	Madiany Sub County Hospital	Infant Death (SAE)	26	44	85
21073090	Kabondo Sub County Hospital	Infant Death (SAE)	33	8	45
21073127	Kabondo Sub County Hospital	Infant Death (SAE)	32	57	78
21094059	Miriu Health Centre	Infant Death (SAE)	19	50	91
21145531	Gita Sub County Hospital	Infant Death (SAE)	20	48	112
21156092	Akala Health Centre	Infant Death (SAE)	24	11	49
21156107	Akala Health Centre	Infant Death (SAE)	23	36	69
21166542	Usigu Health Centre	Infant Death (SAE)	35	38	168
21166559	Usigu Health Centre	Infant Death (SAE)	31	57	197
21177015	Ramula Health Centre	Infant Death (SAE)	18	0	168
21187505	Simenya Health Centre	Infant Death (SAE)	24	13	63
21187522	Simenya Health Centre	Infant Death (SAE)	23	14	75
21187627	Simenya Health Centre	Infant Death (SAE)	21	18	70
21187630	Simenya Health Centre	Infant Death (SAE)	24	20	24
21187637	Simenya Health Centre	Infant Death (SAE)	23	20	63
21198037	Airport Health Centre (Kisumu)	Infant Death (SAE)	18	8	39
21198045	Airport Health Centre (Kisumu)	Infant Death (SAE)	22	11	50
21198138	Airport Health Centre (Kisumu)	Infant Death (SAE)	26	16	0
21208555	Nyalenda Health Centre	Infant Death (SAE)	21	20	39
21208581	Nyalenda Health Centre	Infant Death (SAE)	41	56	85
21208621	Nyalenda Health Centre	Infant Death (SAE)	25	33	36
21229609	Ndiru Level 4 Hospital	Infant Death (SAE)	16	1	35
21229621	Ndiru Level 4 Hospital	Infant Death (SAE)	15	49	106

21020522	Sigomere Sub County Hospital	Stillbirth (SAE)	20	4	28
21020575	Sigomere Sub County Hospital	Stillbirth (SAE)	26	29	128
21041565	Got Agulu Sub-District Hospital	Stillbirth (SAE)	26	46	98
21052061	Ukwala Sub County Hospital	Stillbirth (SAE)	35	6	33
21052089	Ukwala Sub County Hospital	Stillbirth (SAE)	32	2	43
21073011	Kabondo Sub County Hospital	Stillbirth (SAE)	26	2	47
21145514	Gita Sub County Hospital	Stillbirth (SAE)	39	8	132
21166529	Usigu Health Centre	Stillbirth (SAE)	27	2	9
21177121	Ramula Health Centre	Stillbirth (SAE)	22	5	6
21177126	Ramula Health Centre	Stillbirth (SAE)	22	7	0
21187649	Simenya Health Centre	Stillbirth (SAE)	20	12	35
21198003	Airport Health Centre (Kisumu)	Stillbirth (SAE)	28	89	98
21208607	Nyalenda Health Centre	Stillbirth (SAE)	31	25	130
21219053	Mirogi Health Centre	Stillbirth (SAE)	24	36	49
21020565	Sigomere Sub County Hospital	New/prolonged hospitalization (SAE)	31	4	148
21073028	Kabondo Sub County Hospital	New/prolonged hospitalization (SAE)	23	0	51
21041564	Got Agulu Sub-District Hospital	Self-harm/Suicidal behavior	29	28	132
21229560	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	15	7	97
21229601	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	17	12	86
21229611	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	27	7	157
21229642	Ndiru Level 4 Hospital	Self-harm/Suicidal behavior	23	7	95
21041564	Got Agulu Sub-District Hospital	Persistent or significant psychosocial distress	29	28	132
21052106	Ukwala Sub County Hospital	Persistent or significant psychosocial distress	27	73	147
21156108	Akala Health Centre	Persistent or significant psychosocial distress	38	17	151
21156116	Akala Health Centre	Persistent or significant psychosocial distress	37	10	110
21229529	Ndiru Level 4 Hospital	Persistent or significant psychosocial distress	21	38	57
21156116	Akala Health Centre	Experienced violence or abuse	37	10	110
21229529	Ndiru Level 4 Hospital	Experienced violence or abuse	21	38	57
21156116	Akala Health Centre	Kicked out of home	37	10	110
21010120	Rwambwa Sub-county Hospital	Injury	35	52	54