

Relaxations of the Maximum Cut Problem

Demetrios V. Papazaharias, Carter Mann, Luca Wrabetz

Department of Industrial and Systems Engineering

University at Buffalo, Bell Hall, Buffalo, New York, 14260

{dvpapaza, cjmann3, lucawrab}@buffalo.edu

December 2019

Abstract

Maximum cut is a classic NP-Hard problem in combinatorial optimization. In this work we present several relaxations for the maximum cut problem.

1 Introduction

2 Mathematical Formulations

3 Relaxations for Max Cut

3.1 Eigenvalue Relaxation

Proposition 1. *The optimal value of z^* in the maxcut problem defined on graph $G = (V, E)$ satisfies*

$$z^* \leq z^{ev} := \frac{n}{4} \lambda_{\max}(L(G))$$

Proof. Consider the maximization problem

$$\max\{\mathbf{z}^\top L(G) \mathbf{z} \mid \mathbf{z} \in \{-1, 1\}^n\}$$

Since $L(G) = D(G) - A(G)$, then $L(G)_{ii} = \sum_{j \in V} w_{ij}$ for $i \in V$ and $L(G)_{ij} = -w_{ij}$ for $i, j \in V$ such that $i \neq j$. We expand the objective function and obtain

$$\max_{\mathbf{z} \in \{-1, 1\}^n} \sum_{i \in V} \sum_{j \in V} w_{ij} z_i z_j - \sum_{i \in V} \sum_{j \in V} w_{ij} z_i z_j$$

Furthermore,

$$\max_{\mathbf{z} \in \{-1, 1\}^n} \sum_{i \in V} \sum_{j \in V} w_{ij} (z_i z_i - z_i z_j)$$

Since $z_i \in \{-1, 1\}$ then $z_i z_i = 1$, we can simplify the objective function to

$$\max_{\mathbf{z} \in \{-1, 1\}^n} \sum_{i \in V} \sum_{j \in V} w_{ij} (1 - z_i z_j)$$

From the previous section we saw an almost identical formulation for max cut. The only difference being a multiplier of $\frac{1}{4}$. Let z^* represent the optimal solution for max cut

$$z^* = \frac{1}{4} \max\{\mathbf{z}^\top L(G) \mathbf{z} \mid \mathbf{z} \in \{-1, 1\}^n\} \quad (1)$$

Taking the continuous relaxation of (1), namely $z_i \in [-1, 1]$ for $i \in V$, is equivalent to maximizing over the norm infinity, $\|\mathbf{z}\|_\infty \leq 1$. We can relax this further by maximizing over a ball of radius \sqrt{n} . In other words, our region is now defined where $\|\mathbf{z}\| \leq \sqrt{n}$.

$$z^* \leq \frac{1}{4} \max\{\mathbf{z}^\top L(G) \mathbf{z} \mid \mathbf{z} \leq \sqrt{n}\} \quad (2)$$

We can define our problem over the unit ball with a simple transformation. Let $\|\mathbf{z}\| = \sqrt{n} \mathbf{x}$ and we now have

$$z^* \leq \frac{n}{4} \max\{\mathbf{x}^\top L(G) \mathbf{x} \mid \|\mathbf{x}\| \leq 1\} \quad (3)$$

Since for any symmetric matrix \mathbf{A} , $\max\{\mathbf{x}^\top \mathbf{A} \mathbf{x} \mid \|\mathbf{x}\| \leq 1\} = \lambda_{\max}(\mathbf{A})$ and $L(G)$ is symmetric, then we have

$$z^* \leq z^{ev} := \frac{n}{4} \lambda_{\max}(L(G)) \quad (4)$$

□

Proposition 2. *The optimal value of z^* in the maxcut problem defined on graph $G = (V, E)$ satisfies*

$$z^* \leq z^{ev} := -\frac{1}{4} \sum_{i=1}^n u_i + \frac{n}{4} \lambda_{\max}(L(G) + \text{diag}(\mathbf{u}))$$

for all $\mathbf{u} \in \mathbb{R}^n$

Proof. Consider the maximization problem

$$\max\{\mathbf{z}^\top (L(G) + \text{diag}(\mathbf{u}))\mathbf{z} \mid \mathbf{z} \in \{-1, 1\}\} \quad (5)$$

Note that no matter the value of $\mathbf{u} \in \mathbb{R}^n$, $L(G) + \text{diag}(\mathbf{u})$ remains to be a symmetric square matrix. As a result,

$$\lambda_{\max}(L(G) + \text{diag}(\mathbf{u})) = \max\{\mathbf{z}^\top (L(G) + \text{diag}(\mathbf{u}))\mathbf{z} \mid \mathbf{z} \in \{-1, 1\}\}$$

From the previous proof, we perform a similar technique to (5) to obtain the following optimization problem

$$\max_{\mathbf{z} \in \{-1, 1\}} \sum_{i \in V} \left(\sum_{j \in V} w_{ij} + u_i \right) z_i z_i - \sum_{i \in V} \sum_{j \in V} w_{ij} z_i z_j$$

Rearranging terms and using the fact that $z_i z_i = 1$,

$$\max_{\mathbf{z} \in \{-1, 1\}} \sum_{i=1}^n \sum_{j=1}^n w_{ij} (1 - z_i z_j) + \sum_{i=1}^n u_i \quad (6)$$

Which is closely related to the objective function in our previous formulation. We obtain the following relationship between (6) and z^* :

$$z^* = \frac{1}{4} \max\{\mathbf{z}^\top (L(G) + \text{diag}(\mathbf{u}))\mathbf{z} \mid \mathbf{z} \in \{-1, 1\}\} - \frac{1}{4} \sum_{i=1}^n u_i$$

We repeat the process from the previous proof of relaxing the variables so that they are continuous and then further relaxing this problem onto a ball of radius \sqrt{n} .

$$z^* \leq \frac{1}{4} \max\{\mathbf{z}^\top (L(G) + \text{diag}(\mathbf{u}))\mathbf{z} \mid \|\mathbf{z}\| \leq \sqrt{n}\} - \frac{1}{4} \sum_{i=1}^n u_i$$

Let $\mathbf{z} = \sqrt{n}\mathbf{x}$.

$$z^* \leq \frac{n}{4} \max\{\mathbf{x}^\top (L(G) + \text{diag}(\mathbf{u}))\mathbf{x} \mid \|\mathbf{x}\| \leq 1\} - \frac{1}{4} \sum_{i=1}^n u_i$$

Finally, we obtain the inequality

$$z^* \leq z^{ev}(\mathbf{u}) = -\sum_{i=1}^n u_i + \frac{n}{4} \lambda_{\max}(L(G) + \text{diag}(\mathbf{u})) \quad (7)$$

□