Measurement of CP Violation in  $B^{\pm} \to \pi^{\pm}\pi^{+}\pi^{-}$  Decay Channal at Large Hadron Collider

Qichen Dong, Harriet Watson

School of Physics and Astronomy, University of Manchester, Manchester, M13 9PL.

**Abstract:** A set of selected data samples which were collected by LHCb[1] in 2011 are studied. Contained  $B^{\pm} \to \pi^{\pm}\pi^{+}\pi^{-}$  decays in magnet "up" and "down" polarity are constructed. Global CP asymmetry in this channal is measured to be  $A_{CP} = 0.135 \pm 0.023 \pm 0.010$ , in which the first and second uncertainties are statistical and systematical respectively.

## 1 Introduction

Standard Model[2] of particle physics, which developed in early 1970s, has been successfully explaining almost all experiment results and making precise preditions including

## 2 Theory

meaningless texts meaningless texts




Figure 1: this is a figure demo

meaningless texts meaningless texts

## 3 Conclusions

meaningless texts meaningless texts

#### 4 Results

meaningless texts meaningless texts

meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts meaningless texts




Figure 2: this is a figure demo

meaningless texts meaningless texts

Start	Character Block Name
3400	graphs Extension A
4E00	CJK Unified Ideographs

Table 1: asdsada

# **Bibliography**

## References

- [1] A. A. Alves Jr. et al. The lhcb detector at the lhc. JINST 3 (2008) S08005.
- [2] Tom W.B. Kibble. The standard model of particle physics. arXiv:1412.4094.