Measurement of CP Violation in  $B^{\pm} \to \pi^{\pm}\pi^{+}\pi^{-}$  Decay Channal at Large Hadron Collider Qichen Dong, Harriet Watson

School of Physics and Astronomy, University of Manchester, Manchester, M13 9PL.

**Abstract:** A set of selected data samples which were collected by LHCb[1] in 2011 are studied. Contained  $B^{\pm} \to \pi^{\pm}\pi^{+}\pi^{-}$  decays in magnet "up" and "down" polarity are constructed. Global CP asymmetry in this channal is measured to be  $A_{CP} = 0.135 \pm 0.023 \pm 0.010$ , in which the first and second uncertainties are statistical and systematical respectively.

## 1 Introduction

meaningless texts meaningless texts

# 2 Theory

meaningless texts meaningless texts




Figure 1: this is a figure demo

meaningless texts meaningless texts

# 3 Conclusions

meaningless texts meaningless texts

### 4 Results

meaningless texts meaningless texts




Figure 2: this is a figure demo

 $M^2_{\pi^+\pi^-low}\;(GeV^2)$ 

Start	Character Block Name
3400	graphs Extension A
4E00	CJK Unified Ideographs

Table 1: asdsada

meaningless texts meaningless texts

### 5 Ref

#### References

[1] A. A. Alves Jr. et al. The lhcb detector at the lhc,. JINST 3 (2008) S08005.