

BMJ Open Respiratory Research

Association between air pollutant exposure, body water distribution and sleep disorder indices in individuals with low-arousal-threshold obstructive sleep apnoea

Cheng-Yu Tsai,^{1,2} Ming Liu,³ Huei-Tyng Huang,⁴ Wen-Hua Hsu,⁵ Yi-Chun Kuan,^{6,7,8,9} Arnab Majumdar,¹ Kang-Yun Lee,^{2,10} Po-Hao Feng,^{2,10} Chien-Hua Tseng,^{2,10} Kuan-Yuan Chen,² Jiunn-Horng Kang,^{11,12,13} Hsin-Chien Lee,¹⁴ Cheng-Jung Wu,¹⁵ Wen-Te Liu (1) 2,5,6,11

To cite: Tsai C-Y, Liu M, Huang H-T, et al. Association between air pollutant exposure, body water distribution and sleep disorder indices in individuals with low-arousalthreshold obstructive sleep apnoea. BMJ Open Respir Res 2023;10:e001802. doi:10.1136/ bmjresp-2023-001802

Additional supplemental material is published online only. To view, please visit the iournal online (http://dx.doi. org/10.1136/bmjresp-2023-001802).

Received 30 April 2023 Accepted 27 October 2023



@ Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by

For numbered affiliations see end of article.

Correspondence to Dr Wen-Te Liu; lion5835@gmail.com

BMJ

ABSTRACT

Background Air pollution may alter body water distribution, it may also be linked to low-arousal-threshold obstructive sleep apnoea (low-ArTH OSA). Here, we explored the mediation effects of air pollution on body water distribution and low-ArTH OSA manifestations. **Methods** In this retrospective study, we obtained sleep centre data from healthy participants and patients with low-ArTH OSA (N=1924) in northern Taiwan. Air pollutant exposure at different time intervals (1, 3, 6 and 12 months) was estimated using the nearest station estimation method, and government air-quality data were also obtained. Regression models were used to assess the associations of estimated exposure, sleep disorder indices and body water distribution with the risk of low-ArTH OSA. Mediation analysis was performed to explore the relationships between air pollution, body water distribution and sleep disorder indices.

Results First, exposure to particulate matter (PM) with a diameter of ≤10 µm (PM₁₀) for 1 and 3 months and exposure to PM with a diameter of ≤2.5 µm (PM_{2.5}) for 3 months were significantly associated with the Apnoea-Hypopnoea Index (AHI), Oxygen Desaturation Index (ODI), Arousal Index (Arl) and intracellular-to-extracellular water ratio (I-E water ratio). Significant associations were observed between the risk of low-ArTH OSA and 1-month exposure to PM₁₀ (OR 1.42, 95% CI 1.09 to 1.84), PM₂ (OR 1.33, 95% Cl 1.02 to 1.74) and ozone (OR 1.27, 95% Cl 1.01 to 1.6). I-E water ratio alternation caused by 1-month exposure to PM₁₀ and 3-month exposure to PM₂₅ and PM₁₀ had partial mediation effects on AHI and ODI.

Conclusion Air pollution can directly increase sleep disorder indices (AHI, ODI and ArI) and alter body water distribution, thus mediating the risk of low-ArTH OSA.

INTRODUCTION

Obstructive sleep apnoea (OSA) is a sleeprelated breathing disorder affecting approximately 1 billion people globally.¹ disorder is characterised by the partial or

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Exposure to air pollution may lead to alterations in body water distribution and exacerbate the sleep disorder indices (ie, Apnoea-Hypopnoea Index, Oxygen Desaturation Index and Arousal Index).

WHAT THIS STUDY ADDS

 \Rightarrow Air pollution (particulate matter (PM₁₀) and PM₂₅ exposure) worsens obstructive sleep apnoea (OSA) severity and alters body water distribution, affecting sleep disorder indices. One-month PM_{10} , PM_{25} and ozone exposure mediates the risk of low-arousalthreshold (low-ArTH) OSA.

HOW THIS STUDY MIGHT AFFECT RESEARCH. PRACTICE OR POLICY

⇒ Reducing exposure to air pollutants may reduce sleep disorder indices, thus decreasing the risk of low-ArTH OSA.

complete obstruction of the upper airway during sleep, which reduces oxygen delivery.² In approximately one-third of patients, OSA disrupts sleep continuity and limits the accumulation of respiratory stimuli that are required to restore upper airway patency and airflow during sleep.³ This propensity to be woken prematurely in response to respiratory events is defined as low-arousal-threshold (low-ArTH) OSA. Several risk factors for low-ArTH OSA have been identified, including ageing,⁵ body water distribution⁶ and air pollutant exposure. However, whether these factors exert a synergistic effect on low-ArTH OSA manifestations requires further investi-

OSA manifestations are evaluated through polysomnography (PSG), followed by the





measurement of sleep disorder indices-namely the Apnoea-Hypopnoea Index (AHI), Oxygen Desaturation Index (ODI) and Arousal Index (ArI). A study suggested that when screening for the risk of low-ArTH OSA, patients with OSA should fulfil at least two of these three criteria: (1) AHI<30 events/hour, (2) lowest oxygen saturation values >82.5\% (measured through pulse oximetry) and (3) proportion of hypopnoea (ie, hypopnoea-to-total respiratory event) >58.3%. Notably, air pollution affects the AHI, ODI, ArI and ArTH. For instance, in a review of 15 studies involving approximately 133 000 participants from 10 countries, the authors noted that air pollutants such as particulate matter (PM) with a diameter of $\leq 2.5 \,\mu\text{m}$ (PM_{9.5}) and $10 \,\mu\text{m}$ (PM₁₀), nitrogen oxide (NO) and nitrogen dioxide (NO₉) might be associated with increased mean AHI, ODI and ArI values, thereby resulting in sleep quality impairment. A related study on the effects of air pollution on the central nervous system reported that air pollutant exposure may increase sleep arousal frequency, and that air pollutant exposure may be associated with low-ArTH OSA. Moreover, patients with low-ArTH OSA tend to exhibit sleep arousal due to respiratory disturbances, particularly minor events such as hypopnoea rather than apnoea. 10 Several mechanisms associated with this manifestation of low-ArTH OSA, which is partially linked to air pollutant exposure, have been identified. For example, the brain circuitry involved in neural substrates and glutamatergic signalling potentially causes sleep arousal.¹¹ An animal study suggested that exposure to PM_{9.5} interferes with the circuitry of the brain and alters its ultrastructure. 12 Moreover, neuroinflammation and increased oxidative stress in the brain may interfere with the sleep-wake cycle by inducing sleep arousal, and these physiological responses are exacerbated by air pollution. ¹³ PM and NO₂ may affect the central nervous system by altering neurotransmitter levels and breaking down protective epithelial barriers, thus disrupting sleep-regulated brain functions. 15 16 Hence, air pollutant exposure may be a risk factor for the worsening of sleep disorder indices and low-ArTH OSA; thus, further relevant investigation is required.

Air pollutant exposure may affect the balance or fluctuation of body water distribution, namely intracellular water (ICW) and extracellular water (ECW), and may thus worsen sleep disorder indices. Chronic air pollutant exposure may trigger airway mucosa inflammation and osmotic pressure changes, leading to inflammation and oedema of the proximal upper airways and thus changing the body water distribution balance and increasing sleep disorder indices. ^{17 18} In addition, fine particles have been reported to trigger an oxidative mechanism that increases the oxidative stress of cells and alters the water balance between intracellular and extracellular compartments. ¹⁹ Consequently, interactions may exist between air pollutants, body water distribution and OSA severity, but their relationships have yet to be comprehensively explored.

Considering the necessity to further examine the associations between air pollution, body water distribution and




Figure 1 Workflow of data acquisition and statistical analysis of this study. This study first retrospectively accessed the PSG database to obtain the addresses, anthropometric data, body composition data and sleep parameters of patients at a sleep centre. Next, based on the obtained addresses and data from government air monitoring stations, air pollutant exposure levels were estimated. The retrieved data were subsequently subjected to further association and mediation analyses. CO, carbon oxide; NO₂, nitrogen dioxide; NO, nitrogen oxide; O₃, ozone; PM₁₀, aerodynamic diameter of <10 μm; PM_{2.5}, aerodynamic diameter of <2.5 μm; PSG, polysomnography; SO₂, sulfur dioxide (SO₃).

sleep disorder indices in low-ArTH OSA, the objective of this retrospective study was to investigate the relationships among these variables. This study hypothesised that air pollutant exposure influences sleep disorder indices directly and modifies body water distribution indirectly, thereby leading to increases in the AHI, ODI and ArI. To investigate this hypothesis, we collected and analysed the data of both healthy individuals and patients with low-ArTH OSA from a sleep centre in northern Taiwan. Information regarding their residential address, body composition, body water distribution and PSG parameters was collected for examination. Using the obtained addresses, exposure volumes to various air pollutants were estimated, and the relationships between air pollution and the aforementioned variables were investigated. The potential influence of air pollution on the risk of low-ArTH OSA was also evaluated.

METHODS

Study population

Figure 1 presents the workflow of data acquisition and statistical analysis in this study. First, this retrospective study analysed patient data between June 2019 and December 2021 from the PSG database of the sleep centre of Shuang Ho Hospital (New Taipei City, Taiwan). Participants were included in the analysis if they (1) had

Table 1 Demographic characteristics and body composition details of the sample (N=1924)

| Variable | Mean±SD | Variable | N (%) |
|--------------------------|--------------|--------------|--------------|
| Age (year) | 46.8±13.63 | Sex | |
| BMI (kg/m²) | 26.13±4.72 | Male | 1170 (60.81) |
| Neck circumference (cm) | 36.8±3.92 | Female | 754 (39.19) |
| Waist circumference (cm) | 89.62±12.19 | OSA severity | |
| Body composition | | Normal | 326 (16.94) |
| Visceral fat level | 10.85±4.86 | Mild | 541 (28.12) |
| Fat proportion (%) | 28.42±8.88 | Moderate | 431 (22.4) |
| Muscle proportion (%) | 17.81±4.63 | Severe | 626 (32.54%) |
| TBW (%) | 49.9±5.56 | | |
| Body water distribution | | | |
| ICW (%) | 58.46±2.61 | | |
| ECW (%) | 41.54±2.61 | | |
| I-E water ratio | 141.65±14.95 | | |

ICW (%)=(ICW (kg)/TBW (kg))×100.

ECW (%)=(ECW (kg)/TBW (kg))×100.

I-E water ratio=(ICW/ECW)×100.

BMI, body mass index; ECW, extracellular water; ICW, intracellular water; I-E water ratio, ratio of ICW to ECW; OSA, obstructive sleep apnoea; TBW, total body water.

complete data on PSG parameters and a total recording time >6 hours, (2) were aged from 18 to 85 years, (3) had not undergone any invasive surgery or non-invasive treatment for OSA, (4) had not regularly used hypnotic or psychotropic medications, (5) had not been diagnosed as having central nervous system disorders (eg, brain tumour, stroke or major head trauma) and (6) exhibited normal AHI (<5 events/hour) or satisfied the screening criteria for low-ArTH OSA. The residential address and physical profiles (which included information on age, sex, body mass index (BMI), and neck and waist circumferences) were obtained from the clinical records of all eligible individuals.

Body composition and body water distribution

Data on body composition and body water distribution were obtained from the aforementioned medical database. Data determination was conducted using the Tanita MC-780 system (Tanita, Tokyo, Japan), and the procedures were as follows. Individuals were instructed to undergo data determination (the measurement of bioelectrical impedance) before PSG. All participants were required to fast for at least 3 hours and empty their bladders before undergoing bioelectrical impedance measurement. During data collection, patients were instructed to stand still and shoulder-width apart while holding the induction metal handles with both arms placed straight down. Next, the visceral fat level and percentages of fat, muscle and water in the whole-body scale were obtained automatically. The ECW and ICW percentages and ICWto-ECW ratio (I-E water ratio: ICW/ECW×100) were calculated.

Sleep profile-PSG parameter

Three recording systems were used to perform PSG at the sleep centre of Shuang Ho Hospital, including the Embla N7000 (ResMed, San Diego, California, USA), Embletta MPR (Natus Medical, Pleasanton, California, USA), and Nox-A1 (Nox Medical, Alpharetta, Georgia, USA). Two types of scoring interface, including RemLogic software (V.3.41; Embla Systems, Thornton, Colorado, USA) and Noxturnal system (V.6.2.2; Nox Medical), were employed corresponding to the various PSG hardware systems. The procedures for scoring physiological signals in PSG were all conducted by a certificated PSG technologist in accordance with the American Academy of Sleep Medicine manual published in 2017. 20 Specifically, apnoea (≥90%) reduction in the signals of an oronasal thermistor), hypopnoea (≥30% reduction in the signals of nasal prong pressure combined with ≥3% oxygen desaturation or with occurring arousal) and the frequency of oxygen desaturation (≥3% oxygen desaturation) were scored. The PSG technologists scored intervals that comprised alterations in brainwave signals (≥3s) with high-frequency patterns that did not spindle (alpha wave: 8–12Hz, theta wave: 4–8 Hz, high-frequency: >16 Hz) and that were preceded by stable sleep ($\geq 10 \, \text{s}$) as arousal events. Next, the AHI and the ArI were calculated accordingly, and OSA severity was defined as follows: none, AHI<5 events/hour; mild, AHI: 5–15 events/hour; moderate, AHI=15–30 events/hour and severe, AHI≥30 events/hour.²¹ To reduce the individual scoring bias, another technologist was requested to examine the scoring outcomes independently, and the varying parts were extracted for discussions to reach a scoring consensus.

Table 2 Sleep parameters on polysomnography of all participants (N=1924)

| Categorical variable | Mean±SD |
|-----------------------------------|---------------|
| Sleep architecture | |
| Sleep efficiency (%) | 76.76±12.8 |
| Wake (% of SPT) | 16.57±11.53 |
| NREM (% of SPT) | 71.3±10.27 |
| REM (% of SPT) | 12.1±6.1 |
| WASO (min) | 55.85±39.49 |
| TST (min) | 281.32±47.54 |
| Sleep quality index (events/hour) | |
| ODI | 18.43±20.61 |
| Snoring index | 198.35±220.59 |
| Arl | 18.08±10.94 |
| Low-ArTH criteria | |
| AHI (events/hour) | 23.88±21.04 |
| minSpO ₂ (%) | 86.16±7.25 |
| F-hypopnoea (%) | 89.07±13.41 |

ODI=the cumulative frequency of desaturation episodes (≥3%) occurring per hour.

F-hypopnoea=hypopnoea events/(hypopnoea events+apnoea events).

AHI, Apnoea–Hypopnoea Index; ArI, Arousal Index; ArTH, arousal threshold; F-hypopnoea, fraction of hypopnoea; minSpO₂, minimum values in oxygen saturation as measured using pulse oximetry; NREM, non-rapid eye movement; ODI, Oxygen Desaturation Index; REM, rapid eye movement; SPT, sleep period of time; TST, total sleep time; WASO, wake time after sleep onset.

Estimation of air pollutant exposure level

This retrospective study acquired hourly monitoring data on air pullulation levels, temperature, humidity and rainfall from 16 air quality stations subsidised and organised by Taiwan Environmental Protection Administration (19 stations, excluding 1 national park station and 2 background stations). The air pollutants included PM₁₀, PM_{9.5}, carbon monoxide (CO), NO, NO, Sulfur dioxide (SO,) and ozone (O₃). All the exposure data for the participants were calculated backward from the time of visiting the sleep centre. This study used an adjusted form of a previously reported estimation approach for air pollutant exposure values, which were determined using the data from the nearest station.²² Specifically, in this method, we employed the data obtained from air pollution monitoring stations situated within a 3 km radius of the homes of the participants; these stations were considered to be neighbouring stations. Next, as shown in online supplemental figure S1, weights were assigned to these selected stations based on the distances between the stations and the participants' homes; then, the weighted average of daily exposure levels was computed. Online supplemental figure S2 provides a comparison of PM₁₀ and PM_{9.5} exposure outcomes obtained from two methods: the original nearest station estimation method and the

revised method with reference data from the neighbouring stations. The distance was restricted, so that the nearest station was selected, thereby causing imprecise estimates. This study calculated the mean from 1, 3, 6 and 12 months prior to the date of PSG for determining the short-term, medium-term and long-term exposure, respectively.

Statistical analysis

To explore the associations between air pollution, body water distribution, arousal response and OSA severity, multiple linear regression models were employed. To determine the risk of low-ArTH from air pollutant exposure, this study classified the individuals into two groups, namely healthy and low-ArTH OSA groups, on the basis of the aforementioned criteria using clinical screening variables. Multivariable logistic regression models were employed to investigate mean air pollutant exposure between the healthy (AHI<5 events/hour) and low-ArTH groups. The effects of air pollutants on sleep parameters were analysed using the unit of the IQR alterations for individual pollutant exposure. The level of significance was set to p<0.05. SPSS (V.20.0; IBM) was used for all statistical analyses.

RESULTS

Description of basic characteristics, PSG parameters and exposure level to air pollutants of the recruited population

The demographic characteristics of participants (N=1924) are summarised in table 1, and their PSG parameters are presented in table 2. The mean age and BMI of participants were 46.8 years and 26.13 kg/m², respectively, and 60.81% of them were men. Regarding body water distribution, ECW% versus ICW% was 41.54%-58.46%, and the mean derived I-E water ratio was 141.65 (SD: 14.95). Regarding OSA severity, we recorded 326 participants with low OSA severity (16.94%), 541 with mild OSA (28.12%), 431 with moderate OSA (22.4%) and 626 with severe OSA (32.54%). Participants exhibited mean sleep efficiency of 76.76%, ODI of 18.43 events/hour, and ArI of 18.08 events/hour. Regarding the low-ArTH screening criteria, the means of AHI, minSpO2, and F-hypopnoea were 23.88 events/hour, 86.16% and 89.07%, respectively.

Table 3 details the exposure level to air pollutants presented as medians with IQRs. The medians of 1-month, 3-month, 6-month and 12-month exposure were similar, whereas the IQRs indicated a relative decreasing pattern over time.

Associations between air pollution and sleep disorder indices

This study analysed the associations between air pollutant levels at different time scales and different sleep disorder indices. The findings are summarised in table 4 (in the form of per one unit of increment in IQR and 95% CI). We observed that the IQRs of PM_{10} and $\mathrm{PM}_{2.5}$ for both

Exposure values to air pollution and background conditions of the sample

| | Median (IQR) | | | | |
|---------------------------|--------------|--------------|--------------|--------------|--|
| Categorical variables | 1 month | 3 months | 6 months | 12 months | |
| Air pollutant | | | | | |
| PM ₁₀ (μg/m³) | 24.03 (3.4) | 23.45 (2.69) | 24.12 (2.13) | 24.97 (1.75) | |
| PM _{2.5} (µg/m³) | 12.75 (2.41) | 12.41 (1.68) | 12.61 (1.45) | 13.03 (0.66) | |
| CO (ppm) | 0.44 (0.07) | 0.43 (0.06) | 0.43 (0.05) | 0.44 (0.02) | |
| NO (ppb) | 4.47 (0.85) | 4.55 (0.72) | 4.47 (0.43) | 4.52 (0.29) | |
| NO ₂ (ppb) | 14.94 (2.07) | 14.96 (2.21) | 14.7 (1.6) | 15.22 (1.15) | |
| SO ₂ (ppb) | 1.86 (0.21) | 1.89 (0.24) | 1.89 (0.21) | 1.94 (0.22) | |
| O ₃ (ppb) | 29.28 (3.99) | 29.16 (2.95) | 28.75 (1.42) | 28.91 (0.55) | |
| Background | Mean (SD) | | | | |
| Ambient temperature (°C) | 23.32 (4.49) | 23.44 (4.31) | 23.9 (3.04) | 24.07 (0.33) | |
| Relative humidity (%) | 73.21 (4.24) | 73.27 (3.75) | 73.08 (2.6) | 72.86 (1.48) | |

Date are expressed as median (IQR).

CO, carbon monoxide; NO₂, nitrogen dioxide; NO, nitrogen oxide; O₃, ozone; PM₁₀, particulate matter with an aerodynamic diameter of≤10 μm; PM_{2,5}, particulate matter with an aerodynamic diameter of ≤2.5 μm; SO₂, sulfur dioxide.

1-month and 3-month exposure were significantly associated with AHI values (p<0.01). The short-term exposure IQRs of CO, NO₉ and O₈ were positively associated with AHI values (p<0.05). The association patterns of ODI were identical to those of AHI. Significant associations were observed between ArI and PM₁₀, PM_{9.5} and O₃ in the short and medium term (p<0.05).

Associations between air pollution and body water distribution

This study then analysed the associations between air pollutant levels and the body water distribution of participants. The obtained outcomes are presented in table 5 (per one unit of increment in IQR and 95% CI). Total body water (TBW) was positively associated with the IQRs of PM₁₀ and NO₉ in both the short and medium term, whereas PM_{9.5} was only associated with the medium-term IQRs (p<0.05). PM₁₀ was associated with the I-E water ratio in the short term, whereas both PM₁₀ and PM₉₅ were associated with the I-E water ratio in the medium term. The associations between exposure to air pollutants and body water distribution were similar to those with the sleep disorder indices; specifically, the associations were only observed in the short (1 month) and medium term (3 months).

Associations between PM exposure and sleep disorder indices considering the effects of body water distribution

Significant associations were found between 1 month PM₁₀ exposure, 3-month PM exposure, sleep disorder indices and the I-E water ratio. Therefore, the synergistic associations between PM exposure and sleep disorder indices were further analysed considering the effects of body water distribution. As presented in table 6, the results revealed that an IQR increase in 1-month PM₁₀

exposure was positively associated with increases of 4.24, 3.91 and 1.44 events/hour in the AHI, ODI and ArI values, respectively (95% CIs 2.82 to 5.66, 2.51 to 5.31 and 0.56 to 2.32 events/hour, respectively; all p<0.01). Moreover, an IQR increase in 3-month PM₁₀ exposure was positively associated with increases of 2.06, 2.01 and 1.06 events/hour in the AHI, ODI and ArI values, respectively (95% CIs 0.69 to 3.43, 0.67 to 3.35 and 0.22 to 1.89 events/hour; p<0.01, p<0.01 and p<0.05, respectively). Similarly, an IQR increase in 3-month PM_{9.5} exposure was positively associated with increases of 2.57, 2.43 and 1.07 events/hour in the AHI, ODI and ArI values, respectively (95% CIs 0.98 to 4.15, 0.88 to 3.99 and 0.11 to 2.04 events/hour; p<0.01, p<0.01 and p<0.05, respectively). Moreover, significant associations were observed between 1-month PM_{9.5} exposure and sleep disorder indices in the analysis considering body water distribution (ie, the I-E water ratio). However, the difference when examining the associations of 1-month PM₉₅ exposure with the I-E water ratio was nonsignificant (table 5). Therefore, only 1-month PM₁₀ exposure and 3-month PM₁₀ and PM₂₅ exposure were included in the mediation analysis.

Mediation analysis of associations between air pollution, body water distribution and sleep disorder index

Next, we investigated the associations between air pollution, body water distribution and OSA severity using mediation analysis (figure 2). This analysis included PM₁₀ (short and medium term), PM_{9.5} (short term), I-E water ratio, AHI and ODI because they demonstrated significant associations after adjustment for age, sex and BMI; for fine PM exposure, ambient temperature and humidity were further adjusted for. The indirect (mediated) effects of PM₁₀ (short and medium term) and PM_{9.5} (short term) on the I-E water ratio (path 1) were positive

Table 4 Associations between the sleep disorder indices and an IQR alteration in exposure to short-term, medium-term and long-term air pollution

| Categorical | Beta coefficient (95% CI) | | | | |
|--|---------------------------|-----------------------|-----------------------|----------------------|--|
| variables | 1 month | 3 months | 6 months | 12 months | |
| AHI (events/hour) | | | | | |
| PM ₁₀ (μg/m ³) | 4.33 (2.91 to 5.75)** | 2.16 (0.79 to 3.53)** | .62 (-0.48 to 1.73) | 0.67 (-0.56 to 1.9) | |
| PM _{2.5} (µg/m ³) | 4.48 (3.02 to 5.93)** | 2.67 (1.09 to 4.25)** | 0.18 (-1.3 to 1.65) | -0.79 (-1.9 to 0.32) | |
| CO (ppm) | 2.22 (0.93 to 3.52)** | 1.21 (-0.3 to 2.72) | -0.21 (-1.42 to 1.01) | 0.38 (-0.34 to 1.11) | |
| NO (ppb) | 0.53 (-0.17 to 1.22) | 0.3 (-0.3 to 0.9) | 0.12 (-0.37 to 0.62) | 0.14 (-0.33 to 0.61) | |
| NO ₂ (ppb) | 2.4 (1.07 to 3.73)** | 1.29 (-0.26 to 2.84) | 0.16 (-1.19 to 1.5) | 0.03 (-0.94 to 1.01) | |
| SO ₂ (ppb) | -0.36 (-1.08 to 0.36) | 0.38 (-0.52 to 1.28) | 0.18 (-0.75 to 1.11) | 0.16 (-0.86 to 1.17) | |
| O ₃ (ppb) | 1.66 (0.43 to 2.88)* | 0.52 (-0.58 to 1.61) | 0.71 (-0.43 to 1.84) | 0.1 (-0.73 to 0.92) | |
| ODI (events/hour) | | | | | |
| PM ₁₀ (μg/m ³) | 4.01 (2.62 to 5.41)** | 2.12 (0.78 to 3.46)** | 0.7 (-0.39 to 1.78) | 1.02 (-0.19 to 2.22) | |
| PM _{2.5} (µg/m ³) | 4.21 (2.78 to 5.64)** | 2.55 (0.99 to 4.1)** | 0.27 (-1.18 to 1.72) | -0.69 (-1.78 to 0.4) | |
| CO (ppm) | 2.4 (1.13 to 3.67)** | 1.31 (-0.17 to 2.79) | -0.04 (-1.23 to 1.15) | 0.27 (-0.44 to 0.98) | |
| NO (ppb) | 0.61 (-0.07 to 1.29) | 0.3 (-0.29 to 0.88) | 0.16 (-0.33 to 0.64) | 0.25 (-0.21 to 0.71) | |
| NO ₂ (ppb) | 2.48 (1.17 to 3.78)** | 1.32 (-0.2 to 2.85) | 0.24 (-1.08 to 1.57) | 0.18 (-0.78 to 1.14) | |
| SO ₂ (ppb) | -0.35 (-1.06 to 0.36) | 0.51 (-0.38 to 1.39) | 0.34 (-0.57 to 1.25) | 0.34 (-0.66 to 1.34) | |
| O ₃ (ppb) | 1.44 (0.24 to 2.64)* | 0.32 (-0.76 to 1.4) | 0.85 (-0.26 to 1.96) | 0.06 (-0.75 to 0.87) | |
| Arl (events/hour) | | | | | |
| PM ₁₀ (µg/m ³) | 1.48 (0.6 to 2.36)** | 1.09 (0.26 to 1.93)* | 0.35 (-0.32 to 1.02) | -0.55 (-1.29 to 0.19 | |
| PM _{2.5} (µg/m ³) | 1.63 (0.73 to 2.53)** | 1.11 (0.14 to 2.07)* | -0.1 (-0.99 to 0.8) | -0.17 (-0.84 to 0.49 | |
| CO (ppm) | 0.5 (-0.3 to 1.3) | 0.58 (-0.34 to 1.5) | -0.25 (-0.99 to 0.48) | -0.08 (-0.51 to 0.36 | |
| NO (ppb) | 0.23 (-0.2 to 0.65) | 0.16 (-0.2 to 0.53) | -0.08 (-0.38 to 0.23) | -0.26 (-0.54 to 0.03 | |
| NO ₂ (ppb) | 0.99 (0.17 to 1.81)* | 1.06 (0.11 to 2.01)* | 0.17 (-0.65 to 0.99) | -0.23 (-0.82 to 0.36 | |
| SO ₂ (ppb) | -0.27 (-0.72 to 0.17) | -0.36 (-0.91 to 0.19) | -0.44 (-1.01 to 0.12) | -0.5 (-1.11 to 0.12) | |
| O ₃ (ppb) | -0.31 (-1.06 to 0.45) | 0.36 (-0.31 to 1.03) | 0.62 (-0.07 to 1.31) | -0.04 (-0.54 to 0.46 | |

Multivariable linear regression models were adjusted for age, sex, body mass index, temperature and relative humidity. *p<0.05, **p<0.01.

AHI, Apnoea-Hypopnoea Index; ArI, Arousal Index; CO, carbon monoxide; NO, nitrogen dioxide; NO, nitrogen oxide; O,, ozone; ODI, Oxygen Desaturation Index; PM₁₀, particulate matter with an aerodynamic diameter of ≤10 µm; PM₂₅, particulate matter with an aerodynamic diameter of ≤2.5 µm; SO₂, sulfur dioxide.

and significant, as indicated by the regression coefficients (p<0.05). Next, the total effect (mediated and direct) of short-term exposure to PM₁₀ and medium-term exposure to PM₁₀ on AHI (regression coefficients of 4.33 and 2.16) and on ODI (4.01 and 2.12) were positive and significant, respectively (path 3). For the total effect (mediated and direct) of medium-term exposure to PM_{9,5}, the regression coefficients were 2.67 for AHI and 2.55 for ODI. After considering the indirect (mediated) effects between air pollutants and the I-E water ratio on AHI or ODI (path 3), the partial mediation effects were determined. In mediation analysis, the associations of other air pollutants with ArI was nonsignificant.

Associations between air pollution and the risk of low-ArTH

To explore the associations between air pollution and the risk of low-ArTH OSA, we compared the exposure levels to various air pollutants between healthy participants and patients with low-ArTH OSA using logistic regression (ORs)). As summarised in table 7, we observed that shortterm PM₁₀, PM_{2.5} and O₃ exposure was significantly associated with an elevated OR of low-ArTH OSA incidence. In particular, an IQR increment in PM₁₀, PM₉₅ and O₃ exposure was associated with an increased OR of 1.42, 1.33 and 1.27, respectively (95% CIs 1.09 to 1.84, 1.02 to 1.74 and 1.01 to 1.6, respectively).

DISCUSSION

Sleep disorder indices are associated with air pollution and body water distribution. However, the relationships between them and the effects of air pollution on arousal require further investigation. Thus, this study collected data regarding PSG, body composition and estimated exposure levels to various air pollutants and investigated

Associations between the body water distribution and an IQR alteration in exposure to short-term, medium-term and long-term air pollution

| Categorical | Beta coefficient (95% CI) | | | | |
|--|---------------------------|-----------------------|------------------------|-----------------------|--|
| variables | 1 month | 3 months | 6 months | 12 months | |
| TBW (%) | | | | | |
| PM ₁₀ (μg/m ³) | 0.33 (0.01 to 0.65)* | 0.45 (0.14 to 0.76)** | 0.25 (-0.0 to 0.49) | 0.19 (-0.08 to 0.47) | |
| PM _{2.5} (µg/m ³) | 0.21 (-0.12 to 0.54) | 0.43 (0.07 to 0.79)* | 0.26 (-0.07 to 0.59) | -0.03 (-0.28 to 0.22) | |
| CO (ppm) | 0.15 (-0.14 to 0.45) | 0.2 (-0.14 to 0.54) | 0.11 (-0.16 to 0.39) | -0.03 (-0.19 to 0.13) | |
| NO (ppb) | 0.09 (-0.07 to 0.25) | 0.02 (-0.11 to 0.16) | 0.01 (-0.1 to 0.12) | -0.03 (-0.13 to 0.08) | |
| NO ₂ (ppb) | 0.35 (0.05 to 0.65)* | 0.38 (0.04 to 0.73)* | 0.25 (-0.06 to 0.55) | -0.03 (-0.25 to 0.19) | |
| SO ₂ (ppb) | 0.05 (-0.11 to 0.21) | 0.15 (-0.05 to 0.36) | 0.15 (-0.06 to 0.36) | 0.17 (-0.05 to 0.4) | |
| O ₃ (ppb) | -0.04 (-0.31 to 0.24) | 0.11 (-0.14 to 0.36) | -0.05 (-0.31 to -0.21) | 0.04 (-0.15 to 0.22) | |
| I-E water ratio | | | | | |
| $PM_{10} (\mu g/m^3)$ | 1.13 (0.22 to 2.04)* | 1.18 (0.3 to 2.06)* | 0.56 (-0.15 to 1.26) | 0.48 (-0.31 to 1.26) | |
| PM _{2.5} (µg/m ³) | 0.88 (-0.06 to 1.82) | 1.2 (0.19 to 2.22)* | 0.66 (-0.29 to 1.6) | -0.07 (-0.78 to 0.63) | |
| CO (ppm) | 0.36 (-0.46 to 1.19) | 0.47 (-0.5 to 1.44) | 0.2 (-0.57 to 0.98) | -0.12 (-0.58 to 0.33) | |
| NO (ppb) | 0.13 (-0.32 to 0.57) | 0.01 (-0.38 to 0.39) | -0.01 (-0.33 to 0.3) | -0.09 (-0.39 to 0.21) | |
| NO ₂ (ppb) | 0.86 (0.01 to 1.71) | 0.86 (-0.14 to 1.85) | 0.46 (-0.4 to 1.33) | -0.15 (-0.77 to 0.47) | |
| SO ₂ (ppb) | 0.09 (-0.37 to 0.55) | 0.24 (-0.33 to 0.82) | 0.14 (-0.46 to 0.73) | 0.26 (-0.38 to 0.91) | |
| O ₃ (ppb) | 0.34 (-0.44 to 1.12) | 0.25 (-0.45 to 0.96) | 0.16 (-0.57 to 0.88) | 0.07 (-0.45 to 0.6) | |

Multivariable linear regression models were adjusted for age, sex, body mass index, temperature and relative humidity; calculation of I-E water ratio: ICW/ECW×100.

CO, carbon monoxide; I-E water ratio, ratio of intracellular water to extracellular water; NO, nitrogen oxide; NO₀, nitrogen dioxide; O₀, ozone; PM₁₀, particulate matter with an aerodynamic diameter of ≤10 µm; PM_{2.5}, particulate matter with an aerodynamic diameter of ≤2.5 µm; SO₂, sulfur dioxide; TBW, total body water.

their relationships using multivariable linear regression models and mediation analysis. Moreover, the associations between air pollution and the risk of low-ArTH OSA were explored.

In this study, we estimated the long-term exposure levels of PM₂₅ and PM₁₀ to be 13.03 and 24.97 μ g/m³, respectively, both of which were higher than their annual means in the 2021 air quality guidelines of the WHO (5 and 15 μg/m³, respectively). 23 Here, we proposed a revised method where we used data from stations in the proximity of the participants' homes by limiting the maximum distance of the stations rather than using data only from the nearest stations.²⁴ The supplemental results suggested that this revised approach offers a more precise approximation to the actual data than methods that solely reference the air pollution readings of the nearest station. The proposed approach may enhance data reliability by reducing potential estimation errors that occur when relying on only one information source. For instance, if the nearest station is at a considerable distance, the obtained results may be less reliable.

In this study, 1-month PM₁₀, PM₉₅, CO, NO₉ and O₃ exposure was significantly associated with increased AHI and ODI values. Similar outcomes have also been observed for the 3-month exposure to PM₁₀ and PM₉₅. Pertaining to the arousal response, exposure to PM₁₀,

PM_{9,5} and NO₉ in the 1-month and 3-month was positively and significantly associated with ArI values. Consequently, air pollution potentially affects the physiology of both the respiratory and central nervous systems. Specifically, exposure to air pollutants is associated with inflammatory responses in the upper and lower respiratory tract, which may increase airway resistance and thereby aggravate OSA severity.²⁵ Fine particles and gaseous pollutants (eg, NO₉) may also irritate the respiratory system and cause oedema in the nasopharyngeal and oropharyngeal tracts and subsequent airway narrowing, thereby increasing sleep disorder indices. 26 A study examined the relationships between sleep variables from PSG data and mean PM₁₀ exposure levels in winter (December and January) and summer (June and July). 27 Its results demonstrated the AHI values were higher in winter than in summer, aligning with increased PM₁₀ levels. Various reasons could account for these observed relationships. Air pollution may also mediate the development of an inflammatory response and may cause the bronchial physiology to become unstable, which has been identified as a risk factor for OSA. 28 29 PM may accumulate in the lower respiratory tract and induce inflammation and an oxidative response, possibly increasing oxidative stress, overwhelming the antioxidant defence and increasing hypoxaemia risk.³⁰ Consistent with the current findings,

^{*}p<0.05; **p<0.01.

Table 6 Associations between sleep disorder indices and IQR variations for 1-month and 3-month exposure to fine PM considering the effects of body water distribution

| Beta coefficient |
|-----------------------|
| (95% CI) |
| |
| 4.24 (2.82 to 5.66)** |
| 3.91 (2.51 to 5.31)** |
| 1.44 (0.56 to 2.32)** |
| |
| 4.40 (2.95 to 5.86)** |
| 4.13 (2.70 to 5.56)** |
| 1.61 (0.71 to 2.51)** |
| |
| 2.06 (0.69 to 3.43)** |
| 2.01 (0.67 to 3.35)** |
| 1.06 (0.22 to 1.89)* |
| |
| 2.57 (0.98 to 4.15)** |
| 2.43 (0.88 to 3.99)** |
| 1.07 (0.11 to 2.04)* |
| |

Multivariable linear regression models were adjusted for age, sex, body mass index, temperature and relative humidity.

Unit of outcome (ie, sleep disorder indices) was subjected to Box-Cox transformation.

*p<0.05; **p<0.01.

AHI, Apnoea–Hypopnoea Index; ArI, Arousal Index; ODI, Oxygen Desaturation Index; PM $_{10}$, particulate matter with an aerodynamic diameter of \leq 10 μ m; PM $_{2.5}$, particulate matter with an aerodynamic diameter of \leq 2.5 μ m.

a study reported positive associations between air pollution and worsened OSA severity, as follows: $PM_{2.5}$ –AHI, NO_2 –ArI and NO_2 –AHI. 31 Another multiethnic study observed that higher annual exposure concentrations of NO_2 (10 ppb) and $PM_{2.5}$ (5 $\mu g/m^3$) were associated with an increased risk of OSA incidence. 32 Public awareness regarding the effects of air pollutant exposure on sleep disorder indices should be increased.

Regarding the effects of air pollution on the alterations of body water distribution, we determined that 1-month PM₁₀ and 3-month PM₁₀ and PM_{2.5} exposure was associated with increased TBW and I-E water ratio after adjustment for age, sex and BMI. Although no direct evidence or mechanisms support that air pollution may increase water retention, particularly for ICW, some explanations may account for these observations. Specifically, following exposure to air pollutants, the defence mechanisms of the epithelium and mucosa in the respiratory system are deactivated, including excessive inflammation in cells and reduced barrier function. ³³ A study investigated the molecular mechanisms in the cell membrane of airway epithelial cells and the results indicated that PM exposure might affect the permeability of these cells, leading




Figure 2 Mediation analysis of air pollutants between the I-E water ratio and sleep disorder indices. Mediation analysis of I-E water between PM with a 1-month and 3-month PM₁₀ or PM₂₅ exposure and sleep disorder indices (AHI and ODI). The beta coefficients of the adjusted linear regression models and derived p values are presented in the path diagram. The outcomes demonstrated that I-E water ratio partially mediates the effects of 1-month PM₁₀, 3-month PM₁₀ and 3-month PM₂₅ exposure on the AHI and ODI values. All four requirements or a mediation effect were satisfied: path-1, path-2, path-3 and path-3* were statistically significant. Note: path-1, path-3 and path-3*: adjusted for age, sex, body mass index, temperature and relative humidity; path-2: adjusted for age, sex and body mass index. AHI, Apnoea-Hypopnoea Index; I-E water ratio, the ratio of intracellular water to extracellular water: ODI, Oxygen Desaturation Index; PM₁₀, aerodynamic diameter of <10 µm; PM₂₅, aerodynamic diameter of $< 2.5 \mu m$.

to tissue swelling.³⁴ These physiological responses to air pollutants may indirectly contribute to the alteration of the water ratio between the intracellular and extracellular compartments. Exposure to fine particles may also exacerbate oxidative stress in the intracellular environment, resulting in inflammation and oedema. 35 Another possible explanation is the osmotic pressure changes in the cell membrane caused by air pollution. Studies have reported that the organic components of PM_{9.5} may contribute to cell cycle dysregulation as well as several other responses (ie, elevated osmotic pressure and exacerbation of cell inflammation).³⁶ Additionally, the toxic effects of fine particle exposure may increase the vascular permeability of endothelial cells and may further cause cell swelling because of the excess fluid in the intracellular environment.³⁷ On the basis of the association of the 1-month and 3-month $\mathrm{PM}_{2.5}$ and PM_{10} exposure with the I-E water ratio of body water distribution and sleep disorder indices, we performed mediation analysis to examine their interactions and their partial mediation effects. To the best of our knowledge, this study may be the first to explore these mediation effects. No study has reported similar results or indicated the mechanisms

Table 7 Associations (ORs) of IQR alterations in exposure to short-term, medium-term and long-term air pollution between the healthy and OSA with low-ArTH groups

| Categorical variables | OR (95% CI) | | | | |
|---------------------------------------|----------------------|---------------------|---------------------|---------------------|--|
| | 1 month | 3 months | 6 months | 12 months | |
| PM ₁₀ (μg/m ³) | 1.42 (1.09 to 1.84)* | 1.31 (0.99 to 1.72) | 1.02 (0.83 to 1.26) | 0.92 (0.74 to 1.15) | |
| PM _{2.5} (μg/m³) | 1.33 (1.02 to 1.74)* | 1.36 (1.0 to 1.84) | 1.03 (0.78 to 1.36) | 1.03 (0.85 to 1.26) | |
| CO (ppm) | 1.06 (0.85 to 1.33) | 1.1 (0.84 to 1.44) | 1.02 (0.83 to 1.26) | 1.13 (0.99 to 1.29) | |
| NO (ppb) | 1.04 (0.93 to 1.18) | 1.09 (0.97 to 1.21) | 1.07 (0.98 to 1.17) | 1.04 (0.96 to 1.13) | |
| NO ₂ (ppb) | 1.16 (0.92 to 1.48) | 1.24 (0.94 to 1.65) | 1.17 (0.92 to 1.49) | 0.95 (0.79 to 1.13) | |
| SO ₂ (ppb) | 1.1 (0.97 to 1.25) | 1.05 (0.89 to 1.24) | 1.04 (0.88 to 1.24) | 1.05 (0.88 to 1.26) | |
| O ₃ (ppb) | 1.27 (1.01 to 1.6)* | 0.96 (0.79 to 1.17) | 0.89 (0.72 to 1.09) | 0.94 (0.82 to 1.09) | |

^{*}Multivariable linear regression models were adjusted for age, sex, body mass index, temperature and relative humidity. p<0.05.

CO, carbon monoxide; low-ArTH, low-arousal-threshold; NO, nitrogen oxide; NO₂, nitrogen dioxide; O₃, ozone; OSA, obstructive sleep apnoea; PM₁₀, particulate matter with an aerodynamic diameter of ≤10 μm; PM₂₅, particulate matter with an aerodynamic diameter of ≤2.5 µm; SO, sulfur dioxide.

of these synergistic effects. However, these underlying mechanisms may explain how fine particles directly aggravate OSA severity and increase the I-E water ratio of body water, which may indirectly and further aggravate OSA severity. Taken together, the results indicate that air pollution may alter the distribution and volume of body water by affecting cell physiology.

This study highlighted a significant relationship of 1-month PM₁₀ and PM₉₅ exposure with an increase in ArI. Additionally, exposure to air pollutants (ie, PM_{9.5}, PM₁₀ and O₂) was strongly associated with an elevated risk of low-ArTH OSA, suggesting that the aforementioned pollutants are risk factors for low-ArTH OSA. Although the mechanism underlying the specific association between air pollution and the declined ArTH remains unclear, some reasons may account for these findings. Specifically, the ArTH is influenced by the interactions of both the respiratory and central nervous systems. This threshold can be considered as the action point for generating an arousal response when experiencing airway occlusion. Therefore, abnormalities in both the respiratory and central nervous systems may be responsible for the impaired ArTH, which may result in awakening in patients with OSA because of the arousal responses induced by slight hypoxia stimuli.³⁸ In addition, PM and O_s exposure may impair respiratory and central nervous system function by causing inflammation, cell swelling and systemic oedema.³⁹ For example, PM exposure may mediate toxicological mechanisms, causing neuroinflammation in the central nervous system and indirectly increasing sleep arousal frequency. 41 A comprehensive review revealed that elevated levels of PM exposure, even for short durations (ie, a few hours to days), can result in unstable brain haemodynamics. 42 A study suggested that short-term PM exposure damaged neurons, thus negatively affecting the central nervous system. 43 Another study noted positive associations between monthly short-term exposure to fine PM and ArI⁴⁴; this result corroborated

the current results. An animal model study indicated that PM might reach the central nervous system indirectly through the peripheral system, eventually increasing the permeability of the blood-brain barrier after PM exposure for 4–24 hours. 45 Another study proposed that shortterm (1-3 months) exposure to PM results in metal and inflammatory biomarker accumulation in rat brains.⁴⁶ These physiological disturbances within the central nervous system may lead to heightened sleep arousal or disrupted sleep patterns. These potential underlying mechanisms and the related outcomes indicate that PM exposure provokes sleep arousal responses, affecting the central nervous system and thus increasing sleep arousal frequency and the risk of low-ArTH OSA. Nevertheless, further research verifying the cause-and-effect relationship between short-term PM exposure and its adverse influence on the central nervous system, which may affect the ArTH, is warranted.

This study has some limitations. Indoor and outdoor air quality may differ, and we did not measure indoor pollution. Relatedly, air pollution levels estimated based on residential addresses might not reflect the actual exposure accurately. In terms of modelling, we did not identify the effects of short-term PM_{9.5} exposure on sleepdisordered measurements by using a single-pollutant model. Using air pollution measurements for each patient may improve the air pollutant exposure accuracy and enable the detailed evaluation of short-term relationships between pollution and sleep disorders. Regarding the social demography perspective of this study, self-payment at the sleep centre may have led to sampling bias. Our sample may not be representative of the general population. The socioeconomic status or lifestyle habits of participants were not obtained, which this may have may affected OSA manifestations or severity and may have resulted in potential residual confounding. 47 48 Data on psychosocial stressors, which may also confound the associations between air pollution

and OSA, were also missing, such as noise, medication and continuous positive airway pressure therapy. Further questionnaire surveys should collect this information for future research.

CONCLUSION

This study observed that air pollution, including 1-month exposure to PM_{9.5}, PM₁₀, CO, NO₉ and O₃ and 3-month exposure to $PM_{2.5}^{-3}$ and PM_{10} , was significantly associated with increased AHI and ODI values. Both 1-month and 3-month exposure to PM_{2.5}, PM₁₀ and NO₂ was significantly associated with increased ArI values. Significant associations were observed between the I-E water ratio, 1-month and 3-month PM₁₀ exposure, and 1-month PM_{9.5} exposure. Next, sleep disorder indices and 1-month and 3month PM exposure exhibited significant associations even in the context of the effects of I-E water ratio. Similarly, 1-month and 3-month PM_{10} and 1-month $PM_{2.5}$ exposure had mediation effects on the I-E water ratio, which in turn partially worsened sleep disorder indices by increasing the AHI and ODI values. Moreover, 1-month PM₁₀, PM_{9.5} and O₃ exposure was significantly associated with the risk of low-ArTH OSA.

Taken together, these results suggest that air pollutant exposure directly worsens sleep disorder indices and influences body water distribution, thereby indirectly increasing AHI and ODI values. Furthermore, PM_{10} , $PM_{2.5}$ and O_3 exposure may impact the central nervous system, increasing arousal response frequency and thus potentially increasing the risk of low-ArTH OSA. Therefore, mitigating air pollutant exposure may improve sleep disorder indices and reduce the risk of low-ArTH OSA.

Author affiliations

¹Department of Civil and Environmental Engineering, Imperial College London, London, UK

²Division of Pulmonary Medicine, Department of Internal Medicine, Taipei Medical University-Shuang Ho Hospital Ministry of Health and Welfare, New Taipei City, Taiwan

³Department of Biology, University of Oxford, Oxford, UK

⁴Department of Medical Physics and Biomedical Engineering, University College London, London, UK

⁵School of Respiratory Therapy, College of Medicine, Taipei Medical University, Taipei, Taiwan

⁶Sleep Center, Taipei Medical University-Shuang Ho Hospital Ministry of Health and Welfare, New Taipei City, Taiwan

⁷Department of Neurology, Taipei Medical University-Shuang Ho Hospital Ministry of Health and Welfare, New Taipei, Taiwan

⁸Department of Neurology, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan

⁹Neuroscience Institute, Taipei Medical University, Taipei, Taiwan

¹⁰Division of Pulmonary Medicine, Department of Internal Medicine, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan

¹¹Research Center of Artificial Intelligence in Medicine, Taipei Medical University College of Medicine, Taipei, Taiwan

¹²Department of Physical Medicine and Rehabilitation, Taipei Medical University Hospital, Taipei, Taiwan

¹³Graduate Institute of Nanomedicine and Medical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei, Taiwan

¹⁴Department of Psychiatry, Taipei Medical University Hospital, Taipei, Taiwan
¹⁵Department of Otolaryngology, Taipei Medical University-Shuang Ho
Hospital Ministry of Health and Welfare, New Taipei, Taiwan

Acknowledgements We thank all of the participants for their involvement with this study as well as the technologists in the sleep centre of Shuang Ho Hospital for collecting the raw data.

Contributors C-YT, ML and H-TH carried out the findings analysis and drafted the main manuscript text. W-HH and Y-CK performed data curation and investigation. AM validated the results. K-YL and J-HK led the project administration and offered opinions. P-HF, C-HT and K-YC made suggestions to conceptualise the project. H-CL and C-JW provided feedback and adjustment. W-TL conceptualised the project, review and editing the manuscript. Additionally, W-TL is responsible for the overall content.

Funding This study was funded by Taiwan Ministry of Science and Technology (grant number: MOST 110-2634-F-002-049) and Taiwan National Science and Technology Council (grant number NSTC 111-2634-F-002-021).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval The procedure of this retrospective study was reviewed and approved by Taipei Medical University-Joint Institutional Review Board (TMU-JIRB: N202212067). All the procedures related to data assessment, deidentification of individual information, statistical analysis and data storage or maintenance were conducted in accordance with the approved protocol.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. All the data of this study were collected at the Sleep Center of Taipei Medical University—Shuang Ho Hospital (New Taipei City, Taiwan) between June 2019 and December 2021. Because our data set contains personal information, it is not available in online supplement file. For access to the data set or relevant documents, please contact the corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID ID

Wen-Te Liu http://orcid.org/0000-0003-1281-8718

REFERENCES

- 1 Benjafield AV, Ayas NT, Eastwood PR, et al. Estimation of the global prevalence and burden of obstructive sleep apnoea: a literaturebased analysis. Lancet Respir Med 2019;7:687–98.
- 2 Maspero C, Giannini L, Galbiati G, et al. Obstructive sleep apnea syndrome: a literature review. Minerva Stomatol 2015;64:97–109.
- 3 Subramani Y, Singh M, Wong J, et al. Understanding phenotypes of obstructive sleep apnea: applications in anesthesia, surgery, and perioperative medicine. Anesth Analg 2017;124:179–91.
- 4 Eckert DJ, White DP, Jordan AS, et al. Defining phenotypic causes of obstructive sleep apnea. identification of novel therapeutic targets. Am J Respir Crit Care Med 2013;188:996–1004.
- 5 Zinchuk A, Edwards BA, Jeon S, et al. Prevalence, associated clinical features, and impact on continuous positive airway pressure use of a low respiratory arousal threshold among male United States veterans with obstructive sleep apnea. J Clin Sleep Med 2018;14:809–17.
- 6 Hsu W-H, Yang C-C, Tsai C-Y, et al. Association of low arousal threshold obstructive sleep apnea manifestations with body fat and water distribution. Life (Basel) 2023;13:1218.
- 7 Qiu H, Liu W-T, Lin S-Y, et al. Association of air pollution exposure with low arousal threshold obstructive sleep apnea: a crosssectional study in Taipei, Taiwan. *Environ Pollut* 2022;306:119393.



- 8 Edwards BA, Eckert DJ, McSharry DG, et al. Clinical predictors of the respiratory arousal threshold in patients with obstructive sleep apnea. Am J Respir Crit Care Med 2014;190:1293–300.
- 9 Cao B, Chen Y, McIntyre RS. Comprehensive review of the current literature on impact of ambient air pollution and sleep quality. Sleep Med 2021:79:211–9.
- 10 Holley AB, Walter RJ. Does a low arousal threshold equal low PAP adherence J Clin Sleep Med 2018;14:713–4.
- 11 Chamberlin NL. Brain circuitry mediating arousal from obstructive sleep apnea. *Curr Opin Neurobiol* 2013;23:774–9.
- 12 Zhang Q, Li Q, Ma J, et al. Pm2.5 impairs neurobehavior by oxidative stress and myelin sheaths injury of brain in the rat. Environ Pollut 2018;242:994–1001.
- 13 Vallianatou T, Lin W, Bèchet NB, et al. Differential regulation of oxidative stress, microbiota-derived, and energy metabolites in the mouse brain during sleep. J Cereb Blood Flow Metab 2021;41:3324–38.
- 14 Simon MJ, Iliff JJ. Regulation of cerebrospinal fluid (CSF) flow in neurodegenerative, neurovascular and neuroinflammatory disease. *Biochim Biophys Acta* 2016;1862:442–51.
- 15 Brockmeyer S, D'Angiulli A. How air pollution alters brain development: the role of neuroinflammation. *Transl Neurosci* 2016;7:24–30.
- 16 Oberdörster G, Sharp Z, Atudorei V, et al. Translocation of inhaled ultrafine particles to the brain. *Inhal Toxicol* 2004;16:437–45.
- 17 Trevino RJ. Air pollution and its effect on the upper respiratory tract and on allergic rhinosinusitis. Otolaryngol Head Neck Surg 1996;114:239–41.
- 18 Evans S-A, Al-Mosawi A, Adams RA, et al. Inflammation, edema, and peripheral blood changes in lung-compromised rats after instillation with combustion-derived and manufactured nanoparticles. Exp Lung Res 2006;32:363–78.
- 19 Sun J, Yu J, Shen Z, et al. Oxidative stress-inducing effects of various urban Pm2. 5 road dust on human lung epithelial cells among 10 Chinese Megacities. Ecotoxicol Environ Saf 2021;224:112680.
- 20 Berry RB, Brooks R, Gamaldo C, et al. AASM scoring manual updates for 2017 (version 2.4). American Academy of Sleep Medicine, 2017.
- 21 Quan S, Gillin JC, Littner M, et al. Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. Sleep 1999;22:667–89.
- 22 He Y, Liu W, Lin S, et al. Association of traffic air pollution with severity of obstructive sleep apnea in urban areas of northern Taiwan: a cross-sectional study. Sci Total Environ 2022;827:154347.
- 23 Salud OMdl, Weltgesundheitsorganisation, Organization WH, Environment ECf. WHO global air quality guidelines: particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. World Health Organization, 2021.
- 24 Meng Y-Y, Wilhelm M, Rull RP, et al. Traffic and outdoor air pollution levels near residences and poorly controlled asthma in adults. Ann Allergy Asthma Immunol 2007;98:455–63.
- 25 Zanobetti A, Redline S, Schwartz J, et al. Associations of Pm10 with sleep and sleep-disordered breathing in adults from seven US urban areas. Am J Respir Crit Care Med 2010;182:819–25.
- 26 Ghorani-Azam A, Riahi-Zanjani B, Balali-Mood M. Effects of air pollution on human health and practical measures for prevention in Iran. J Res Med Sci 2016;21:65.
- 27 Yıldız Gülhan P, Güleç Balbay E, Elverişli MF, et al. Do the levels of particulate matters less than 10 μm and seasons affect sleep Aging Male 2020;23:36–41.

- 28 D'Amato G, Liccardi G, D'Amato M, et al. Outdoor air pollution, climatic changes and allergic bronchial asthma. Eur Respir J 2002;20:763–76.
- 29 Dixit R. Asthma and obstructive sleep apnea: more than an association *Lung India* 2018;35:191–2.
- Wang Y, Tang M. 5 induces ferroptosis in human endothelial cells through iron overload and redox imbalance. *Environmental Pollution* 2019;254:112937.
- 31 Tung NT, Lee Y-L, Lin S-Y, *et al.* Associations of ambient air pollution with overnight changes in body composition and sleep-related parameters. *Sci Total Environ* 2021;791:148265.
- 32 Billings ME, Gold D, Szpiro A, et al. The Association of ambient air pollution with sleep apnea: the multi-ethnic study of atherosclerosis. Ann Am Thorac Soc 2019;16:363–70.
- 33 Zhao C, Wang Y, Su Z, et al. Respiratory exposure to Pm2. 5 soluble extract disrupts Mucosal barrier function and promotes the development of experimental asthma. Science of The Total Environment 2020;730:139145.
- 34 Liang Y, Chu PH, Tian L, et al. Targeting mitochondrial permeability transition pore ameliorates PM_{2.5}-Induced mitochondrial dysfunction in airway epithelial cells. *Environ Pollut* 2022;295:118720.
- 35 Gangwar RS, Bevan GH, Palanivel R, et al. Oxidative stress pathways of air pollution mediated toxicity: recent insights. Redox Biol 2020;34:101545.
- 36 An J, Tang W, Wang L, et al. Transcriptomics changes and the candidate pathway in human Macrophages induced by different PM₂₅ extracts. Environ Pollut 2021;289:117890.
- 37 Long Y-M, Yang X-Z, Yang Q-Q, et al. PM_{2.5} induces vascular permeability increase through activating MAPK/ERK signaling pathway and ROS generation. J Hazard Mater 2020;386:121659.
- 38 Eckert DJ, Malhotra A. Pathophysiology of adult obstructive sleep apnea. *Proc Am Thorac Soc* 2008;5:144–53.
- 39 Kim H, Kim W-H, Kim Y-Y, et al. Air pollution and central nervous system disease: a review of the impact of fine particulate matter on neurological disorders. Front Public Health 2020;8:575330.
- 40 Nuvolone D, Petri D, Voller F. The effects of ozone on human health. *Environ Sci Pollut Res Int* 2018;25:8074–88.
- 41 Genc S, Zadeoglulari Z, Fuss SH, et al. The adverse effects of air pollution on the nervous system. J Toxicol 2012;2012:782462.
- 42 Wang Y, Eliot MN, Wellenius GA. Short-term changes in ambient particulate matter and risk of stroke: a systematic review and metaanalysis. JAHA 2014;3:25103204.
- 43 Song J, Qu R, Sun B, et al. Associations of short-term exposure to fine particulate matter with neural damage biomarkers: a panel study of healthy retired adults. *Environ Sci Technol* 2022;56:7203–13.
- 44 Lo K, Chiang L-L, Hsu S-M, et al. Association of short-term exposure to air pollution with depression in patients with sleeprelated breathing disorders. Sci Total Environ 2021;786:147291.
- 45 Sharma HS, Ali SF, Hussain SM, et al. Influence of engineered nanoparticles from metals on the blood-brain barrier permeability, cerebral blood flow, brain edema and neurotoxicity. An experimental study in the rat and mice using biochemical and morphological approaches. J Nanosci Nanotechnol 2009;9:5055–72.
- 46 Ljubimova JY, Braubach O, Patil R, et al. Coarse particulate matter (Pm2. 5–10) in Los Angeles Basin air induces expression of inflammation and cancer biomarkers in rat brains. Sci Rep 2018;8:5708.
- 47 Krishnan V, Dixon-Williams S, Thornton JD. Where there is smoke... there is sleep apnea: exploring the relationship between smoking and sleep apnea. *Chest* 2014;146:1673–80.
- 48 Kolla BP, Foroughi M, Saeidifard F, et al. The impact of alcohol on breathing parameters during sleep: a systematic review and metaanalysis. Sleep Med Rev 2018;42:59–67.