

Probability Distributions and Moment Matching

ESS 575 Models for Ecological Data

N. Thompson Hobbs

January 31, 2017



Housekeeping

Errors in text:


```
http://www.stat.colostate.edu/~hooten/papers/pdf/  
Hobbs_Hooten_Bayesian_Models_2015_errata.pdf
```

Also in root directory of ESS_575_2017

Roadmap

- ▶ The rules of probability
 - ▶ conditional probability
 - ▶ independence
 - ▶ the law of total probability
- ▶ Factoring joint probabilities
- ▶ Probability distributions for discrete and continuous random variables
- ▶ Marginal distributions
- ▶ Moment matching

Motivation: A general approach to scientific research





Motivation: Why do we need to know this stuff?


Concept to be taught	Why do you need to understand this concept?
Conditional probability	It is the foundation for Bayes' Theorem and all inferences we will make.
The law of total probability	Basis for the denominator of Bayes' Theorem $[y]$
Factoring joint distributions	This is the procedure we will use to build models.
Independence	Allows us to simplify fully factored joint distributions.
Probability distributions	Our toolbox for representing uncertainty
Moments	The way we summarize distributions.
Marginal distributions	Bayesian inference is based on marginal distributions of unobserved quantities.
Moment matching	Allows us to embed the predictions of models into any statistical distribution.

Motivation: The essence of Bayes

Bayesian analysis is the *only* branch of statistics that treats all unobserved quantities as random variables. We seek to understand the characteristics of the probability distributions governing the behavior of these random variables.

Motivation: models of data

$$\mu_i = g(\boldsymbol{\theta}, x_i)$$



A model of the data describes our ideas about how the data arise.

Motivation: flexibility in analysis

Deterministic models

general linear
nonlinear
differential equations
difference equations
auto-regressive
occupancy
state-transition
integral-projection

Types of data

real numbers
non-negative real numbers
counts
0 to 1
0 or 1
counts in categories
proportions in categories

univariate and
multivariate


Motivation: flexibility in analysis

Probability model	Support for random variable
normal	real numbers
lognormal	non-negative real numbers
gamma	non-negative real numbers
beta	0 to 1 real numbers
Bernoulli	0 or 1
binomial	counts in 2 categories
Poisson	counts
multinomial	counts in > 2 categories
negative binomial	counts
Dirichlet	proportions in ≥ 2 categories
Cauchy	real numbers

Motivation: flexibility in analysis



$p = 0.5$








Work flow: probability distributions

- ▶ General properties and definitions (today)
 - ▶ discrete random variables
 - ▶ continuous random variables
- ▶ Specific distributions (in lab)
- ▶ Marginal distributions (Thursday and lab exercise)
- ▶ Moment matching (Thursday and lab exercise)



How will we use probability distributions?



Used to fit models to data, to represent uncertainty in processes and parameters, and to portray prior information

Used to make inferences about models.



Board work on probability distributions for discrete random variables

- ▶ Probability mass function, z , PMF (also called probability function, discrete destiny function)
 - ▶ notation $[z], f(z)$
 - ▶ $[z]$ is a function that returns the probability of a specific value of the random variable $= z$
 - ▶ Support of random variable z is defined as all values of z for which $[z] > 0$ and defined.
 - ▶ requirements to be a PMF
 - ▶ $[z] \geq 0$
 - ▶ $\sum_{z \in s} z = 1$, where s is the support of the random variable
 - ▶ moments of PMF
 - ▶ first moment, the expected value (or mean) $= E(z) = \mu = \sum_{z \in s} z[z]$, approximated from many (n) random draws from $[z]$ using $E(z) \simeq \frac{1}{n} \sum_{i=1}^n z_i$
 - ▶ second central moment, the variance $= E((z - \mu)^2) = \sigma^2 = \sum_{z \in s} (z - \mu)^2 [z]$, approximated from many (n) random draws from $[z]$ using $E((z - \mu)^2) \simeq \frac{1}{n} \sum_{i=1}^n (z_i - \mu)^2$
- ▶ cumulative distribution function for z
- ▶ quantile function for z



Board work on probability distributions for continuous random variables

- ▶ Probability density function, PDF, $[z]$
 - ▶ notation $[z], f(z), z \sim \text{normal}()$
 - ▶ $[z]$ gives the *probability density* of a specific value of the random variable $= z$.
 - ▶ Support of random variable z is defined as all values of z for which $[z] > 0$ and defined.
 - ▶ requirements
 - ▶ $[z] \geq 0$
 - ▶ $\int_{-\infty}^{\infty} [z] dz = 1$
 - ▶ $\Pr(a < z < b) = \int_a^b [z] dz$
 - ▶ What is probability density?
 - ▶ moments
 - ▶ first moment, the expected value (or mean) =
 $E(z) = \mu = \int_{-\infty}^{\infty} z[z] dx$, approximated from many (n) random draws from $[z]$ using $E(z) \simeq \frac{1}{n} \sum_{i=1}^n z_i$
 - ▶ second central moment, the variance =
 $E((z - \mu)^2) = \sigma^2 = \int_{-\infty}^{\infty} (z - \mu)^2 [z] dx$, approximated from many (n) random draws from $[z]$ using $E((z - \mu)^2) \simeq \frac{1}{n} \sum_{i=1}^n (z_i - \mu)^2$
- ▶ cumulative distribution function, CDF

dx should be dz



Marginal distributions of discrete random variables

	a_1	a_2	a_3	a_4	$\Pr(B) \downarrow$
b_1	$\frac{2}{20}$	0	$\frac{1}{20}$	$\frac{3}{20}$	$\frac{6}{20}$
b_2	0	$\frac{4}{20}$	0	$\frac{6}{20}$	$\frac{10}{20}$
b_3	$\frac{1}{20}$	$\frac{1}{20}$	$\frac{1}{20}$	0	$\frac{3}{20}$
b_4	0	0	0	$\frac{1}{20}$	$\frac{1}{20}$
$\Pr(A) \rightarrow$	$\frac{3}{20}$	$\frac{5}{20}$	$\frac{2}{20}$	$\frac{10}{20}$	$\frac{20}{20}$



Marginal distributions of discrete random variables

If we have a function $[A, B]$ specifying the joint probability of the discrete random variables A and B, then

$\sum_A [A, B]$ is the marginal probability of B
and

$\sum_B [A, B]$ is the marginal probability of A.

This same idea applies to any number of jointly distributed random variables. We simply sum over all but one.




Marginal distributions of continuous random variables

Exercise: If A and B are continuous random variables and we have a function $[A,B]$ that gives their joint probability density, what is the marginal distribution of A ? Of B ?



Marginal distributions of continuous random variables



Motivation: flexibility in analysis

Deterministic models

general linear
nonlinear
differential equations
difference equations
auto-regressive
occupancy
state-transition
integral-projection

univariate and
multivariate

Types of data

real numbers
non-negative real numbers
counts
0 to 1
0 or 1
counts in categories
proportions in categories

Motivation: flexibility in analysis


Probability model	Support for random variable
normal	all real numbers
lognormal	non-negative real numbers
gamma	non-negative real numbers
beta	0 to 1 real numbers
Bernoulli	0 or 1
binomial	counts in 2 categories
Poisson	counts
multinomial	counts in > 2 categories
negative binomial	counts
Dirichlet	proportions in ≥ 2 categories
Cauchy	non-negative real numbers

$$\mu_i = g(\theta, x_i)$$

A familiar approach


$$\boldsymbol{\theta} = (\beta_0, \beta_1)'$$

$$\mu_i = g(\boldsymbol{\theta}, x_i) = \beta_0 + \beta_1 x_i$$



The problem

All distributions have parameters:



α and β are parameters of the distribution of the random variable z .

Types of parameters

Parameter name	Function
intensity, centrality, location	sets position on x axis
shape	controls dispersion and skew
scale, dispersion parameter	shrinks or expands width
rate	scale ⁻¹

The problem

The normal and the Poisson are the only distributions for which the parameters of the distribution are the *same* as the moments. For all other distributions, the parameters are *functions* of the moments.

$$\alpha = m_1(\mu, \sigma^2)$$

$$\beta = m_2(\mu, \sigma^2)$$

We can use these functions to “match” the moments to the parameters.

Moment matching

$$\mu_i = g(\theta, x_i)$$

$$\alpha = m_1(\mu_i, \sigma^2)$$

$$\beta = m_2(\mu_i, \sigma^2)$$

$$[y_i | \alpha, \beta]$$

Moment matching the gamma distribution

The gamma distribution: $[z|\alpha, \beta] = \frac{n^\alpha z^{\alpha-1} e^{-\beta z}}{\Gamma(\alpha)}$

The mean of the gamma distribution is

$$\mu = \frac{\alpha}{\beta}$$

and the variance is

$$\sigma^2 = \frac{\alpha}{\beta^2}.$$

Discover functions for α and β in terms of μ and σ^2 .

Note: $\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} \frac{dt}{t}$

Moment matching the beta distribution

The beta distribution gives the probability density of random variables with support on $0, \dots, 1$.

$$[z|\alpha, \beta] = \frac{z^{\alpha-1}(1-z)^{\beta-1}}{B(\alpha, \beta)}$$

$$B = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}$$

$$\mu = \frac{\alpha}{\alpha + \beta}$$

$$\sigma^2 = \frac{\alpha \beta}{(\alpha + \beta)^2 (\alpha + \beta + 1)}$$

$$\alpha = \frac{\mu^2 - \mu^3 - \mu \sigma^2}{\sigma^2}$$

$$\beta = \frac{\mu - 2\mu^2 + \mu^3 - \sigma^2 + \mu \sigma^2}{\sigma^2}$$

You need some functions...

```
#BetaMomentMatch.R
# Function for parameters from moments
shape_from_stats <- function(mu, sigma){
  a <- (mu^2-mu^3-mu*sigma^2)/sigma^2
  b <- (mu-2*mu^2+mu^3-sigma^2+mu*sigma^2)/sigma^2
  shape_ps <- c(a,b)
  return(shape_ps)
}
# Functions for moments from parameters
beta.mean=function(a,b)a/(a+b)
beta.var = function(a,b)a*b/((a+b)^2*(a+b+1))
```

Problems continued

Do section on Moment Matching