

# THE PRIMITIVE AND DIAMOND SURFACES LOCALLY MINIMIZE THE VARIANCE OF GAUSS CURVATURE

HAO CHEN

**ABSTRACT.** We prove that Schwarz' Primitive and Diamond surfaces are local minimizers for the variance of Gauss curvature among local deformations within the space of Triply Periodic Minimal Surfaces of genus 3. This is done by viewing the branched values of the Gauss map as a point configuration on the sphere, and rewriting the variance of Gauss curvature as a product of two integrals involving exponentials of Green functions.

Triply Periodic Minimal Surfaces (TPMSs) are minimal surfaces in flat 3-tori. The genus of a TPMS is at least three. A TPMS of genus three (TPMSg3) has a Weierstrass parameterization defined on a branched double cover of the sphere with eight branch points. The branched covering map is provided by the Gauss map, so the branch points correspond to eight *flat points* on the TPMS (where the Gauss curvature vanishes). We refer the readers to [?] for general reference.

TPMSs find applications in natural sciences as a model for periodic bicontinuous structures [?]. In nature and in laboratories, the chance to observe Schwarz' Primitive (P), Diamond (D) and Schoen's Gyroid (G) surfaces is much higher than finding other TPMSs. Note that these surfaces belong to the same Bonnet family. They are therefore locally isometric, have the same branch points, and the same variance of Gauss curvature. For all the three surfaces, the branch points are vertices of a cube. Based on numerical evidence [?], it was conjectured that

**Conjecture.** *Schwarz' Primitive, Diamond, and Schoen's Gyroid surfaces have the minimal variance of Gauss curvature among all Triply Periodic Minimal Surfaces of genus 3.*

In this short note, we prove that the Primitive and Diamond are indeed local minimizers.

**Theorem.** *Schwarz' Primitive and Diamond surfaces are local minimizers for the variance of Gauss curvature among local deformations within the space of Triply Periodic Minimal Surfaces of genus 3.*

We will relax the problem to study the variance for the Gauss curvature defined on the 13-dimensional space  $V$  of configurations of eight points on the sphere up to global rotations. They correspond to the normal vectors at the eight flat points on a TPMSg3. For Diamond, Primitive, and Gyroid surfaces, the eight points are at the vertices of a cube. However, only a 5-dimensional variety in  $V$  correspond to TPMSg3s. In particular, the 5-dimensional subspace of antipodal configurations correspond to embedded TPMSg3s known as the *Meeks family*.

In [?], it was proved that, up to homotheties, the Primitive, Diamond, and Gyroid all belong to a unique locally rigid 5-parameter smooth family of pairwise non-homothetic triply periodic minimal surfaces. In particular, the family that

---

*Date:* May 2, 2026.

*2010 Mathematics Subject Classification.* Primary 53A10.

*Key words and phrases.* minimal surfaces.

contains the Diamond and the Primitive surfaces is the Meeks family. Schoen's Gyroid, however, does not belong to Meeks family.

At the cubic configuration, we compute the Hessian of the variance for the Gauss curvature on  $V$  and find, unfortunately, a 2-dimensional negative eigenspace generated by "twists". So the Hessian is indefinite, and the cubic configuration is not a local minimizer in  $V$ . Nevertheless, we are able to prove positive definiteness for the deformations preserving antipodality, so the cubic configuration is a local minimizer among antipodal deformations, thereby proving the Theorem.

The 5-parameter family of local deformations of the Gyroid is not fully understood. Among the known deformations of the Gyroid, it is indeed a local minimizer. But in the presence of the negative eigenspace, we can not rule out a possibility that it might not be a local minimizer along other deformations.

**Acknowledgement.** The author thanks Chengjian Yao and Thierry De Pauw for very helpful and inspiring discussions.

## 1. REFORMULATION

**1.1. Weierstrass parameterization.** Given a TPMSg3, let  $p_1, \dots, p_8$  be (the stereographic projection of) the Gauss map at the eight flat points. Consider the hyperelliptic Riemann surface  $\Sigma$  of genus three defined by

$$w^2 = P(z) = \prod_{i=1}^8 (z - p_i).$$

It can be seen as a branched double cover of the Riemann sphere  $\mathbb{S}^2 \simeq \mathbf{CP}^1$  with eight branch points. Then we have the following *Weierstrass parameterization* for the TPMS (up to translations)

$$(1) \quad \Sigma \ni (z, w) \mapsto \operatorname{Re} \int^{(z, w)} \frac{(1 - z^2, i(1 + z^2), 2z)}{w} e^{i\vartheta} dz.$$

Switching the two branches by  $(z, w) \mapsto (z, -w)$  correspond to the inversion with respect to any of the eight flat points. The parameter  $\vartheta$  is the *Bonnet angle*; changing  $\vartheta$  gives a locally isometric deformation of the surface known as *Bonnet rotation*. Two surfaces whose  $\vartheta$  differ by  $\pi/2$  are called *conjugate pairs*.

For the minimal surface to be triply periodic and globally well-defined, one needs to solve the period problem (aka, *close the periods*). More precisely, one finds  $p_i$ 's and  $\vartheta$  so that the real parts of the integrals over closed cycles on  $\Sigma$  form a three dimensional lattice. The period problem involves three real equations for each of the three coordinates, hence nine real equations in total. Since we have 17 real parameters ( $p_i$ 's and  $\vartheta$ ), the set of TPMSg3s form a 8-dimensional variety up to translations, or 5-dimensional variety up to homotheties.

Closing the periods is usually very difficult. But Meeks proved in [?] that, when the branch points form four antipodal pairs, there exists a conjugate pair embedded TPMSg3s. Up to homotheties, these TPMSg3s form a 5-dimensional manifold termed *Meeks family*.

In this paper, we will not insist on closing the periods. The variance of Gauss curvature is invariant under local isometries such as Bonnet rotations. So it is a function defined on a 13-dimensional space of configurations of eight points on the sphere up to rotations. The Meeks surfaces correspond to the 5-dimensional subspace of antipodal configurations, for which we know the existence of  $\vartheta$  that closes the periods. Our main Theorem is equivalent to that the cubic configuration is a local minimizer of the Gauss variance on this 5-dimensional subspace.

**1.2. Variance of Gauss curvature.** More precisely, we want to minimize the (dimensionless) variance of Gauss curvature, defined by

$$\sigma(K) = \frac{\langle K^2 \rangle - \langle K \rangle^2}{\langle K \rangle^2} = \left[ \frac{\int_{\Sigma} K^2 dA}{\int_{\Sigma} dA} - \left( \frac{\int_{\Sigma} K dA}{\int_{\Sigma} dA} \right)^2 \right] / \left( \frac{\int_{\Sigma} K dA}{\int_{\Sigma} dA} \right)^2.$$

By Gauss-Bonnet,  $\int_{\Sigma} K dA$  is a topological invariant. It is then equivalent to minimize

$$(2) \quad \int_{\Sigma} K^2 dA \times \int_{\Sigma} dA.$$

From the Weierstrass parameterization, we compute

$$dA = \frac{(1 + |z|^2)^2}{|w|^2} dx \wedge dy, \quad K = -4 \frac{|w|^2}{(1 + |z|^2)^4},$$

where  $z = x + yi$ . Indeed, one verifies that

$$\int_{\Sigma} K dA = -4 \int_{\Sigma} \frac{dx \wedge dy}{(1 + |z|^2)^2} = -8\pi$$

as expected. The area element and the Gauss curvature are invariant after switching the branches. It then suffices to compute the invariance on one of the branch. So (2) becomes (up to a constant factor)

$$(3) \quad \int_{\mathbf{CP}^1} \frac{|P(z)|}{(1 + |z|^2)^6} dx \wedge dy \times \int_{\mathbf{CP}^1} \frac{(1 + |z|^2)^2}{|P(z)|} dx \wedge dy.$$

Note that the second integral is improper: The integrand is not defined at the zeros of  $P(z)$ . So the integral is defined as the limit

$$\int_{\mathbf{CP}^1} \frac{(1 + |z|^2)^2}{|P(z)|} dx \wedge dy = \lim_{\varepsilon \rightarrow 0} \int_{\mathbf{CP}^1 \setminus \cup_i D_{\varepsilon}^i} \frac{(1 + |z|^2)^2}{|P(z)|} dx \wedge dy,$$

where  $D_{\varepsilon}^i$  denote the  $\varepsilon$ -disks around  $p_i$  under the Fubini–Study metric. Fortunately, the limit converges.

**1.3. Fubini–Study metric.** Note that

$$dA_{\text{FS}} = \frac{4 dx \wedge dy}{(1 + |z|^2)^2}$$

is the area element for the Fubini–Study metric on  $\mathbf{CP}^1$ , which is (normalized to) exactly the standard metric on the unit sphere  $\mathbb{S}^2$ . Moreover, we can identify  $P(z)$  to a global section  $s$  of the holomorphic line bundle  $\mathcal{O}(8) = \mathcal{O}(1)^{\otimes 8}$  over  $\mathbf{CP}^1$ , and

$$\|s\|_{\text{FS}} = \frac{|P(z)|}{(1 + |z|^2)^4}$$

is its point-wise Fubini–Study norm. See, for instance, [?]. Then (3) becomes (up to a constant factor)

$$(4) \quad J_8 := \int_{\mathbf{CP}^1} \|s\|_{\text{FS}} dA_{\text{FS}} \times \int_{\mathbf{CP}^1} \|s\|_{\text{FS}}^{-1} dA_{\text{FS}}.$$

Note again that the second integral is improper. It is defined as the convergent limit of the integral over  $\mathbf{CP}^1 \setminus \cup_i D_{\varepsilon}^i$  as  $\varepsilon \rightarrow 0$ .

**1.4. Lelong–Poincaré Equation.** Define

$$u = \log \|s\|_{\text{FS}}.$$

Note that the Fubini–Study metric on  $\mathcal{O}(8)$  is Hermitian. One then easily verifies the Lelong–Poincaré Equation

$$\Delta u = 2\pi \sum_{i=1}^8 \delta(z - p_i) - 4 \frac{4}{(1 + |z|^2)^2},$$

or, using the Laplacian under the Fubini–Study metric,

$$\Delta_{\text{FS}} u = \frac{(1 + |z|^2)^2}{4} \Delta u = 2\pi \sum_{i=1}^8 \delta_i(z) - 4,$$

where

$$\delta_i(z) = \frac{(1 + |z|^2)^2}{4} \delta(z - p_i)$$

denote the Dirac delta function under the Fubini–Study metric centered at  $p_i$ .

The solution is given by

$$u = 2\pi \sum_{i=1}^8 G_i(z),$$

where  $G_i(z)$  is the Green function solving

$$(5) \quad \Delta_{\text{FS}} G_i(z) = \delta_i(z) - \frac{1}{4\pi}.$$

On the unit sphere  $\mathbb{S}^2 \simeq \mathbf{CP}^1$ , we have

$$G_i(z) = -\frac{1}{2\pi} \log \sin \frac{\text{dist}(z, p_i)}{2} + C$$

where  $\text{dist}(z, p_i)$  is the spherical distance between  $z$  and  $p_i$ . Then (4) becomes

$$(6) \quad J_8 = \int_{\mathbb{S}^2} \exp(u) \, dA_{\mathbb{S}^2} \times \int_{\mathbb{S}^2} \exp(-u) \, dA_{\mathbb{S}^2},$$

where  $dA_{\mathbb{S}^2}$  is the standard spherical area element on the unit sphere. Note that (6) does not depend on the constant  $C$  in  $G_i$ , so we assume  $C = 0$  from now on. Note again that the second integral is improper and is defined as the convergent limit

$$\int_{\mathbb{S}^2} \exp(-u) \, dA_{\mathbb{S}^2} = \lim_{\varepsilon \rightarrow 0} \int_{\Omega_\varepsilon} \exp(-u) \, dA_{\mathbb{S}^2},$$

where

$$\Omega_\varepsilon = \mathbb{S}^2 \setminus \cup_{i=1}^n D_\varepsilon^i.$$

The Conjecture would be proved if  $J_8$  is minimized when  $p_i \in \mathbb{S}^2$  are the vertices of a cube.

*Remark 1.* With this reformulation, we actually expand our consideration beyond TPMSg3s. We are now considering all minimal surfaces with the Weierstrass parameterization (1). In particular, we do not need the period problems to be solved, not to mention the embeddedness.

**1.5. Generalization.** The analysis above can be generalized as follows. Fix  $n$  points  $p_1, \dots, p_n$ , and define the polynomial

$$P(z) = \prod_{i=1}^n (z - p_i).$$

Consider the product of integrals

$$\int_{\mathbf{CP}^1} \frac{|P(z)|}{(1 + |z|^2)^{\frac{n}{2}+2}} dx \wedge dy \times \int_{\mathbf{CP}^1} \frac{(1 + |z|^2)^{\frac{n}{2}-2}}{|P(z)|} dx \wedge dy.$$

Note that

$$\|s\|_{\text{FS}} = \frac{|P(z)|}{(1 + |z|^2)^{n/2}}$$

is the point-wise Fubini–Study norm of  $P(z)$  as a global section  $s$  of  $\mathcal{O}(n)$  over  $\mathbf{CP}^1$ . Then the product can be rewritten (up to a constant factor) into the form

$$J_n := \int_{\mathbf{CP}^1} \|s\|_{\text{FS}} dA_{\text{FS}} \times \int_{\mathbf{CP}^1} \|s\|_{\text{FS}}^{-1} dA_{\text{FS}}.$$

Now the Lelong–Poincaré Equation is

$$\Delta_{\text{FS}} u = 2\pi \sum_{i=1}^n \delta_i(z) - n/2,$$

which is solved by

$$u = 2\pi \sum_{i=1}^n G_i(z).$$

So

$$J_n = \int_{\mathbb{S}^2} \exp(u) dA_{\mathbb{S}^2} \times \int_{\mathbb{S}^2} \exp(-u) dA_{\mathbb{S}^2}.$$

And we could ask the following problem

**Problem.** *What are the positions of  $p_1, \dots, p_n$  that minimize  $J_n$ ?*

When  $n = 8$ , this is exactly the problem of finding branch values to minimize the variance of the Gauss curvature.

## 2. GRADIENT AND HESSIAN

**2.1. The gradient.** Recall that

$$\Omega_\varepsilon = \mathbb{S}^2 \setminus \cup_{i=1}^n D_\varepsilon^i,$$

where  $D_\varepsilon^i$  denote the  $\varepsilon$ -disks around  $p_i$  under the spherical metric. Define

$$I_\varepsilon^\pm = \int_{\Omega_\varepsilon} \exp(\pm u) dA_{\mathbb{S}^2},$$

and two probability measure  $d\mu_\varepsilon^\pm = \rho_\varepsilon^\pm dA_{\mathbb{S}^2}$  on  $\Omega_\varepsilon$ , where

$$\rho_\varepsilon^\pm = \frac{\exp(\pm u)}{\int_{\Omega_\varepsilon} \exp(\pm u) dA_{\mathbb{S}^2}} = \frac{\exp(\pm u)}{I_\varepsilon^\pm}.$$

We compute

$$\begin{aligned} \langle \nabla_{p_i} \log I_\varepsilon^\pm, v_i \rangle &= \frac{\langle \nabla_{p_i} I_\varepsilon^\pm, v_i \rangle}{I_\varepsilon^\pm} \\ &= \frac{1}{I_\varepsilon^\pm} \left( \int_{\Omega_\varepsilon} \langle \pm 2\pi \nabla_{p_i} G_i, v_i \rangle \exp(\pm u) dA_{\mathbb{S}^2} + \oint_{\partial D_\varepsilon^i} \exp(\pm u) \langle n_i, \tilde{v}_i \rangle ds \right) \\ &= \int_{\Omega_\varepsilon} \langle \pm 2\pi \nabla_{p_i} G_i, v_i \rangle \rho_\varepsilon^\pm dA_{\mathbb{S}^2} + \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm ds \end{aligned}$$

where  $v_i \in T_{p_i} \mathbb{S}^2$ ,  $n_i$  is the unit normal vectors on the boundary of  $D_\varepsilon^i$  pointing to the interiors of the disks, and  $\tilde{v}_i$  is a Killing field such that  $\tilde{v}_i(p_i) = v_i$ . So for

$$v = (v_1, \dots, v_8) \in \bigoplus_{i=1}^8 T_{p_i} \mathbb{S}^2,$$

we have

$$(7) \quad \langle \nabla_p \log I_\varepsilon^\pm, v \rangle = \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu_\varepsilon^\pm + \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm ds$$

Note that the formula does not depend on the choice of the Killing fields  $\tilde{v}_i$ . To see this, assume another choice  $\tilde{v}'_i$ . Then  $\tilde{w}_i = \tilde{v}'_i - \tilde{v}_i$  is a Killing field that vanishes at  $p_i$ , which is given as a rotation around  $p_i$ . Consequently, we have  $\langle \mp 2\pi \nabla_z G_i, \tilde{w}_i \rangle = 0$  and  $\langle n_i, \tilde{w}_i \rangle = 0$ .

Note also that, when  $\tilde{v}_i = \tilde{v}_*$  is the same Killing field for all  $i$ , then

$$\begin{aligned} \langle \nabla_p \log I_\varepsilon^\pm, v \rangle &= \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_* \rangle d\mu_\varepsilon^\pm + \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_* \rangle \rho_\varepsilon^\pm ds \\ &= \int_{\Omega_\varepsilon} \langle \mp \nabla_z u, \tilde{v}_* \rangle d\mu_\varepsilon^\pm + \int_{\Omega_\varepsilon} \langle \pm \nabla_z u, \tilde{v}_* \rangle d\mu_\varepsilon^\pm = 0 \end{aligned}$$

So a deformation along a Killing field does not change  $I_\varepsilon^\pm$ .

Recall that  $2\pi G_i = \log \sin(\frac{1}{2} \text{dist}(z, p_i))$ , so

$$2\pi \nabla_z G_i \sim \frac{\hat{w}_i}{\text{dist}(z, p_i)}, \quad \rho_\varepsilon^\pm \sim \text{dist}(z, p_i)^{\pm 1}$$

as  $z \rightarrow p_i$ , where  $\hat{w}_i = \nabla_z \text{dist}(z, p_i)$ . It is then obvious that (7) converges as  $\varepsilon \rightarrow 0$  for  $\mu_\varepsilon^\pm$ . As for  $\mu_\varepsilon^-$ , in the standard spherical coordinate with the pole  $p_i$ , polar angle  $\theta$  and azimuth angle  $\phi$ , the integrands

$$\begin{aligned} \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu_\varepsilon^- &\sim \frac{2}{\theta I^-} \langle \hat{w}_i(\theta, \phi), \tilde{v}_i(\theta, \phi) \rangle d\theta d\phi, \\ \langle n_i, \tilde{v}_i \rangle \rho_\varepsilon^- ds &\sim \frac{-2}{I^-} \langle \hat{w}_i(\varepsilon, \phi), \tilde{v}_i(\varepsilon, \phi) \rangle d\phi \end{aligned}$$

as  $\theta, \varepsilon \rightarrow 0$ . The integrals in (7) then converge as  $\varepsilon \rightarrow 0$  because

$$\lim_{\theta \rightarrow 0} \int_0^{2\pi} \langle \hat{w}_i(\theta, \phi), \tilde{v}_i(\theta, \phi) \rangle d\phi \rightarrow 0.$$

More specifically, we have

$$\langle \nabla_p \log I^\pm, v \rangle := \lim_{\varepsilon \rightarrow 0} \langle \nabla_p \log I_\varepsilon^\pm, v \rangle = \int_{\mathbb{S}^2} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu^\pm.$$

As a sanity check, one easily verifies that this vanishes when  $\tilde{v}_i = \tilde{v}_*$  is the same Killing field for all  $i$ .

Finally, by symmetry, one easily verifies that  $\langle \nabla_p \log I_\varepsilon^\pm, v \rangle = 0$  for any  $v$  when  $p_1, \dots, p_8$  are the vertices of the cube. So the cube is indeed a critical point of  $J_8$ .

**2.2. The Hessian.** We compute,

$$\begin{aligned} \langle \nabla_{p_i} \rho_\varepsilon^\pm, v_i \rangle &= \frac{\langle \pm 2\pi \nabla_{p_i} G_i, v_i \rangle \exp(\pm u)}{I_\varepsilon^\pm} - \frac{\exp(\pm u) \langle \nabla_{p_i} I_\varepsilon^\pm, v_i \rangle}{(I_\varepsilon^\pm)^2} \\ &= \rho_\varepsilon^\pm \left( \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle - \int_{\Omega_\varepsilon} \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu_\varepsilon^\pm - \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm ds \right) \end{aligned}$$

The Hessian

$$\begin{aligned}
(8) \quad \nabla_p^2 \log I_\varepsilon^\pm(v, v) &= \langle \nabla_p \langle \nabla_p \log I_\varepsilon^\pm, v \rangle, v \rangle \\
&= \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \nabla_{p_i} \langle \pm 2\pi \nabla_{p_i} G_i, v_i \rangle, v_i \rangle \rho_\varepsilon^\pm \, dA_{\mathbb{S}^2} \\
(9) \quad &+ \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \pm 2\pi \nabla_{p_i} G_i, v_i \rangle \sum_{j=1}^n \langle \nabla_{p_j} \rho_\varepsilon^\pm, v_j \rangle \, dA_{\mathbb{S}^2} \\
(10) \quad &+ \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \sum_{j=1}^n \langle \nabla_{p_j} \rho_\varepsilon^\pm, v_j \rangle \, ds \\
(11) \quad &+ \sum_{j=1}^n \oint_{\partial D_\varepsilon^j} \langle n_j, \tilde{v}_j \rangle \sum_{i=1}^n \langle \pm 2\pi \nabla_{p_i} G_i, v_i \rangle \rho_\varepsilon^\pm \, ds \\
&+ \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \langle \nabla_z \rho_\varepsilon^\pm, \tilde{v}_i \rangle \, ds
\end{aligned}$$

We expand each term

$$\begin{aligned}
(8) &= \int_{\Omega_\varepsilon} \sum_{i=1}^n \pm 2\pi \nabla_z^2 G_i(\tilde{v}_i, \tilde{v}_i) \, d\mu_\varepsilon^\pm \\
(9) &= \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle \sum_{j=1}^n \langle \nabla_{p_j} \rho_\varepsilon^\pm, v_j \rangle \, dA_{\mathbb{S}^2} \\
&= \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle \sum_{j=1}^n \langle \mp 2\pi \nabla_z G_j, \tilde{v}_j \rangle \, d\mu_\varepsilon^\pm \\
&\quad - \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle \, d\mu_\varepsilon^\pm \int_{\Omega_\varepsilon} \sum_{j=1}^n \langle \mp 2\pi \nabla_z G_j, \tilde{v}_j \rangle \, d\mu_\varepsilon^\pm \\
&\quad - \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle \, d\mu_\varepsilon^\pm \sum_{j=1}^n \oint_{\partial D_\varepsilon^j} \langle n_j, \tilde{v}_j \rangle \rho_\varepsilon^\pm \, ds \\
(10) &= \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \sum_{j=1}^n \langle \mp 2\pi \nabla_z G_j, \tilde{v}_j \rangle \rho_\varepsilon^\pm \, ds \\
&\quad - \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm \, ds \int_{\Omega_\varepsilon} \sum_{j=1}^n \langle \mp 2\pi \nabla_z G_j, \tilde{v}_j \rangle \, d\mu_\varepsilon^\pm \\
&\quad - \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm \, ds \sum_{j=1}^n \oint_{\partial D_\varepsilon^j} \langle n_j, \tilde{v}_j \rangle \rho_\varepsilon^\pm \, ds,
\end{aligned}$$

and

$$(11) = \sum_{j=1}^n \int_{\partial D_\varepsilon^j} \langle n_j, \tilde{v}_j \rangle \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm \, ds.$$

Putting them together gives

$$(12) \quad \nabla_p^2 \log I_\varepsilon^\pm(v, v) = \int_{\Omega_\varepsilon} \left( \sum_{i=1}^n \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle \right)^2 d\mu_\varepsilon^\pm$$

$$(13) \quad + \int_{\Omega_\varepsilon} \sum_{i=1}^n \pm 2\pi \nabla_z^2 G_i(\tilde{v}_i, \tilde{v}_i) d\mu_\varepsilon^\pm$$

$$(14) \quad - \left( \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu_\varepsilon^\pm + \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm ds \right)^2$$

$$(15) \quad + \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \sum_{j=1}^n \langle \mp 2\pi \nabla_z G_j, \tilde{v}_j \rangle \rho_\varepsilon^\pm ds$$

$$(16) \quad + \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \sum_{j=1}^n \langle \mp 2\pi \nabla_z G_j, \tilde{v}_j - \tilde{v}_i \rangle \rho_\varepsilon^\pm ds$$

Note again the independence to the choice of the Killing fields  $\tilde{v}_i$  (by the same argument as before) and, when  $\tilde{v}_i = \tilde{v}_*$  is the same Killing field for all  $i$ ,

$$\begin{aligned} \nabla_p^2 \log I_\varepsilon^\pm(v, v) &= \int_{\Omega_\varepsilon} \langle \nabla_z u, \tilde{v}_* \rangle^2 d\mu_\varepsilon^\pm + \int_{\Omega_\varepsilon} \pm \nabla_z^2 u(\tilde{v}_*, \tilde{v}_*) d\mu_\varepsilon^\pm \\ &\quad + \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_* \rangle \langle \mp \nabla_z u, \tilde{v}_* \rangle \rho_\varepsilon^\pm ds = 0. \end{aligned}$$

The boundary term (16) converges to 0 because  $D_\varepsilon^i$  only contains the singularity of  $G_j$  when  $j = i$ . For the same reason, in the limit  $\varepsilon \rightarrow 0$ , we may rewrite the “coupling” boundary term (15) into the form

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \sum_{i=1}^n \oint_{\partial D_\varepsilon^i} \langle n_i, \tilde{v}_i \rangle \langle \mp 2\pi \nabla_z G_i, \tilde{v}_i \rangle \rho_\varepsilon^\pm ds \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{i=1}^n \left[ - \int_{\Omega_\varepsilon} \langle \nabla_z u, \tilde{v}_i \rangle \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu_\varepsilon^\pm \mp \int_{\Omega_\varepsilon} 2\pi \nabla_z^2 G_i(\tilde{v}_i, \tilde{v}_i) d\mu_\varepsilon^\pm \right], \end{aligned}$$

where the second term cancels with (13).

We have seen that the second term in (14) converges to 0 as  $\varepsilon \rightarrow 0$ . At a critical point, its first term (14) vanishes. If this is the case, we have

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \nabla_p^2 \log I_\varepsilon^\pm(v, v) \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\Omega_\varepsilon} \left( \sum_{i=1}^n \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle \right)^2 d\mu_\varepsilon^\pm - \int_{\Omega_\varepsilon} \sum_{i=1}^n \langle \nabla_z u, \tilde{v}_i \rangle \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu_\varepsilon^\pm \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\Omega_\varepsilon} \sum_{i=1}^n \sum_{j=1}^n \langle 2\pi \nabla_z G_j, \tilde{v}_j - \tilde{v}_i \rangle \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu_\varepsilon^\pm. \end{aligned}$$

When  $i \neq j$ , the convergence follows from the same argument for the convergence of the gradient (7). The possible divergence with  $i = j$  is ruled out by  $\tilde{v}_j - \tilde{v}_i$ . In the following, we will simply write

$$\nabla_p^2 \log I^\pm(v, v) = \int_{\mathbb{S}^2} \sum_{i=1}^n \sum_{j=1}^n \langle 2\pi \nabla_z G_j, \tilde{v}_j - \tilde{v}_i \rangle \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu^\pm.$$



### 3. LOCAL MINIMIZER

Without loss of generality, we assume that the vertices of the cube, in the standard coordinates  $(\theta, \phi)$  of the sphere, are as follows:

$$\begin{aligned} p_1 &= (\theta_0, 0), & p_5 &= (\pi - \theta_0, 0), \\ p_2 &= (\theta_0, \pi/2), & p_6 &= (\pi - \theta_0, \pi/2), \\ p_3 &= (\theta_0, \pi), & p_7 &= (\pi - \theta_0, \pi), \\ p_4 &= (\theta_0, 3\pi/2), & p_8 &= (\pi - \theta_0, 3\pi/2). \end{aligned}$$

where  $\theta_0 = \arccos(1/\sqrt{3})$ .

**3.1. Decomposition into irreducible representations.** We have seen that the Hessian vanishes on Killing fields, compatible with the fact the  $J$  is invariant under global rotations. So we study the Hessian on the 13-dimensional space

$$V = \left( \bigoplus_{i=1}^8 T_{p_i} \mathbb{S}^2 \right) / \text{SO}(3)$$

at the cubic configuration. For this purpose, we decompose  $V$  into irreducible representations (irreps) of the symmetry group of the cube, as follows:

- The 3-dimensional space  $T_{1u}$  generated by slides (that moves the center of mass). One generator is given by

$$v_i = \frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{1, 2, 5, 6\}, \quad v_i = -\frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{3, 4, 7, 8\}.$$

- The 2-dimensional space  $E_g$  generated by orbit of

$$v_i = \frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{1, 3, 5, 7\}, \quad v_i = -\frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{2, 4, 6, 8\}.$$

Every face remain rectangular under these deformations.

- The 2-dimensional space  $E_u$  generated by orbit of

$$v_i = \frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{1, 2, 3, 4\}, \quad v_i = -\frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{5, 6, 7, 8\}.$$

The face centers of the cube remain symmetry centers under these deformations.

- The 3-dimensional space  $T_{2g}$  generated by the orbit of

$$v_i = \frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{1, 2, 7, 8\}, \quad v_i = -\frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{3, 4, 5, 6\}.$$

- The 3-dimensional space  $T_{2u}$  generated by orbit of

$$v_i = \frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{1, 3, 6, 8\}, \quad v_i = -\frac{\partial \phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{2, 4, 5, 7\}.$$

The Hessian is invariant under the action of  $G$ . By Schur's lemma, it must be a multiple of identity on each of the irreps. So we only need to verify that the Hessian is non-negative on one generator for each of the irreps.

**3.2. The definiteness on  $T_{1u}$ ,  $E_u$  and  $E_g$ .** We have chosen the generator vectors, intentionally, with the following property. The eight vertices are partitioned into two subsets  $A$  and  $B$ , with four vertices each, such that vertices in  $A$  extend to the Killing field  $\partial_\phi$ , while vertices in  $B$  extend to  $-\partial_\phi$ . In this case, we have

$$\begin{aligned} \nabla_p^2 \log I^\pm(v, v) &= \int_{\mathbb{S}^2} \sum_{i=1}^n \sum_{j=1}^n \langle 2\pi \nabla_z G_j, \tilde{v}_j - \tilde{v}_i \rangle \langle 2\pi \nabla_z G_i, \tilde{v}_i \rangle d\mu^\pm \\ &= -4 \int_{\mathbb{S}^2} \sum_{i \in A} \sum_{j \in B} (2\pi \partial_\phi G_j)(2\pi \partial_\phi G_i) d\mu^\pm = -16\pi^2 \int_{\mathbb{S}^2} \partial_\phi G_A \partial_\phi G_B d\mu^\pm, \end{aligned}$$

where

$$G_A = \sum_{i \in A} G_i, \quad G_B = \sum_{i \in B} G_i.$$

To study the definiteness of the Hessian, we need to first study the sign of

$$H_{ij}^\pm = \int_{\mathbb{S}^2} \partial_\phi G_i \partial_\phi G_j d\mu^\pm.$$

When  $p_i$  and  $p_j$  are antipodal (e.g.  $(i, j) = (1, 7)$ ) or on a horizontal face diagonal (e.g.  $(i, j) = (1, 3)$ ), one easily sees that the integrand  $\partial_\phi G_i \partial_\phi G_j \leq 0$  everywhere on the sphere, so  $H_{ij}^\pm$  is negative. When  $p_i$  and  $p_j$  are vertical neighbors (e.g.  $(i, j) = (1, 5)$ ), one easily sees that the integrand  $\partial_\phi G_i \partial_\phi G_j \geq 0$  everywhere on the sphere, so  $H_{ij}^\pm$  is positive. Otherwise, the following lemma would be useful:

**Lemma.** *Let  $f(\theta, \phi)$  be a function defined on  $\mathbb{S}^2$  that is odd in  $\phi$ , and  $\rho$  be a positive function on  $\mathbb{S}^2$  with*

$$\rho(\theta, \phi) = \rho(\theta, \phi + \frac{\pi}{2}) = \rho(\pi - \theta, \phi).$$

*Assume that  $f(\theta, \phi)$  is negative and strictly monotonically increasing in  $\phi$  on the interval  $(0, \pi)$ . Then the integrals*

$$\begin{aligned} K_1 &= \int_{\mathbb{S}^2} f(\theta, \phi) f(\theta, \phi - \frac{\pi}{2}) \rho dA_{\mathbb{S}^2} < 0, \\ K_2 &= \int_{\mathbb{S}^2} f(\theta, \phi) f(\pi - \theta, \phi - \frac{\pi}{2}) \rho dA_{\mathbb{S}^2} < 0. \end{aligned}$$

*Moreover, if  $f(\theta, \phi) - f(\pi - \theta, \phi)$  is negative and monotonically increasing in  $\phi$  on the interval  $(0, \pi)$  for  $0 < \theta < \frac{\pi}{2}$ , then  $K_1 < K_2 < 0$ .*

*Proof.* We decompose the sphere into four quadrants. Then

$$\begin{aligned} K_1 &= \int_0^\pi \sin \theta d\theta \int_0^{\frac{\pi}{2}} \rho d\phi (f(\theta, \phi) f(\theta, \phi - \frac{\pi}{2}) + f(\theta, \phi + \frac{\pi}{2}) f(\theta, \phi) \\ &\quad + f(\theta, \phi + \pi) f(\theta, \phi + \frac{\pi}{2}) + f(\theta, \phi - \frac{\pi}{2}) f(\theta, \phi + \pi)) \\ &= \int_0^\pi \sin \theta d\theta \int_0^{\frac{\pi}{2}} \rho d\phi (f(\theta, \phi) + f(\theta, \phi + \pi)) (f(\theta, \phi - \frac{\pi}{2}) + f(\theta, \phi + \frac{\pi}{2})) \\ &= \int_0^\pi \sin \theta d\theta \int_0^{\frac{\pi}{2}} \rho d\phi (f(\theta, \phi) - f(\theta, \pi - \phi)) (-f(\theta, \frac{\pi}{2} - \phi) + f(\theta, \frac{\pi}{2} + \phi)) \end{aligned}$$

For  $0 < \phi < \frac{\pi}{2}$ , by the assumed monotonicity, we have  $f(\phi) < f(\pi - \phi) < 0$  and  $g(\frac{\pi}{2} - \phi) < g(\frac{\pi}{2} + \phi) < 0$ . So  $K_1$  is negative. On the other hand,

$$\begin{aligned}
K_2 &= \int_0^{\frac{\pi}{2}} \sin \theta d\theta \int_0^{\frac{\pi}{2}} \rho d\phi (f(\theta, \phi) f(\pi - \theta, \phi - \frac{\pi}{2}) + f(\pi - \theta, \phi) f(\theta, \phi - \frac{\pi}{2}) \\
&\quad + f(\theta, \phi + \frac{\pi}{2}) f(\pi - \theta, \phi) + f(\pi - \theta, \phi + \frac{\pi}{2}) f(\theta, \phi) \\
&\quad + f(\theta, \phi + \pi) f(\pi - \theta, \phi + \frac{\pi}{2}) + f(\pi - \theta, \phi + \pi) f(\theta, \phi + \frac{\pi}{2}) \\
&\quad + f(\theta, \phi - \frac{\pi}{2}) f(\pi - \theta, \phi + \pi) + f(\pi - \theta, \phi - \frac{\pi}{2}) f(\theta, \phi + \pi)) \\
&= \int_0^{\frac{\pi}{2}} \sin \theta d\theta \int_0^{\frac{\pi}{2}} \rho d\phi \\
&\quad \left[ (f(\pi - \theta, \phi) + f(\pi - \theta, \phi + \pi)) (f(\theta, \phi - \frac{\pi}{2}) + f(\theta, \phi + \frac{\pi}{2})) \right. \\
&\quad \left. + (f(\theta, \phi) + f(\theta, \phi + \pi)) (f(\pi - \theta, \phi - \frac{\pi}{2}) + f(\pi - \theta, \phi + \frac{\pi}{2})) \right] \\
&= \int_0^{\frac{\pi}{2}} \sin \theta d\theta \int_0^{\frac{\pi}{2}} \rho d\phi \\
&\quad \left[ (f(\pi - \theta, \phi) - f(\pi - \theta, \pi - \phi)) (-f(\theta, \frac{\pi}{2} - \phi) + f(\theta, \phi + \frac{\pi}{2})) \right. \\
&\quad \left. + (f(\theta, \phi) - f(\theta, \pi - \phi)) (-f(\pi - \theta, \frac{\pi}{2} - \phi) + f(\pi - \theta, \phi + \frac{\pi}{2})) \right]
\end{aligned}$$

and the integrand is negative by the same argument as above.

Moreover,

$$\begin{aligned}
K_1 - K_2 &= \int_0^{\frac{\pi}{2}} \sin \theta d\theta \int_0^{\frac{\pi}{2}} \rho d\phi \\
&\quad (f(\theta, \phi) - f(\pi - \theta, \phi) - f(\theta, \pi - \phi) + f(\pi - \theta, \pi - \phi)) \\
&\quad (-f(\theta, \frac{\pi}{2} - \phi) + f(\pi - \theta, \frac{\pi}{2} - \phi) + f(\theta, \frac{\pi}{2} + \phi) - f(\pi - \theta, \frac{\pi}{2} + \phi)).
\end{aligned}$$

Again the integrand is negative if  $f(\theta, \phi) - f(\pi - \theta, \phi)$  is negative and monotonically increasing in  $\phi$  on the interval  $(0, \pi)$ . So  $K_1 < K_2$ .  $\square$

Using this lemma, one easily proves that if  $p_i$  and  $p_j$  are horizontal neighbors (e.g.  $(i, j) = (1, 2)$ ) or on a vertical face diagonal (e.g.  $(i, j) = (1, 6)$ ), we have  $H_{ij}^\pm < 0$ . Consequently, by

$$\nabla_p^2 \log I^\pm(v, v) = -16\pi^2 \sum_{i \in A} \sum_{j \in B} H_{ij}^\pm$$

we can already conclude that the Hessian is positive definite on the irreps  $T_{1u}$ ,  $E_g$ , but negative definite on  $E_u$ .

**3.3. The definiteness on  $T_{2u}$ .** For  $T_{2u}$  we place the cube so that a pair of antipodal vertices are at the north and the south pole. So

$$\begin{aligned}
p_1 &= (0, 0), & p_5 &= (\pi, 0), \\
p_2 &= (\theta_0, 0), & p_6 &= (\pi - \theta_0, \pi/3), \\
p_3 &= (\theta_0, 2\pi/3), & p_7 &= (\pi - \theta_0, \pi), \\
p_4 &= (\theta_0, 4\pi/3), & p_8 &= (\pi - \theta_0, 5\pi/3).
\end{aligned}$$

where  $\theta_0 = \arccos(-1/3)$ . Then a generator for  $T_{2u}$  is given by

$$v_i = \frac{\partial_\phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{1, 2, 3, 4\}, \quad v_i = -\frac{\partial_\phi}{\sin \theta} \Big|_{p_i} \text{ for } i \in \{5, 6, 7, 8\}.$$

Note that the vertices  $p_1$  and  $p_5$  (at the poles) are actually fixed under this generator. The Hessian becomes

$$\nabla_p^2 \log I^\pm(v, v) = -16\pi^2 \int_{\mathbb{S}^2} \partial_\phi G_A \partial_\phi G_B d\mu^\pm,$$

where  $A = \{2, 3, 4\}$  and  $B = \{6, 7, 8\}$ . One easily notices that the integrand  $\partial_\phi G_A \partial_\phi G_B \leq 0$  everywhere on the sphere. So the Hessian is positive definite on  $T_{2u}$ .

**3.4. The definiteness on  $T_{2g}$ .** For  $T_{2g}$  we orient the cube with all eight vertices at  $\frac{1}{\sqrt{3}}(\varepsilon_1, \varepsilon_2, \varepsilon_3)$  with  $\varepsilon_i \in \{\pm 1\}$ . The  $zx$ -component of  $T_{2g}$  corresponds to the partition

$$A = \left\{ \frac{1}{\sqrt{3}}(\varepsilon_1, \varepsilon_2, \varepsilon_3) : \varepsilon_1 \varepsilon_3 = +1 \right\} = \frac{1}{\sqrt{3}} \{ (1, 1, 1), (1, -1, 1), (-1, 1, -1), (-1, -1, -1) \},$$

$$B = \left\{ \frac{1}{\sqrt{3}}(\varepsilon_1, \varepsilon_2, \varepsilon_3) : \varepsilon_1 \varepsilon_3 = -1 \right\} = \frac{1}{\sqrt{3}} \{ (1, 1, -1), (1, -1, -1), (-1, 1, 1), (-1, -1, 1) \},$$

i.e., the four vertices with  $XZ > 0$  and those with  $XZ < 0$ . The corresponding generator assigns  $+\partial_\phi/\sin\theta$  to each vertex in  $A$  and  $-\partial_\phi/\sin\theta$  to each vertex in  $B$ , giving

$$\nabla_p^2 \log J_8(v, v) = -16\pi^2 \int_{\mathbb{S}^2} \partial_\phi G_A \partial_\phi G_B d(\mu^+ + \mu^-).$$

Because  $A$  is not invariant under the rotation generated by  $\partial_\phi$ , the integrand  $\partial_\phi G_A \cdot \partial_\phi G_B$  is *not* pointwise non-positive, and a direct sign argument fails. Instead we use group averaging.

Group averaging. The measure  $d(\mu^+ + \mu^-)$  is  $O_h$ -invariant, so the integral is unchanged by any  $g \in O_h$ . Consider the coordinate swap  $\sigma: (X, Y, Z) \mapsto (X, Z, Y)$ , which is a symmetry of the cube  $\{\pm 1, \pm 1, \pm 1\}/\sqrt{3}$  and hence lies in  $O_h$ . Under  $\sigma$ :

- the partition  $\{A, B\}$  maps to the partition  $\{A', B'\}$  where

$$A' = \frac{1}{\sqrt{3}} \{ (1, 1, 1), (1, 1, -1), (-1, -1, 1), (-1, -1, -1) \}, \quad B' = \text{complement of } A'$$

(the  $xy$ -component of  $T_{2g}$ , with  $\varepsilon_1 \varepsilon_2 = +1$  on  $A'$ );

- the Killing field  $\partial_\phi = X\partial_Y - Y\partial_X$  maps to  $\partial_\chi = X\partial_Z - Z\partial_X$  (rotation about the  $Y$ -axis).

Both  $v = (\text{sign}(\varepsilon_1 \varepsilon_3) \partial_\phi / \sin\theta)$  and  $v' = (\text{sign}(\varepsilon_1 \varepsilon_2) \partial_\phi / \sin\theta)$  are  $T_{2g}$ -generators of the same norm in the decomposition  $V = \cdots \oplus T_{2g} \oplus \cdots$  (since  $\varepsilon_1 \varepsilon_3$  and  $\varepsilon_1 \varepsilon_2$  are both  $T_{2g}$  basis functions, and both generators have  $\|v\|^2 = \|v'\|^2 = 8$ ). By Schur's lemma (see the remark at the beginning of this section), the Hessian takes the same value on every  $T_{2g}$ -generator of equal norm. Therefore

$$\int_{\mathbb{S}^2} \partial_\phi G_A \partial_\phi G_B d(\mu^+ + \mu^-) = \int_{\mathbb{S}^2} \partial_\phi G_{A'} \partial_\phi G_{B'} d(\mu^+ + \mu^-).$$

It therefore suffices to prove the integral on the right is negative, i.e., to work with the  $xy$ -partition  $(A', B')$ .

Formulas for  $G_{A'}$  and  $G_{B'}$ . The inner products  $p \cdot p_i$  for the four  $A'$ -vertices pair into two antipodal pairs giving inner products  $\pm(X+Y+Z)/\sqrt{3}$  and  $\pm(X+Y-Z)/\sqrt{3}$ . Hence

$$Q_{A'} := \prod_{i \in A'} (1 - p \cdot p_i) = \left(1 - \frac{(X+Y+Z)^2}{3}\right) \left(1 - \frac{(X+Y-Z)^2}{3}\right) =: Q_1 R_1,$$

so  $G_{A'} = -\frac{1}{4\pi} \log Q_{A'} + C_{A'}$ . Since  $B'$ -vertices are  $A'$ -vertices with  $Y \mapsto -Y$ , we get  $Q_{B'}(X, Y, Z) = Q_{A'}(X, -Y, Z) = Q_2 R_2$  where  $Q_2 = 1 - (X - Y + Z)^2/3$  and  $R_2 = 1 - (X - Y - Z)^2/3$ , and  $G_{B'} = -\frac{1}{4\pi} \log Q_{B'} + C_{B'}$ .

Derivation of  $\partial_\phi G_{A'}$ . Write  $Q_{A'} = Q_1 R_1$ . Since  $\partial_\phi(X + Y \pm Z) = X - Y$ , one computes

$$\partial_\phi \log Q_1 = -\frac{2(X+Y+Z)(X-Y)}{3Q_1}, \quad \partial_\phi \log R_1 = -\frac{2(X+Y-Z)(X-Y)}{3R_1}.$$

Therefore

$$\partial_\phi G_{A'} = \frac{X-Y}{6\pi} \left[ \frac{X+Y+Z}{Q_1} + \frac{X+Y-Z}{R_1} \right].$$

Combining the two fractions:  $(X+Y+Z)(1-(X+Y-Z)^2/3) + (X+Y-Z)(1-(X+Y+Z)^2/3) = 2(X+Y)(1-\frac{(X+Y)^2-Z^2}{3})$ . On  $\mathbb{S}^2$ ,  $1-\frac{(X+Y)^2-Z^2}{3} = \frac{2}{3}(2-X^2-XY-Y^2)$ , so

$$\partial_\phi G_{A'} = \frac{2(X^2-Y^2)(2-X^2-XY-Y^2)}{9\pi Q_1 R_1}.$$

The same computation with  $Y \rightarrow -Y$  yields

$$\partial_\phi G_{B'} = -\frac{2(X^2-Y^2)(2-X^2+XY-Y^2)}{9\pi Q_2 R_2}.$$

The integrand is non-positive. Multiplying,

$$\partial_\phi G_{A'} \cdot \partial_\phi G_{B'} = -\frac{4(X^2-Y^2)^2(2-X^2-XY-Y^2)(2-X^2+XY-Y^2)}{81\pi^2 Q_1 Q_2 R_1 R_2}.$$

Note  $(2-X^2-XY-Y^2)(2-X^2+XY-Y^2) = (2-X^2-Y^2)^2 - (XY)^2 = (1+Z^2)^2 - (XY)^2$ . On  $\mathbb{S}^2$ :  $1+Z^2 \geq 1$ , while  $|XY| \leq (X^2+Y^2)/2 = (1-Z^2)/2 \leq 1/2 < 1+Z^2$ , so  $(1+Z^2)^2 - (XY)^2 > 0$  everywhere. Likewise  $(X^2-Y^2)^2 \geq 0$ , and  $Q_1, Q_2, R_1, R_2 > 0$  away from the eight cube vertices. Therefore the integrand is non-positive, and equals zero only when  $X^2 = Y^2$ , a set of measure zero. Consequently,

$$\int_{\mathbb{S}^2} \partial_\phi G_{A'} \partial_\phi G_{B'} d(\mu^+ + \mu^-) < 0,$$

and by the group-averaging equality above,  $\int_{\mathbb{S}^2} \partial_\phi G_A \partial_\phi G_B d(\mu^+ + \mu^-) < 0$  as well. The Hessian is therefore positive definite on  $T_{2g}$ .

## REFERENCES

*Email address:* [chenhao5@shanghaitech.edu.cn](mailto:chenhao5@shanghaitech.edu.cn)

SHANGHAI TECH UNIVERSITY, SHANGHAI, CHINA