



Role of Clinical Pharmacists in Intensive Care Units

Pharm.D Vault Editorial Team | Educational Review Draft

Abstract. Critically ill patients commonly receive complex, high-risk medicine regimens while organ function, physiology, and clinical priorities change rapidly. Clinical pharmacists contribute to intensive care unit (ICU) teams through medication reconciliation, therapeutic optimization, dose adjustment, antimicrobial stewardship, adverse-event prevention, and transitions-of-care support. This educational review outlines key pharmacist activities, implementation considerations, and evidence-informed outcome measures. Pharmacist integration should be matched to local ICU needs and supported by collaborative practice, access to clinical data, and consistent documentation.

Keywords: critical care pharmacy; intensive care unit; medication safety; antimicrobial stewardship; multidisciplinary care

Introduction

The ICU is a high-risk medication environment. Patients may have altered absorption, distribution, metabolism, and elimination; receive vasoactive drugs, sedatives, antimicrobials, anticoagulants, and continuous infusions; and experience frequent transfers between care settings. These features create opportunities for medication errors and for treatment that no longer matches the patient's current physiology. A clinical pharmacist embedded in the multidisciplinary team can help identify and resolve medication-related problems in real time.

Core Patient Care Activities

Core activities include prospective medication review, renal and hepatic dose adjustment, therapeutic drug monitoring, anticoagulation assessment, antimicrobial optimization, evaluation of sedation and analgesia plans, prevention of drug interactions, and medication reconciliation. The relative priority of these activities should be determined by the ICU population and service model. For example, a unit with high antimicrobial use may prioritize stewardship and microbiology review, while a unit with frequent transfers may focus on admission and discharge reconciliation.

Antimicrobial Stewardship and Infection Management

Clinical pharmacists can support selection, dosing, infusion strategy, monitoring, and de-escalation of antimicrobials. In critically ill patients, changing renal function, extracorporeal therapies, fluid shifts, and altered tissue perfusion can complicate dose selection. Pharmacists should work with infectious diseases, microbiology, and ICU clinicians to ensure that antimicrobial decisions are individualized, documented, and reassessed as cultures and clinical response evolve.

Medication Safety and Transitions of Care

Medication safety work includes reviewing high-alert medicines, infusion concentrations, compatibility concerns, duplicate therapy, and monitoring requirements. At ICU admission and discharge, medication reconciliation helps reduce unintended discrepancies. A pharmacist can also prepare patient-specific discharge counselling and communicate medication changes to the receiving team. These activities are especially important when patients move between the ICU, ward, rehabilitation, and outpatient settings.

Measuring Value Without Oversimplifying It

Program evaluation should use measures that fit the service. Potential indicators include the number and type of pharmacist interventions, acceptance of recommendations, time to antimicrobial optimization, high-risk medication review completion, medication discrepancy rates, and adverse drug-event measures. Patient-centered outcomes such as length of stay, mortality, and ventilation duration are important but may be influenced by many factors. A balanced dashboard should therefore combine process measures, safety measures, and team-reported usefulness.

Implementation Considerations

Successful integration requires clarity about the pharmacist's role, consistent participation in rounds, access to laboratory and medication records, and agreed routes for urgent escalation. Services should define coverage hours, handover practices, documentation standards, and referral criteria. Education is also needed so that pharmacists, physicians, nurses, and trainees understand how recommendations will be discussed and implemented. Local quality-improvement cycles can identify which activities deliver the greatest clinical value.

Conclusion

Clinical pharmacists can make a meaningful contribution to ICU medication safety and therapeutic optimization when they are integrated into multidisciplinary care. The strongest model is not a list of isolated interventions; it is a reliable service that anticipates risk, communicates clearly, and follows patients across transitions of care.

References

1. Lee H, Ryu K, Sohn Y, et al. Impact on patient outcomes of pharmacist participation in multidisciplinary critical care teams: a systematic review and meta-analysis. *Critical Care Medicine*. 2019;47(9):1243-1250. doi:10.1097/CCM.0000000000003830.
2. Lat I, Paciullo C, Daley MJ, et al. Position paper on critical care pharmacy services: 2020 update. *Critical Care Medicine*. 2020;48(9):e813-e834. doi:10.1097/CCM.0000000000004437.
3. Arredondo E, Udeani G, Horseman M, et al. Role of clinical pharmacists in intensive care units. *Cureus*. 2021;13(9):e17929. doi:10.7759/cureus.17929.
4. Rice M, Lear A, Kane-Gill S, Seybert AL, Smithburger P. Pharmacy personnel involvement in transitions of care of intensive care unit patients: a systematic review. *Journal of Pharmacy Practice*. 2021;34(1):117-126. doi:10.1177/0897190020911524.

Editorial note: This educational review draft is not a substitute for patient-specific clinical advice. Verify all references, local policy statements, author details, and editorial requirements before publication.