



Are you up to date on the latest evidence for endoscopic weight-loss procedures?



CAUTION: The law restricts these devices to sale by or on the order of a physician. Indications, contraindications, warnings, and instructions for use can be found at www.IFU-BSCI.com. Products shown for INFORMATION purposes only and may not be approved or for sale in certain countries. This material is not intended for use in France.

Results from different clinical investigations are not directly comparable. Information provided for educational purposes only.

ENDO-2428703-AA © 2026 Boston Scientific Corporation or its affiliates. All rights reserved.

CE 2797

endura
Weight Loss Solutions

Endoscopic Sleeve Gastroplasty (ESG)



Efficacy

- In the multicentre randomised controlled **MERIT trial**, ESG achieved a **mean total body weight loss (TBWL) of 13.6% at 12 months**, significantly superior to lifestyle intervention alone.¹

Efficacy has been consistently confirmed in larger prospective cohorts. A prospective multicentre study reported a **mean TBWL of 15.5% at 12 months and 15.1% at 24 months**.²
- In a large prospective cohort, ESG achieved a **mean TBWL of 21.2% at 12 months**, with a high proportion of patients achieving clinically meaningful weight loss.³

Efficacy was further demonstrated in a longitudinal cohort, reporting a **mean TBWL of 15.6% at 12 months**.⁴
- IFSO Bariatric Endoscopy Committee meta-analysis** reported a **mean TBWL of 17.56% at 12 months**, confirming reproducible efficacy across heterogeneous clinical settings.⁵

1

Mean TBWL of
**13.6% at
12 months**
Randomised
Controlled Trial

2

Mean TBWL of
**21.2% at
12 months**
Large Prospective Cohort

3

Mean TBWL of
**17.56% at
12 months**
Meta-Analysis






Long-Term Results

A prospective long-term study reported a **mean TBWL of 18.19% at 4 years**, with the majority of patients maintaining clinically relevant weight loss.³


Five-year follow-up data confirmed a **mean TBWL of 15.9% at 5 years**, supporting long-term maintenance of efficacy.⁴



Safety

In the MERIT trial, the **serious adverse event (SAE) rate was 2%**, with no mortality and full resolution of all events.¹

In the IFSO meta-analysis, the **pooled SAE rate was 1.25%**, confirming a favorable and consistent safety profile across large populations.⁵





Transoral Outlet Reduction (TORe)

Efficacy

- 1 In a large retrospective cohort, TORe achieved a **mean total body weight loss (TBWL) of 8.5% at 12 months**, demonstrating clinically meaningful weight loss in patients with weight regain after Roux-en-Y gastric bypass.⁶
- 2 In patients treated for dumping syndrome and/or weight regain, TORe resulted in a **mean TBWL of 9.9% at 12 months**, confirming efficacy in both weight and symptom control.⁷ Mean **TBWL was 8.1% at 24 months**, with **61% of patients maintaining >5% TBWL**.⁷ Resolution of dumping syndrome was observed in **57.2% of patients at 24 months**.⁷
- 3 In a matched cohort comparison, TORe achieved comparable weight loss to surgical gastrojejunal revision; in the TORe cohort, **mean TBWL was 8.5% at 1 year, 6.9% at 3 years, and 8.8% at 5 years**, while maintaining a low severe adverse event rates.⁸

1

Mean TBWL of
8.5% at
12 months

2

Mean TBWL of
8.1% at
24 months

3


Mean TBWL of
6.9% at
3 years
8.8% at
5 years





Long-Term Results

In a large retrospective study, TORe resulted in a **mean TBWL of 8.8% at 5 years**, with **77% of patients stopping weight regain**.⁶ Mean absolute weight loss at 5 years was **10.3 kg**.⁶



Safety

In the 5-year follow-up study, **no severe adverse events were reported**, with an overall **SAE rate of 3.9%** and no mortality.⁶

In a matched cohort comparison, TORe was associated with **0% severe adverse events**, compared with **19.4% in the surgical revision group**, while achieving comparable long-term weight loss.⁸





REFERENCES

ESG

1. Abu Dayyeh BK. Endoscopic sleeve gastropasty for treatment of class I and II obesity (MERIT). Lancet. 2022. N=85 ESG; 150 exposed to ESG.
2. Alqahtani A. Prospective multicentre study of endoscopic sleeve gastropasty. Gastrointest Endosc. 2023. N=656.
3. Bhandari M. Four-year outcomes of endoscopic sleeve gastropasty. J Minim Access Surg. 2023. N=612.
4. Sharaiha RZ. Five-year outcomes of endoscopic sleeve gastropasty. Clin Gastroenterol Hepatol. 2021. N=216 (5-year follow-up N=56).
5. Abu Dayyeh BK. IFSO evidence-based review and position statement on ESG. Obes Surg. 2024. N=15,714.

TORe

6. Jirapinyo P. Five-year outcomes of transoral outlet reduction after Roux-en-Y gastric bypass. Gastrointest Endosc. 2020. N=331.
7. Pontecorvi V. Long-term outcomes of transoral outlet reduction for dumping syndrome and weight regain. Obes Surg. 2023. N=87.
8. Dolan RD. Endoscopic versus surgical gastrojejunal revision: 5-year safety and efficacy comparison. Gastrointest Endosc. 2021. N=31 TORe vs 31 surgical.

CAUTION: The law restricts these devices to sale by or on the order of a physician. Indications, contraindications, warnings, and instructions for use can be found at www.IFU-BSCI.com. Products shown for INFORMATION purposes only and may not be approved or for sale in certain countries. This material is not intended for use in France.

Results from different clinical investigations are not directly comparable.
Information provided for educational purposes only.

ENDO-2428703-AA © 2026 Boston Scientific Corporation or its affiliates.
All rights reserved.

CE 2797

endura
Weight Loss Solutions