

DEPARTMENT OF RADIO-DIAGNOSIS & IMAGING

Report : MRI

Report Date:

Patient Name :	Amit Kumar Pal		
Age:	28	UHID :	UHID-00328781
Sex:	Male	Bill No :	OPD17-0031387
Ref. Doctor :	Dr.Ashutosh Shrivastava	Bill Date:	02/05/17 11:51
Ward:		Room No:	

MR Imaging of left knee joint was performed on a 1.5 T scanner using SENSE FLEX-M COIL. T1W, T2W, PDW Fat saturated and gradient images were obtained in the coronal, sagittal and axial planes using SE, TSE, STIR, FFE and Grase sequences.

Follow up case of GCT lower end of femur, status- post curettage with bone grafting.

Mildly lobulated intramedullary altered signal intensity with internal fluid signal as well as osseous signal involving epimetaphysis of lower end of left femur with relative sparing of lateral femoral condyle consistent with post operative fluid with bone chips, lesion measures 77 x 53 x 36mm. Lesion is sharply demarcated without obvious surrounding STIR hyperintensity suggestive of no perilesional marrow edema(favours benign pathology). Mildly thickened medial cortex of lower end of femur seen due to periosteal reaction however small focal breach of cortex is also seen on medial aspect with associated mild edema in adjacent soft tissue. No obvious intra articular extension of lesion is seen however marked cortical thinning of intercondylar roof near the attachment site of PCL is seen.

Medial patellofemoral ligament is not well seen with marked surrounding STIR hyperintensity ? trauma of variable degree/ preoperative trauma.

Mild marginal osteophytes seen to involving medial/ lateral tibiofemoral and patellofemoral joints with associated patchy articular cartilage eburnation of both tibiofemoral and medial patellar facet cartilages- s/o degenerative disease.

Rest of visualised sections of femur, tibia, fibula and patella are normal in signal intensity pattern.

The tibio-femoral, patello-femoral and proximal tibio-fibular joints are normal in alignment, joint space and signal intensity. (Please correlate with Plain X-rays/CT Scan to assess cortical bony lesions/injuries.)

No significant joint effusion seen.

Hoffas fat pad appears normal.

The anterior and posterior cruciate ligament appear normal.

Medial and lateral menisci are normal in morphology and signal intensity.

The collateral ligaments are normal in integrity and signal intensity.

Ligamentum patellae appears normal.

Please correlate clinically and with other investigations.

.....End of Report.....

DR.SANDEEP MAKEN,MD

DR. RAVI SHANKER, MD

DR. KRISHNA BHARDWAJ, MD, DMRD

ASSO. DIRECTOR IMAGING

ASSO. CONSULTANT RADIOLOGY

ASSO. CONSULTANT RADIOLOGIST

Note: The information in this report is based on interpretation of images. This report is not the diagnosis and should be correlated with clinical details and other investigations.