

NAME: AMIT KUMAR PAL - 524279**AGE: 27 YRS / M ABH****REF BY: AIMS****DATE: 17/06/2016** DATE NO.MIS-2016-0023**MRI OF LEFT KNEE WITH SCREENING LEG**

MRI of left knee was done on HDXT MR scanner & following sequences were obtained.

Axial: T2W & T2W FS

Coronal: T1W & PD FS

Sagittal: T2, GRE & PD FS

Clinical profile: Patient is a known case of GCT with post operative bone grafting status.

FINDINGS:

Epimetaphyseal region of distal femur reveals lytic expansile intraosseous lesion, measuring ~ 3.5 (AP) x 5.7 (TR) x 7.5 (CC) cm in size with altered signal intensities within (T1W hypointense and STIR mixed signal intensities) and associated cortical breach in the antero-medial aspect with adjoining soft tissue thickening. Minimal free fluid and altered signal intensity is noted in the prepatellar and suprapatellar region without any obvious involvement of the patella.

Few well defined T2W / STIR hyperintense and T1W isointense to mildly hyperintense areas are also noted within the above mention intraosseous lesion. One of the area showing fluid fluid level (likely with hemorrhagic component).

The superior margin is ~9.5 cm proximal to medial femoral condyle and lower margins are ~ 1.4 cm proximal to medial femoral condyle.

No intraarticular extension is noted.

Medial retinaculum is not visualized.

Mild hyperintense signals are noted in the anterior cruciate ligament.

The medial and lateral menisci are normal with no evidence of any tear or any area of signal alteration. The articular cartilage too is normal.

The posterior cruciate as well as the lateral and medial collateral ligaments are unremarkable with no evidence of a tear.