

- Beta diversity: try separating into treated vs. untreated
- For indicator species, show how many samples are 0
 - Maybe do pseudo log
 - Half of the smallest non-zero sample
 - `geom_jitter()`
 - Write in legend how many are 0
 - Compare analysis: otu vs. AVS?, filtered out low abundance first
 - `Scale_y_log10` to make the y-axis lines even intervals
 - Turn bottom graph into boxplot (supp vs. no supp) and add both
- Differential abundance: see which taxa correlated with vitamin D
 - Merge first differential abundance with indicator species
- picrust2
 - cutoff at 0.005
 - Can calculate by hand to check if maaslin expected
 - Are log 2 fold changes too low or p values too low
- Correlation matrix
 - Run at genus level
 - Add *Frisingicoccus* to correlation matrix
 - Show numbers on correlation matrix
- Don't need to filter for treatment vs. no treatment (since all analysis have been run without)
- Figure 1: Alpha, beta diversity
- Figure 2: Differential abundance, indicator value (stat numbers) + boxplots [decide based on which analysis we chose]
- Figure 3: picrust (individual functions), pathway flowchart (drawn on powerpoint), {add functions with all significance into supplementary figures}
- Figure 4: [change to genus level] genus + function + edss {, can include top 10 abundance in supplemental figures}
- (maybe) Figure 5: random forest (maybe)