

Stability Of Paths

In what conditions is a path through a dynamic system
stable?

Mathematics Extended Essay

Colonel By Secondary School

Word Count: 3666

Jeffrey Li

January 24, 2021

Abstract

In this paper, we first reestablish the foundations of classical Lyapunov point stability. We then extend Lyapunov's arguments and establish a rigorous definition of Path Stability, in line with Lyapunov's original reasoning. Next, we note some important properties derived from the definition of Path Stability which allow the emergent behavior of the path to be decomposed and analyzed individually on each point. Finally, by linearizing the differential equation describing change of state, we arrive at insights linking the stability of the path to the Quadratic Form of the Jacobian of the system, another problem that has extensive literature in the field of linear algebra, demonstrating the concept of Path Stability has deep conceptual roots reaching into the heart of mathematics.

Contents

1 Research Question	4
2 Introduction	4
3 Motivation	5
4 Dynamical Systems	5
4.1 Definition	5
4.2 Path Notation	6
4.3 Evolution Function	6
5 Lypaunov Point Stability	7
6 Path Stability	7
7 General Conditions For Path Stability	9
7.1 Motivation	9
7.2 Time Invariance	10
7.3 Local Attractiveness	11
8 Conditions For Local Attractiveness	13
8.1 Local Distance Derivative	13
8.2 Local Dot Product	14
9 Linearization	16
10 Conclusion	18
References	19

1 Research Question

Given a particle travelling under a dynamic system, under what conditions will an arbitrarily small disturbance to the initial position always result in an arbitrarily small displacement to the new path, regardless of the time?

2 Introduction

In life, we can often describe the phenomenon we observe as a product of two factors: ever-present change and an element of randomness.

Firstly, the way in which things change often can be described as a dynamic system, which are systems whose development in time is dependent upon their current state. Dynamic systems are integral to many fields of study; through Hamiltonian mechanics, we can understand motion as a dynamic system whose state is defined as the combination of position and velocity. In ecological models, populations will grow if resources are abundant, and shrink if resources are scarce. Hence, the change from one year to the next is largely dependent upon the population of a species from the year before.

Secondly, no idealized model is ever perfectly accurate to its real world counterpart because of the prevalence of random disturbances of various magnitudes throughout every aspect of life. The work of the Russian mathematician Aleksandr Lyapunov at the turn of the 20th century has given mathematicians frameworks and methods for understanding the stability of a system; in other words, to what degree the system is resilient to small initial disturbances. In the following century, this topic has been extensively studied with respect to stable *points*, but in comparison, I found far fewer works that deal with the stability of entire *paths*: studying how the state of a system develops over an arbitrarily large duration of time.

3 Motivation

For me, the consideration of the problem of stability arose from contemplating a ball in a rounded crater. The ball may start at any point along within the semi-circular cavity, but would always roll down and come to rest at the center. A ball at the top of the hill, however would roll in very different directions when given two slightly different initial positions. Similar concepts of stability can be found in prices, which are sometimes subject to extreme fluctuation and other times are extremely inert, or celestial motion, which was the origin of Chaos Theory through the three-body problem, while at the same time illustrating unexpected stability in the form of Lagrange Points within an orbit. All of these led me to wonder if Stability could somehow be quantified, and be described as a numerical property of a field, much like potential energy for example. The ball in the valley will be used as the main reference point because of its simplicity, but I constantly referred back to the other cases given as a source of intuition, to examine properties of Stability that transcended any one example.

4 Dynamical Systems

4.1 Definition

To begin, I believe it is important to clearly define what is meant by a system. In the example of the ball in a valley, the system can be said to be defined by the set of points bounded within the valley in combination with the laws of motion that dictate the ball's path. Thus in this paper, a unique dynamic system of dimension n , denoted as \mathcal{S} , will refer to a unique domain $\mathcal{D} \in \mathbb{R}^n$ in conjunction with an associated Evolution Function $f_s(\mathbf{x}) : \mathcal{D} \rightarrow \mathbb{R}^n$ that gives the 'trajectory' of a particle when given its current 'position'. Inside of the domain \mathcal{D} , a point can be described using the vector $\mathbf{x} \in \mathcal{D}$. Again referring to

the initial example, the area encompassing the valley can be thought of as the system's domain \mathcal{D} , while the Evolution Function can be thought of as the laws of motion. A more formal description will be given in the following paragraph.

4.2 Path Notation

After considering the initial premise, I realized that these conditions meant that, knowing the initial position \mathbf{x}_0 of a particle at time 0, then using the aforementioned Evolution Function, we are able to derive the position of the particle at all future times. Therefore, I created a notation of Paths to describe this idea. For the particle beginning at the point \mathbf{x}_0 at time 0, the corresponding Path Function $P_{\mathbf{x}_0}(t)$ is a function from $[0, \infty) \rightarrow \mathcal{D}$ such that for any time $t_0 \in [0, \infty)$, the position of the particle at that instant is given by $P_{\mathbf{x}_0}(t_0)$. For example, by definition, $P_{\mathbf{x}_0}(0)$ will always give the particle's initial position - or in other words, $P_{\mathbf{x}_0}(0) = \mathbf{x}_0$.

4.3 Evolution Function

The time derivative of the Path Function can be used to formally define a system's Evolution Function. By convention, the time derivative $\frac{d}{dt}(P_{\mathbf{x}_0}(t))$ will be notated as $\dot{P}_{\mathbf{x}_0}(t)$, with the single dot above the function indicating a first derivative. Earlier, it was said that the Evolution Function will return a particle's trajectory in time given its current position. Therefore, the Evolution Function $f_s(\mathbf{x})$ of the system \mathcal{S} is defined as satisfying

$$f_s(P_{\mathbf{x}_0}(t)) = \dot{P}_{\mathbf{x}_0}(t) \tag{1}$$

meaning that the change with respect to time of a particle's position is given by the Evolution Function evaluated at the particle's current location.

5 Lyapunov Point Stability

In his original article from 1892, Lyapunov (1992) layed out a definition for stability in a Dynamical System which is mathematically equivalent to the following:

Definition 5.1 (Lyapunov Stability). Let there exist a system \mathcal{S} as previously defined. Within this system, suppose we claim there is an equilibrium point $\mathbf{x}_e \in \mathcal{D}$. Then the point \mathbf{x}_e is stable iff, for every positive real number $\epsilon > 0$, we can find $\delta > 0$ such that, for any point \mathbf{x}_0 where $\|\mathbf{x}_e - \mathbf{x}_0\| < \delta$, the path $P_{\mathbf{x}_0}(t)$ satisfies the inequality $\|\mathbf{x}_e - P_{\mathbf{x}_0}(t)\| < \epsilon$ for all $t \in [0, \infty)$.

In essence, what this definition proposes is that a point is stable if a particle that starts ‘close’ to it will always remain ‘close’ to the initial point, even as time goes to infinity. I found this formalization very compelling for two main reasons. First of all, it captures the core of what the intuitive understanding of stability. Secondly, Lyapunov’s definition allows for stability to be treated in a much more rigorous and analytical way, as the conditions it sets out can definitively proven to be true or false using strictly mathematical tools. Not only that, but because of its usage of the epsilon-delta formalization at the core of calculus, the further developments that I made followed very naturally, and I was able to use familiar methods of attack despite the novelty of the problem.

6 Path Stability

Despite the ingenuity of Lyapunov’s definition, there was still one major difference between my conception of stability and Lyapunov’s. While Lyapunov treated stability in terms of the behavior of local points, I wanted to describe stability in terms of local *paths*. Because I really liked methods Lyapunov used in his approach, the first thing I did was to take Lyapunov’s original conclusions

and rephrase the formalization to relate to the behavior of paths. That is how I came to my definition of Path Stability, as follows:

Definition 6.1 (Path Stability). Within the system \mathcal{S} , suppose there exists the path P_{x_0} which starts at \mathbf{x}_e . This path is Unbounded Stable iff, for every positive real number $\epsilon > 0$, we can find $\delta > 0$ such that, for any point \mathbf{x}_0 where $\|\mathbf{x}_e - \mathbf{x}_0\| < \delta$, the paths P_{x_e} and P_{x_0} satisfy the inequality $\|P_{x_e}(t) - P_{x_0}(t)\| < \epsilon$ for all $t \in [0, \infty)$.

The first thing to remark from definition 6.1 is the similarities it shares with the original definition 5.1. In fact, the two definitions are mathematically identical except for, where the stable **point** \mathbf{x}_e is used by Lyapunov, my definition substitutes the stable **path** P_{x_e} . This subtle yet powerful difference will be discussed in the following sections.

However, to conclude the current section, I will draw attention to the obvious corollary resulting from the parallel definitions - that any point which satisfies the conditions of Lyapunov Stability must also be the origin of a corresponding path that satisfies Path Stability. To illustrate this, note that in order for a point to be Lyapunov Stable, it must be a point of equilibrium - in other words, its path never moves and is defined as the constant function $P_{x_e}(t) = \mathbf{x}_e$ for every time t . Therefore, any other nearby path that permanently stays ‘close’ to the *point* \mathbf{x}_e will consequently also stay ‘close’ to the *path* $P_{x_e}(t)$. This corollary serves to clearly show that Path Stability is not only fully compatible with Lyapunov’s definition, but in fact generalizes the scope of the original result.

7 General Conditions For Path Stability

7.1 Motivation

After laying out a rigorous definition for what I had originally conceptualized in Path Stability, my next goal, and the focus of the rest of this paper, was to derive the general conditions that were needed to satisfy definition 6.1. As has been shown, Lyapunov Stable equilibrium points are always Path Stable, but far more interesting to me was studying Path Stability for non-equilibrium points of origin.

In particular, I noted that the path a particle travels can be defined through the set of all points that the path passes through. For a given path, any point that lies in the system can be definitively shown to either lie on the path or not lie on the path. The path itself is a real-valued continuous function from \mathcal{R}^+ to \mathcal{D} , which is both not finitely bounded and can be arbitrarily complex, making general considerations on path functions not limited to special cases exceedingly difficult. Meanwhile, the constituent *points* that comprise the path are well defined vector elements, which are much simpler to study. In addition, while the set of points that lie along a given path may be uncountably infinite, because of the conditions on the system of continuity and differentiability, properties of points along a path will also in general be continuous and differentiable, meaning that generalized and meaningful results can still be drawn about the entire set. For this reason, I directed my focus to finding what *local* properties on the individual points along a path must be satisfied in order for the path to be stable.

7.2 Time Invariance

Firstly, I wanted to note one particular corollary derived from the fact that paths were composed of a set of points: each point is also the origin of its own corresponding path. If the path originating at point \mathbf{x}_0 arrives at point \mathbf{x}_1 at the time t_0 , then the two will trace identical trajectories from that point onward; in other words,

$$P_{\mathbf{x}_0}(t_0 + s) = P_{\mathbf{x}_1}(t_1 + s)$$

The following is the rigorous proof of this idea, which I refer to as Time Invariance.

Theorem 7.1 (Time Invariance). *For any two paths $P_{\mathbf{x}_0}$ and $P_{\mathbf{x}_1}$, if there exists $t_0, t_1 \geq 0$ such that $P_{\mathbf{x}_0}(t_0) = P_{\mathbf{x}_1}(t_1)$, then $P_{\mathbf{x}_0}(t_0 + s) = P_{\mathbf{x}_1}(t_1 + s)$ for all $s \geq 0$.*

Proof. From the Taylor series expansion of a function $f(t)$ as found in Taylor and Mann (1983),

$$\begin{aligned} P_{\mathbf{x}_0}(t_0 + s) &= P_{\mathbf{x}_0}(t_0) + \sum_{n=1}^{\infty} \frac{s^n}{n!} P_{\mathbf{x}_0}^{(n)}(t_0) \\ &= P_{\mathbf{x}_0}(t_0) + \sum_{n=1}^{\infty} \frac{s^n}{n!} \left(\frac{d}{dt} P_{\mathbf{x}_0}^{(n-1)}(t) \Big|_{t=t_0} \right) \\ &= P_{\mathbf{x}_0}(t_0) + \sum_{n=1}^{\infty} \frac{s^n}{n!} \left(f_s^{(n-1)}(P_{\mathbf{x}_0}(t_0)) \right) \\ &= P_{\mathbf{x}_1}(t_1) + \sum_{n=1}^{\infty} \frac{s^n}{n!} \left(f_s^{(n-1)}(P_{\mathbf{x}_1}(t_1)) \right) \\ &= P_{\mathbf{x}_1}(t_1 + s) \end{aligned}$$

□

7.3 Local Attractiveness

The criteria I layed out for Path Stability in definition 6.1 can be stated as: if a given path P_{x_0} is stable, then any other path that starts ‘nearby’ P_{x_0} must remain ‘nearby’ for time $t \rightarrow \infty$.

I considered what this meant for each individual point. If, at each point along the path, all ‘nearby’ paths would be guaranteed to remain ‘nearby’ for an arbitrarily small period of time, then at no point along the entire path could a ‘nearby’ path diverge. Intuitively, this might be obvious, but to specify precisely what is meant by ‘nearby’ and ‘diverge’, I developed the concept of Local Attractiveness.

Definition 7.1 (Local Attractiveness). A point $x_0 \in \mathcal{D}$ is said to be locally attractive if, for every $\epsilon > 0$, there exists $\delta \in (0, \epsilon]$ such that for every x_δ where $\|x_\delta\| < \delta$, there exists $\sigma > 0$ for which the following is true for all $\Delta t \in [0, \sigma)$.

$$\|P_{x_0}(0) - P_{(x_0+x_\delta)}(0)\| \geq \|P_{x_0}(\Delta t) - P_{(x_0+x_\delta)}(\Delta t)\|. \quad (2)$$

Note the similarities between this definition and the epsilon-delta definition of the limit, which arose because I tried to use the concept of the limit to ‘squeeze’ all ‘nearby’ points to converge to the stable path.

Theorem 7.2. *For some path P_{x_0} , if every point on the path denoted by $x_t = P_{x_0}(t)$, where $t \in [0, \infty)$, is locally attractive, then the path P_{x_0} is Path Stable as defined in 6.1.*

Proof. Let $\tau > 0$ be an arbitrarily large length of time. For a given $\epsilon > 0$ and some $t \in [0, \tau]$, let $\delta_t \in (0, \epsilon]$ be equal to the corresponding δ value given in definition 7.1. We define $\delta_{min} = \min\{\delta_t : 0 \leq t \leq \tau\}$, and let $P_{x_\phi}(t)$ be any path such that $\|P_{x_\phi}(0) - P_{x_0}(0)\| < \delta_{min}$.

In the following, we use proof by contradiction to prove that, for all \mathbf{x}_ϕ where $\|\mathbf{x}_\phi - \mathbf{x}_0\| < \delta_{min}$ and $t \in [0, \tau]$, $\|P_{\mathbf{x}_\phi}(t) - P_{\mathbf{x}_0}(t)\| \leq \|\mathbf{x}_\phi - \mathbf{x}_0\|$.

We let $D(t) = \|P_{\mathbf{x}_\phi}(t) - P_{\mathbf{x}_0}(t)\|$, and assume that there exists some $t_0 \in [0, \tau]$ such that $D(t_0) > D(0) = \|\mathbf{x}_\phi - \mathbf{x}_0\|$. By definition of a path function, $P_{\mathbf{x}_\phi}(t)$ and $P_{\mathbf{x}_0}(t)$ are continuous on domain $[0, \infty)$ and the norm $\|\mathbf{x}\|$ is continuous on domain \mathcal{D} . Therefore, $D(t)$ is also continuous on $t \in [0, \infty)$, and there must exist some $t_1 \in [0, t_0)$ such that $D(t_1) = D(0)$ and,

$$D(t_k) > D(t_1) = D(0) = \|\mathbf{x}_\phi - \mathbf{x}_0\|, \quad \forall t_k \in (t_1, t_0). \quad (3)$$

We let $\mathbf{x}_\alpha = P_{\mathbf{x}_0}(t_1)$ and $\mathbf{x}_\beta = P_{\mathbf{x}_\phi}(t_1)$. Then we have,

$$\begin{aligned} D(t_1) &= \|P_{\mathbf{x}_\phi}(t_1) - P_{\mathbf{x}_0}(t_1)\| \\ &= \|\mathbf{x}_\beta - \mathbf{x}_\alpha\| \\ &= D(0) \\ &= \|P_{\mathbf{x}_\phi}(0) - P_{\mathbf{x}_0}(0)\| \\ &= \|\mathbf{x}_\phi - \mathbf{x}_0\| \\ &< \delta_{min} \leq \delta_{t_1}, \end{aligned} \quad (4)$$

from which there exists some σ such that for all $\Delta t \in [0, \sigma)$, we have,

$$\|P_{\mathbf{x}_\alpha}(0) - P_{\mathbf{x}_\beta}(0)\| \geq \|P_{\mathbf{x}_\alpha}(\Delta t) - P_{\mathbf{x}_\beta}(\Delta t)\| \quad (5)$$

by local attractiveness.

Since $P_{\mathbf{x}_\alpha}(0) = \mathbf{x}_\alpha$, we have $P_{\mathbf{x}_0}(t_1 + s) = P_{\mathbf{x}_\alpha}(s)$ for all $s \geq 0$ by theorem 7.1. Similarly, we have $P_{\mathbf{x}_\phi}(t_1 + s) = P_{\mathbf{x}_\beta}(s)$ for all $s \geq 0$. By setting $s = 0$ on

the left side of equation 5 and $s = \Delta t$ on the right side of equation 5, we obtain,

$$D(t_1) = \|P_{\mathbf{x}_0}(t_1) - P_{\mathbf{x}_\phi}(t_1)\| \geq \|P_{\mathbf{x}_0}(t_1 + \Delta t) - P_{\mathbf{x}_\phi}(t_1 + \Delta t)\|,$$

for some $\Delta t \in (0, t_0 - t_1)$. This is contradictory with equation 3. Therefore, $\|P_{\mathbf{x}_\phi}(t) - P_{\mathbf{x}_0}(t)\| \leq \|\mathbf{x}_\phi - \mathbf{x}_0\| < \delta_{min}$ for all $t \in [0, \tau]$. Since τ can be an arbitrarily large length of time, we can conclude that $P_{\mathbf{x}_0}(t)$ is bounded stable by denoting $\delta = \delta_{min}$ and $\mathbf{x}_\delta = \mathbf{x}_\phi - \mathbf{x}_0$. \square

8 Conditions For Local Attractiveness

8.1 Local Distance Derivative

Having shown that Local Attractiveness directly results in Path Stability, the next question to answer is what are the general conditions for a point to be Locally Attractive? From the definition, it is clear that if all ‘nearby’ paths around a point are getting ‘closer’ for some duration of time, then that point is locally attractive. This can be denoted analytically as follows.

Theorem 8.1. *For some point \mathbf{x}_0 , if there exists $\delta > 0$ such that every \mathbf{x}_α where $\|\mathbf{x}_0 - \mathbf{x}_\alpha\| < \delta$ satisfies the inequality*

$$\frac{d}{dt} \|P_{\mathbf{x}_0}(t) - P_{\mathbf{x}_\alpha}(t)\| < 0 \tag{6}$$

Then \mathbf{x}_0 is locally attractive.

Proof. From equation 6, the derivative of the distance between the two paths is less than 0. Therefore, by the Mean Value Theorem from Taylor and Mann (1983) and the definition of the derivative, there exists $t_1 > 0$ such that for the

following is true for all $\Delta t \in (0, t_1)$.

$$\|P_{\mathbf{x}_\alpha}(\Delta t) - P_{\mathbf{x}_0}(\Delta t)\| < \|P_{\mathbf{x}_\alpha}(0) - P_{\mathbf{x}_0}(0)\|. \quad (7)$$

By definition 7.1, the point \mathbf{x}_0 is locally attractive. \square

8.2 Local Dot Product

Next, what conditions must a point meet such that the derivative of the distance between it and any other local path is less than 0? To answer this, I considered that the displacement between any two paths $\Lambda(t) = (P_{\mathbf{x}_\alpha}(\Delta t) - P_{\mathbf{x}_0}(\Delta t))$ is itself a continuously differentiable vector valued function $\Lambda(t) : \mathbb{R} \rightarrow \mathbb{R}^n$. For a general vector, its magnitude will decrease if another small vector that points in the ‘opposite’ direction is added to it. Mathematically, the ‘opposite’ direction can be denoted by the inner product $\hat{v} \cdot \hat{w} < 0$. Therefore:

Theorem 8.2. *Given the continuous and differentiable vector valued function $\Lambda(t) : \mathbb{R} \rightarrow \mathbb{R}^n$,*

$$\Lambda(t) \cdot \dot{\Lambda}(t) < 0 \implies \frac{d}{dt} \|\Lambda(t)\| < 0$$

Proof. Let $\Lambda(t) : \mathbb{R} \rightarrow \mathbb{R}^n$ be a continuous and differentiable vector valued function such that

$$\Lambda(t) \cdot \dot{\Lambda}(t) < 0 \quad (8)$$

Let the operator $\|\Lambda(t)\|$ be the 2-norm of $\Lambda(t)$.

For any time t , we re-write $\Lambda(t)$ as $(\Lambda_1(t), \Lambda_2(t), \dots, \Lambda_n(t))$ where $\Lambda_i(t) : \mathbb{R} \rightarrow$

\mathbb{R} for $i = 1, 2, \dots, n$. By definition of the norm in \mathbb{R}^n ,

$$\begin{aligned}
 \|\Lambda\|^2 &= \sum_{i=1}^n \Lambda_i(t)^2 && \implies \\
 \frac{d}{dt} \|\Lambda\|^2 &= \frac{d}{dt} \left(\sum_{i=1}^n \Lambda_i(t)^2 \right) && \implies \\
 2\|\Lambda\| \|\Lambda\|' &= 2 \sum_{i=1}^n \Lambda_i(t) \dot{\Lambda}_i(t) && \implies \\
 \|\Lambda\|' &= \frac{\Lambda(t) \cdot \dot{\Lambda}(t)}{\|\Lambda\|} && (9)
 \end{aligned}$$

From equation 8, $\Lambda(t) \cdot \dot{\Lambda}(t) < 0$. From the definition of the norm, $\|\Lambda\| > 0$. Therefore,

$$\frac{d}{dt} \|\Lambda\| = \|\Lambda\|' = \frac{\Lambda(t) \cdot \dot{\Lambda}(t)}{\|\Lambda\|} < 0 \quad (10)$$

□

From theorem 8.1, the path functions $P_{\mathbf{x}_\alpha}(t)$ and $P_{\mathbf{x}_0}(t)$ are both continuously differentiable by definition. Therefore, $\Lambda(t) = (P_{\mathbf{x}_\alpha}(\Delta t) - P_{\mathbf{x}_0}(\Delta t))$ is also continuously differentiable. Therefore, by theorems 8.1 and 8.2, we have:

Theorem 8.3. *For some point \mathbf{x}_0 , if there exists $\delta > 0$ such that every \mathbf{x}_α where $\|\mathbf{x}_0 - \mathbf{x}_\alpha\| < \delta$ satisfies the inequality*

$$(P_{\mathbf{x}_0}(0) - P_{\mathbf{x}_\alpha}(0)) \cdot (\dot{P}_{\mathbf{x}_0}(0) - \dot{P}_{\mathbf{x}_\alpha}(0)) < 0 \quad (11)$$

Then \mathbf{x}_0 is locally attractive.

9 Linearization

Finally, we will evaluate what conditions are sufficient to that the aforementioned local dot product is always negative. When dealing with differentials under an arbitrarily small limit, as is the current case, it is often useful to linearize the function's derivative, which is analogous to taking the slope of a curve. To linearize a vector function, we use the Jacobian operator, as defined in Taylor and Mann (1983).

Definition 9.1 (Jacobian). Given \mathbf{x}_0 and some unit vector \mathbf{x}_δ for which $\|\mathbf{x}_\delta\| = 1$, the Jacobian matrix of the vector differentiable vector-valued function $\mathbf{f}_s : \mathbb{R} \rightarrow \mathbb{R}^n$ evaluated at point \mathbf{x}_0 is the linear transformation from $\mathbb{R}^n \rightarrow \mathbb{R}^n$ denoted by $\mathbf{J}_{\mathbf{f}_s}[\mathbf{x}_0]$ and given in the following equation.

$$\lim_{c \rightarrow 0} \frac{f_s(\mathbf{x}_0 + c\mathbf{x}_\delta) - f_s(\mathbf{x}_0)}{c} = \mathbf{J}_{\mathbf{f}_s}[\mathbf{x}_0](\mathbf{x}_\delta). \quad (12)$$

Theorem 9.1. For some point \mathbf{x}_0 , let $\mathbf{J}_{\mathbf{f}_s}[\mathbf{x}_0]$ be the Jacobian transformation at that point. If, all unit vectors $\{\mathbf{x}_\delta \in \mathbb{R}^n : \|\mathbf{x}_\delta\| = 1\}$ satisfy the inequality

$$\mathbf{x}_\delta \cdot \mathbf{J}_{\mathbf{f}_s}[\mathbf{x}_0]\mathbf{x}_\delta < 0 \quad (13)$$

Then the point \mathbf{x}_0 is Locally Attractive.

Proof. From the definition of the dot product, it is known to be continuous on \mathbb{R}^n . Therefore, for any two vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ where $\mathbf{x} \cdot \mathbf{y} < 0$, by definition of continuity, there exists some $\epsilon_1 > 0$ such that all $\delta\mathbf{y} \in \mathbb{R}^n$ where $\|\delta\mathbf{y}\| < \epsilon_1$ also satisfy the inequality

$$\mathbf{x} \cdot (\mathbf{y} + \delta\mathbf{y}) < 0. \quad (14)$$

From equation 12, we know that for any $\epsilon_1 > 0$, we can find $\delta_1 > 0$ such

that for scalar c ,

$$|c| \in (0, \delta_1) \implies \left\| \frac{f_s(\mathbf{x}_0 + c\mathbf{x}_\delta) - f_s(\mathbf{x}_0)}{c} - \mathbf{J}_{f_s}[\mathbf{x}_0]\mathbf{x}_\delta \right\| < \epsilon_1 \quad (15)$$

Therefore, knowing $\mathbf{x}_\delta \cdot \mathbf{J}_{f_s}[\mathbf{x}_0]\mathbf{x}_\delta < 0$, we can substitute in 14 \mathbf{x} for \mathbf{x}_δ , \mathbf{y} for $\mathbf{J}_{f_s}[\mathbf{x}_0]\mathbf{x}_\delta$ and $\delta\mathbf{y}$ for $(\frac{f_s(\mathbf{x}_0 + c\mathbf{x}_\delta) - f_s(\mathbf{x}_0)}{c} - \mathbf{J}_{f_s}[\mathbf{x}_0]\mathbf{x}_\delta)$, giving

$$\begin{aligned} \mathbf{x}_\delta \cdot \left(\mathbf{J}_{f_s}[\mathbf{x}_0]\mathbf{x}_\delta + \frac{f_s(\mathbf{x}_0 + c\mathbf{x}_\delta) - f_s(\mathbf{x}_0)}{c} - \mathbf{J}_{f_s}[\mathbf{x}_0]\mathbf{x}_\delta \right) &< 0 \implies \\ \mathbf{x}_\delta \cdot \left(\frac{f_s(\mathbf{x}_0 + c\mathbf{x}_\delta) - f_s(\mathbf{x}_0)}{c} \right) &< 0 \implies \\ c\mathbf{x}_\delta \cdot (f_s(\mathbf{x}_0 + c\mathbf{x}_\delta) - f_s(\mathbf{x}_0)) &< 0 \end{aligned} \quad (16)$$

Remembering $f_s(P_{x_0}(t)) = \dot{P}_{x_0}(t)$ from 1, and denoting $\mathbf{x}_\alpha = \mathbf{x}_0 + c\mathbf{x}_\delta$

$$(P_{\mathbf{x}_\alpha}(0) - P_{\mathbf{x}_0}(0)) \cdot (\dot{P}_{\mathbf{x}_\alpha}(0) - \dot{P}_{\mathbf{x}_0}(0)) < 0 \quad (17)$$

Which, by theorem 8.3, proves \mathbf{x}_0 is locally attractive. \square

Finally, we note that linear transforms that satisfy equation 13 are called Negative Definite, a well studied property within linear algebra.

Definition 9.2 (Negative Definite). From Daniel and Noble (1988), a linear transformation denoted by $\mathcal{T}(\mathbf{x}) \in \mathcal{L}_{n,n}$ is said to be Negative Definite if, for each non-zero vector $\mathbf{x} \in \mathbb{R}^n$, the Quadratic Form $\mathbf{x}^T \mathcal{T} \mathbf{x} < 0$.

Therefore, by combining theorems 7.2, 9.1, and definition 9.2, we obtain the final result to answer the original research question:

Theorem 9.2. *For a given path $P_{\mathbf{x}_0}(t)$, if for each point $\mathbf{x}_t = P_{\mathbf{x}_0}(t)$, the linear transformation led by $\mathbf{J}_{f_s}[\mathbf{x}_t]$ is Negative Definite, then $P_{\mathbf{x}_0}(t)$ is Path Stable.*

QUOD ERAT DEMONSTRANDUM.

10 Conclusion

This paper has explored the idea of stability and its standard mathematical representation. Taking Lyapunov's original idea, we have used the same epsilon-delta formalization to expand the concept to include entire paths. To determine if a path meets this definition of stability, it has been shown that if the set of all points on the path meet the criteria for local attractiveness, as defined in the paper, then the path overall must also be stable. In addition, by showing that points where the Quadratic Form of the Jacobian is Negative Determinant are always locally attractive, the paper presents a direct method of calculating the overall stability of a path, and provides areas where this concept can be developed even further, most namely into the complex vector space, as well as into stochastic or time-variant systems.

References

- Daniel, J. W. and Noble, B. (1988). *Applied Linear Algebra*. Englewood Cliffs, NJ, 3rd edition.
- Lyapunov, A. M. (1992). The general problem of the stability of motion. *Control Theory and Applications Series*, 55(3):531–773.
- Taylor, A. E. and Mann, W. R. (1983). *Advanced Calculus*. John Wiley and Sons, Inc., 3rd edition.