PO11Q: Introduction to Quantitative Political Analysis I

Worksheet Week 7 - Solutions

Dr Florian Linke

F.Linke@warwick.ac.uk

a. Billy is looking for the heaviest bag possible and finds one that is 1082 g. What is the probability of finding a heavier bag?

```
\mu = 1000
\sigma = 50
x = 1082
```

Normally distributed, so find a z-score for the observed value. Heavier means right tail.

```
Z = (x - \mu)/\sigma

Z = (1082 - 1000)/50

Z = 1.64
```

Consult tables area under right tail, close to 0.05. Therefore, probability is 5%.

b. What is the probability that Billy will find a bag lighter than 870g?

```
\mu = 1000
\sigma = 50
x = 870
```

Normally distributed so find a z-score for the observed value.

```
Z = (x - \mu)/\sigma

Z = (870 - 1000)/50

Z = -2.6
```

Consult table's area under right tail, probability is equal to 0.0047. For a positive z-score this would indicate the probability of a heavier bag, but because our z score is negative, it shows the probability of a lighter bag. This probability is less than 0.5%.

c. How would the results of a. and b. change if the standard deviation was only 40g?

```
For a.

\mu = 1000

\sigma = 40

x = 1082

Z = (x - \mu)/\sigma

Z = (1082 - 1000)/40

Z = 2.05

Probability is 2% now.

For b.

\mu = 1000
```



```
\sigma = 40

x = 870

Z = (x - \mu)/\sigma

Z = (870 - 1000)/40

Z = -3.25

Probability is now about 0.1%
```

Both of these probabilities are smaller and are a direct reflection of a more narrow distribution.