

The predictive value of NTproBNP on postoperative outcome in patients undergoing offpump CABG

Ibrahim AbuBakr ElSeddiq

July 19, 2020

1 Review of Literature

1.1 Natriuretic Peptides

In 1956, Kisch found that guinea pig atrial, but not ventricular, cells contained highly developed Golgi networks, similar to those observed in secretory cells [Kisch, 1956]. Jamieson and Palade reported that atrial, but not ventricular, myocytes contain spherical, electron opaque granules [Jamieson and Palade, 1964]. At the same time, physiological experiments conducted by Henry and colleagues revealed that balloon distension of the atria correlated with increased urination in dogs [Henry et al., 1956].

In a study published in 1981, de Bold and colleagues elegantly linked the seemingly disparate studies of Kisch and Henry by showing that atrial, but not ventricular, extracts contain a potent blood pressure decreasing component that works by stimulating renal sodium and water secretion [de Bold et al., 1981]. Shortly after the publication of this landmark paper, a number of groups reported the purification and sequencing of atrial peptides of varying sizes that possessed natriuretic, diuretic, and/or smooth muscle relaxing activity [Currie et al., 1984] [Flynn et al., 1983] [Kangawa et al., 1984] [Misono et al., 1984]. Thus, the first natriuretic peptide was discovered.

Several different names were given to these peptides such as atrial natriuretic factor, atriopeptin, cardionatrin, and cardiodilatin. However, atrial natriuretic peptide (ANP) is most often used to describe this peptide in the current literature. The second member of the family to be discovered, B-type natriuretic peptide (BNP), was originally called brain natriuretic peptide because it was purified and sequenced from porcine brain [Sudoh et al., 1988]. However, subsequent studies found that it is more highly concentrated in cardiac ventricles of patients with heart failure [Mukoyama et al., 1991c] [Mukoyama et al., 1990].

Therefore, it is often described as B-type natriuretic peptide today. Finally, the third member of the family, C-type natriuretic peptide (CNP) was purified in 1991 from porcine brain extracts based on its ability to relax smooth muscle. All three members are similar in primary amino acid structure, contain a 17-residue disulfide ring, and are the products of separate genes.[Sudoh et al., 1990]

The biological actions of natriuretic peptides are mediated through membrane-bound natriuretic peptide receptors (NPR) that are linked to a cyclic guanosine monophosphate-dependent signaling cascade, including NPR-A, which preferentially binds ANP and BNP, and NPR-B, which preferentially binds CNP.

Elevated natriuretic peptides levels can be also found in many circumstances involving LV dysfunction or hypertrophy; right ventricular (RV) dysfunction secondary to pulmonary diseases; cardiac inflammatory or infectious diseases; and endocrinologic diseases and high output status without decreased left ventricular ejection fraction (EF), e.g., sepsis, renal failure, cirrhosis of liver, or intracranial pathologies. The causes and mechanisms of elevated natriuretic peptides levels are summarized in Fig. 2.

1.1.1 Atrial Natriuretic Peptide

All natriuretic peptides are synthesized as preprohormones Fig 1. The resulting mRNA gives rise to a 151 amino acid polypeptide, known as preproANP. The first 25 amino acids constitute a signal sequence that is cleaved to yield a 126 amino acid peptide called proANP, which is the major form of ANP stored in the atrial granules [Oikawa et al., 1984]. Upon release from these granules, proANP is rapidly cleaved by corin, a transmembrane cardiac serine protease. Corin is highly expressed on the extracellular surface of atrial cardiomyocytes and cleaves proANP into the biologically active 28-amino acid form of ANP [Yan et al.,




Figure 1: Structure of the human natriuretic peptides. The structure of the preprohormones for ANP, BNP and CNP are outlined on the left of each panel. The final amino acid sequence and structure of the mature peptides along with the major degradation product are shown on the right. The sites of cleavage are indicated with scissors.

2000]. Alternative processing of proANP in the kidney by an unknown protease results in a 32-amino acid peptide called urodilatin that contains four additional amino-terminal residues [Forssmann et al., 1998]. Disruption of the murine ANP gene, results in marked hypertension, which was initially described as salt-sensitive [John et al., 1995b], but later found not to be correlated with dietary salt intake [John et al., 1996].

Release of proANP from the atrial granules is primarily stimulated by stretch of the atrial wall caused by increased intravascular volume [Bilder et al., 1986] [Edwards et al., 1988] [Lang et al., 1985], but pressor hormones also stimulate ANP release [Ruskoaho, 2003c]. Plasma levels of ANP are relatively low (10 fmol/ml), but in patients with congestive heart failure, circulating ANP levels are elevated from 10- to 30-fold [Burnett et al., 1986] [Cody et al., 1986b].

The plasma half-life of ANP in humans is approximately 2 min [Nakao et al., 1986] [Yandle et al., 1986]. Degradation of the active ANP peptide occurs through the actions of neutral endopeptidase (NEP) [Stephenson and Kenny, 1987] [Vanneste et al., 1988] as well as through binding to the natriuretic peptide clearance receptor (NPR-C), a cell surface receptor that lacks guanylyl cyclase activity and controls the local concentrations of natriuretic peptides via constitutive receptor mediated internalization and degradation. Inhibiting NEP, increases the half-life of ANP [Yandle et al., 1989], suggesting that NEP activity contributes to the rapid clearance of ANP. However, it is important to note that mice lacking functional NEP do not exhibit increased natriuretic peptide function [Lu et al., 1995]. In contrast, mice lacking NPR-C are hypotensive, exhibit skeletal overgrowth and have reduced ability to clear ANP compared to wild type mice, suggesting that NPR-C is also a physiologic regulator of circulating natriuretic peptide concentrations [Matsukawa et al., 1999].

ANP is secreted in response to:




Figure 2: The causes and mechanisms of elevated natriuretic peptides levels.

- Stretching of the atrial wall [Widmaier et al., 2008]
- Reduced Sympathetic stimulation of β -adrenoceptors
- Raised sodium concentration (hypernatremia), though sodium concentration is not the direct stimulus for increased ANP secretion. [Widmaier et al., 2008]
- Endothelin, a potent vasoconstrictor
- exercise [Kokkonen et al., 2002]

1.1.2 B-Type Natriuretic Peptide

BNP was initially purified and sequenced from extracts of porcine brain tissue and hence it was named brain natriuretic peptide [Sudoh et al., 1988]. Subsequently, BNP was found at much higher concentrations in cardiac tissues [Mukoyama et al., 1991c] [Mukoyama et al., 1990].

Although low levels of BNP are stored with ANP in atrial granules, BNP is found at greater concentrations in cardiac ventricles. In this tissue, BNP is not stored in granules, but rather transcribed as needed in response to cardiac stress states. [Grepin et al., 1994] [Thuerauf et al., 1994] In normal human subjects, plasma concentrations of BNP are very low (1 fmol/ml), but in response to congestive heart failure, circulating concentrations of BNP are dramatically elevated [Mukoyama et al., 1991c] [Mukoyama et al., 1990].

BNP can be produced in both atria and ventricles, and is upregulated in failing ventricular myocardium. In response to increased myocardial stretch and wall stress, ventricular myocytes secrete the pro-hormone pre-proBNP, which is then cleaved into biologically active BNP and the inactive byproduct N-terminal-proBNP (NT-proBNP). Elevated BNP levels have been demonstrated to be a response to increased angiotensin II and sympathetic tones. [Iwanaga et al., 2006]

BNP is eliminated by binding to the NPR-C or degradation by NEP on endothelial cells, smooth muscle cells, cardiac myocytes, renal epithelium, and fibroblasts. NT-proBNP is cleared mainly by the kidney.[Schrier and Abraham, 1999] Compared to ANP, circulating BNP has a significantly longer half-life of around 20 min in humans [Mukoyama et al., 1991c] [Mukoyama et al., 1990]; the half-life of NT-proBNP is about 60-90 minutes and would be expected to be longer in the setting of renal dysfunction. Unlike ANP, BNP is not initially cleaved by NEP. Instead, the first six amino-terminal amino acids of BNP are first cleaved by the metalloprotease, meprin A in the kidney brush border, which then allows further degradation by NEP [Pankow et al., 2007].

Plastic tubes containing ethylenedinitrotetraacetic acid (EDTA) are desirable for BNP determination and refrigeration is required if the interval between blood collection and analysis is over 4 hours; whereas NT-proBNP can be measured in both serum or plasma, collected in glass or plastic tubes, and has no significant loss of immunoreactivity after 48 hours at room temperature. Although these existing BNP

assays correlate closely, BNP assays are not currently analytical equivalent due to lack of assay standardization. A multicenter collaborative proficiency testing study conducted in 90 Italian laboratories had demonstrated that there are significant differences in analytical characteristics and measured values among the most popular commercial methods for BNP and NT-proBNP. Thus, clinicians should be very careful when comparing results obtained by laboratories that use different methods.

1.1.3 C-Type Natriuretic Peptide

C-type natriuretic peptide (CNP) was initially purified and sequenced from porcine brain extracts [Sudoh et al., 1990]. It is the most highly expressed natriuretic peptide in the brain but is also highly expressed in chondrocytes and endothelial cells. Unlike ANP and BNP, the human gene encoding CNP, NPPC, is not located on chromosome 1 but on chromosome 2 [Ogawa et al., 1994c].

Processing of proCNP to its mature form may occur through the action of the intracellular serine endoprotease, furin. In vitro, furin cleaves the 103 amino acid proCNP into a 53 amino acid carboxyl-terminal biologically active peptide [Wu et al., 2003c]. This 53 amino acid form of CNP (CNP-53) is the major active form of CNP, at the tissue level [Brown et al., 1997]. However, in the systemic circulation, a shorter 22 amino acid form dominates (CNP-22). The protease responsible for this cleavage is not known. Importantly, CNP-53 and CNP-22 appear to bind and activate their cognate receptor, NPR-B, equally well [Yeung et al., 1996].

CNP is not stored in granules and its secretion is increased by growth factors [Suga et al., 1993b] [Suga et al., 1992b] and sheer stress [Chun et al., 1997] in cultured endothelial cells. CNP expression in neonatal vascular smooth muscle cells is increased in response to vascular

injury [Brown et al., 1997]. In normal human subjects, mean CNP concentration is very low (1 fmol/ml). It is elevated in patients with congestive heart failure, although to a much lower extent than ANP and BNP [Charles et al., 2006] [Del-Ry et al., 2005] [Kalra et al., 2003b].

1.2 Natriuretic Peptide Receptors

There are three known natriuretic peptide binding proteins. All members contain a relatively large (450 amino acid) extracellular ligand binding domain and a single membrane-spanning region of about 20 residues. Natriuretic peptide receptors A and B contain an equally large intracellular domain consisting of a so-called kinase homology domain, dimerization domain, and carboxyl-terminal guanylyl cyclase domain. Thus, NPR-A and NPR-B signal by catalyzing the synthesis of the intracellular signaling molecule cGMP. In contrast, NPR-C only contains a 37 residue intracellular domain and lacks guanylyl cyclase activity. It primarily controls local natriuretic peptide concentrations via receptor-mediated internalization and degradation, although many groups have reported signaling functions for NPR-C as well [Rose and Giles, 2008].

1.2.1 Natriuretic Peptide Receptor-A

Natriuretic peptide receptor-A (NPR-A) is the principal receptor of ANP and BNP. NPR-A binds natriuretic peptides at a stoichiometry of 2:1 with a rank natriuretic peptide preference of: $\text{ANP} \geq \text{BNP} > \text{CNP}$ [Bennett et al., 1991] [Koller et al., 1991] [Suga et al., 1992a].

Phosphorylation is essential for activation of NPR-A and dephosphorylation is a mechanism of desensitization in response to prolonged ANP exposure or protein kinase C activation [Potter and Garbers, 1992]




Figure 3: Schematic representation of natriuretic receptors

[Potter and Garbers, 1994]. Although ATP increases ANP-dependent guanylyl cyclase activity, the mechanism for this effect is debatable [Antos et al., 2005] [Antos and Potter, 2007] [Burczynska et al., 2007] [Joubert et al., 2005]. Data indicate that ATP reduces the K_m for NPR-A [Antos and Potter, 2007].

NPR-A internalization and degradation is also controversial. One group consistently reports that the majority of internalized ANP-NPR-A complexes are degraded via a lysosomal pathway with a small portion returning intact to the plasma membrane [Pandey, 2002]. Meanwhile, studies in primary kidney and Chinese Hamster ovary indicate that NPR-A is a membrane resident protein that does not undergo acute internalization and degradation [Fan et al., 2005b] [Koh et al., 1992b] [Vieira et al., 2001b].

NPR-A and/or its mRNA is expressed in kidney, lung, adipose, adrenal, brain, heart, testis, and vascular smooth muscle tissue [Goy et al., 2001]. NPR-A null mice exhibit chronic salt-resistant hypertension and cardiac hypertrophy and fibrosis [Kuhn et al., 2002]. A deletion

in the human NPR-A gene was identified in nine Japanese individuals, of which eight had essential hypertension; the normotensive individual with the altered allele had left ventricular hypertrophy [Nakayama et al., 2000b].

1.2.2 Natriuretic Peptide Receptor-B

Natriuretic peptide receptor-B (NPR-B) is the principal receptor of C-type natriuretic peptide (CNP) and exhibits similar topology, glycosylation, and intramolecular disulfide bonding patterns as NPR-A. NPR-B binds natriuretic peptides with a selectivity preference of $\text{CNP} > \text{ANP} \geq \text{BNP}$ [Bennett et al., 1991] [Koller et al., 1991] [Suga et al., 1992a].

NPR-B dephosphorylation has been shown to mediate desensitization in response to prolonged CNP exposure, protein kinase C activation, and intracellular calcium elevations [Potter, 1998b] [Potter and Hunter, 2000] [Potthast et al., 2004]. ATP increases the guanylyl cyclase activity of NPR-B, by decreasing its Michaelis constant [Antos and Potter, 2007]. NPR-B and/or its mRNA is expressed in bone, brain, fibroblasts, heart, kidney, liver, lung, uterine, and vascular smooth muscle tissue [Bryan et al., 2006] [Dickey et al., 2007]. Mice with a targeted disruption of the NPR-B gene, display dwarfism and female sterility [Tamura et al., 2004].

NPR-B dominant negative mutant transgenic rats have also been generated [Langenickel et al., 2006]. In addition to mild growth retardation of the long bones, the rats displayed progressive, blood pressure-independent cardiac hypertrophy and an elevated heart rate.

Consistent with a prominent role for CNP in the heart, NPR-B, not NPR-A, is the most active natriuretic peptide receptor in the failed heart [Dickey et al., 2007]. Homologous loss-of-function mutations in

human NPR-B result in a rare form of dwarfism called acromesomelic dysplasia, type Maroteaux (AMDM) [Bartels et al., 2004].

1.2.3 Natriuretic Peptide Receptor-C

Natriuretic peptide receptor-C (NPR-C) consists of a large extracellular ligand-binding domain that is approximately 30-35% identical to NPR-A and NPR-B, a single membrane-spanning region, but only 37 intracellular amino acids [Chang et al., 1989] [Fuller et al., 1988] [Porter et al., 1990]. It has no known enzymatic activity but has been suggested to signal in a G protein-dependent manner [Rose and Giles, 2008]. It binds natriuretic peptides with a stoichiometry of two molecules of receptor to one molecule of ligand [Ammarguella et al., 2001]. Its ligand selectivity preference is: ANP > CNP \geq BNP [Bennett et al., 1991] [Suga et al., 1992a].

The main function of NPR-C, also known as the clearance receptor, is to clear circulating natriuretic peptides through the process of receptor-mediated internalization and degradation [Koh et al., 1992b] [Nussenzveig et al., 1990]. Internalization of NPR-C occurs in the absence of ligand; thus, this is a constitutive process [Nussenzveig et al., 1990]. Osteocrin, an endogenous protein with limited homology to members of the natriuretic peptide family, binds NPR-C, but not NPR-A or NPR-B [Moffatt et al., 2007]. Osteocrin is thought to compete with CNP for binding to NPR-C in bone, and therefore, increase local CNP levels during critical periods for bone development [Moffatt et al., 2007].

NPR-C is the most widely and abundantly expressed natriuretic peptide receptor; for example, it constitutes 94% of the total ANP binding sites in endothelial cells [Leitman et al., 1986]. NPR-C and/or its mRNA is expressed in adrenal, brain, heart, kidney, mesentery, and vascular smooth muscle tissue [Nagase et al., 1997] [Porter et al., 1990] [Suga et al., 1992c] [Wilcox et al., 1991]. NPR-C knockout mice exhibit

increased ANP half-lives, long bone overgrowth, hypotension, mild diuresis, dilute urine, and blood volume depletion [Matsukawa et al., 1999]. Mouse strains containing chemically induced loss-of-function mutations in the extracellular domain of NPR-C display skeletal overgrowth from endochondral ossification defects as well [Jaubert et al., 1999].

1.3 Physiologic Effects of Natriuretic Peptides

1.3.1 Natriuretic Peptide Effects on Blood Pressure

ANP binding to NPR-A is a key-signaling pathway, which regulates normal homeostatic blood pressure. This is clearly demonstrated in mice lacking ANP or its receptor NPR-A, which have blood pressures that are elevated 20-40mmHg, compared to control mice [John et al., 1995b] [John et al., 1996] [Lopez et al., 1995] [Oliver et al., 1997]. The link between NPR-A and blood pressure in mice is particularly strong because Smithies and colleagues demonstrated that NPR-A copy number is inversely related to blood pressure in a remarkably linear manner [Oliver et al., 1998]. Conversely, blood pressures in transgenic mice overexpressing ANP or BNP are substantially decreased [Ogawa et al., 1994b] [Steinhilber et al., 1990b]. Although infusion of supraphysiological levels of CNP into animals acutely decreases blood pressure [Clavell et al., 1993] [Sudoh et al., 1990], mice lacking functional CNP or NPR-B are normotensive [Chusho et al., 2001] [Tamura et al., 2004], suggesting that the CNP/NPR-B pathway is not a fundamental regulator of basal blood pressure in mice.

NPR-A dependent decreases in blood pressure are achieved through natriuresis and diuresis, vasorelaxation, increased endothelium permeability, and antagonism of the renin-angiotensin system. Classic experiments showed that atrial extract infusions resulted in rapid renal

excretion of water and sodium [de Bold et al., 1981]. Studies by Garbers and colleagues indicated that the renal response requires NPR-A because mice lacking this receptor do not respond to ANP, BNP, or to acute volume expansion [Kishimoto et al., 1996]. Similar studies found that NPR-A was also required for ANP- or BNP-dependent vasorelaxation in mice [Lopez et al., 1997]. Physiological experiments involving mice with severe reductions of NPR-A in vascular smooth muscle cells demonstrated that while smooth muscle NPR-A is required for acute ANP- or BNP-dependent vasorelaxation, this response does not play a significant role in controlling chronic blood pressure [Holtwick et al., 2002].

The ability of the ANP/NPR-A pathway to increase endothelial permeability is supported by the observation that hematocrit levels are elevated prior to urination and are preserved in nephrectomized animals [Almeida et al., 1986] [Fluckiger et al., 1986] [Richards et al., 1988]. Furthermore, mice with genetically engineered reductions of NPR-A in vascular endothelium exhibit volume expansion, hypertension, and reduced albumin clearance from the vascular system [Sabrane et al., 2005].

1.3.2 Effects of Natriuretic Peptides on Cardiac Hypertrophy and Fibrosis

Although prolonged hypertension can cause hypertrophy, the level of hypertrophy in NPR-A deficient mice is significantly greater than that observed in other genetic models that cause similar levels of hypertension, suggesting that NPR-A elicits a local growth inhibitory signal in the heart. Data for this idea was initially shown in NPR-A knockout mice, which have enlarged hearts even when effectively treated with antihypertensive drugs from birth [Knowles et al., 2001]. Additional studies determined that transgenic re-expression of NPR-A in the hearts of NPR-A $-/-$ mice reduced cardiomyocyte size without affecting heart

rate or blood pressure [Kishimoto et al., 2001]. Finally, mice with reduced cardiomyocyte expression of NPR-A exhibited moderate hypertrophy even though they were slightly hypotensive [Holtwick et al., 2003] [Patel et al., 2005]. In terms of natriuretic peptides, mice lacking ANP have larger hearts, whereas mice transgenically overexpressing ANP have smaller hearts [Barbee et al., 1994] [Steinhilber et al., 1990b]. In contrast, targeted deletion of BNP resulted in normotensive mice with normal heart size but with increased ventricular fibrosis - especially when subjected to pressure overload [Tamura et al., 2000]. Thus, genetic studies in mice strongly support a role for ANP activation of NPR-A in the local inhibition of cardiac hypertrophy and BNP activation of NPR-A in the inhibition of cardiac fibrosis.

Data supporting a role for the CNP/NPR-B pathway in cardiac remodeling has been reported. Although NPR-B inactivation mutations in mice have not been shown to cause hypertrophy [Tamura et al., 2004] [Tsuji and Kunieda, 2005], transgenic rats expressing a dominant negative form of NPR-B exhibit mild blood pressure-independent cardiac hypertrophy and increased heart rate [Langenickel et al., 2006]. In addition, CNP infusion was shown to reduce cardiac remodeling in response to experimentally induced myocardial infarction in rats, and transgenic expression of CNP improved outcomes in mice subjected to ischemia/reperfusion injury or myocardial infarction [Wang et al., 2007].

1.3.3 Effects of CNP and NPR-B on Bone Growth

The most obvious function of the CNP/NPR-B pathway is to stimulate long bone growth. Though undetectable at birth, mice lacking functional CNP or NPR-B develop dwarfism due to impaired endochondrial ossification [Chusho et al., 2001] [Tamura et al., 2001] [Tsuji and Kunieda, 2005]. Conversely, transgenic CNP overexpression or reduced degradation of CNP due to loss-of-function mutations in NPR-C

result in skeletal overgrowth [Jaubert et al., 1999] [Matsukawa et al., 1999] [Yasoda et al., 2004]. Growth plate histology reveals that the endochondral proliferative and hypertrophic zones are reduced in mice with impaired CNP or NPR-B signaling, whereas overexpressing mice have enlarged growth plates [Chusho et al., 2001] [Tamura et al., 2004] [Yasoda et al., 2004].

One cGMP effector involved in the long bone growth pathway is cGMP-dependent protein kinase II, also known as PKGII or cGKII. Loss-of-function mutations in the mouse or rat gene that encodes this kinase also cause dwarfism [Chikuda et al., 2004] [Pfeifer et al., 1996]. Interestingly, the growth plates of rodents with defective cGKII are enlarged, which differs from the diminished growth plates seen in the CNP or NPR-B deleted mice, suggesting that a cGKII-independent pathway is also involved in CNP-dependent long bone growth.

Humans with two loss-of-function alleles for NPR-B suffer from a rare type of autosomal recessive dwarfism, called acromesomelic dysplasia, type Maroteaux [Bartels et al., 2004]. These individuals are characterized by disproportionate limb to torso ratios that are only obvious a year or more after birth. Interestingly, although single copy carriers of a nonfunctional NPR-B allele do not suffer from disease, they are statistically shorter than comparable individuals with two wild type NPR-B alleles [Olney et al., 2006]. Thus, it is possible that NPR-B mutations could have a significant effect on the stature of the general population.

1.4 Biological Action of CNH

Cardiac natriuretic hormones have powerful physiological effects on the cardiovascular system, body fluid, and electrolyte homeostasis [McGrath and de Bold AJ, 2005] [Clerico and Emdin, 2004] [Clerico, 2002a] [de Lemos et al., 2003b] [Ruskoaho, 2003a]. CNH share a direct di-

uretic, natriuretic and vasodilator effect and an inhibitory action on ventricular myocyte contraction [Zhang et al., 2005] as well as remodeling and inflammatory processes of myocardium and smooth muscle cells [Nagaya et al., 1998] [Hayashi et al., 2001] [Magga et al., 2004] [Hardt and Sadoshima, 2004] (Fig. 3.14). Thus, CNH exert a protective effect on endothelial function by decreasing shear stress, modulating coagulation and fibrinolysis pathways, and inhibiting platelet activation (Fig. 3.15). They can also inhibit vascular remodeling process as well as coronary restenosis post-angioplasty [Ma et al., 2004] [Chen and Burnett] [Morishige et al., 2000] [Takeuchi et al., 2003] [Yasuda et al., 2002] [Qian et al., 2002] [Nakanishi et al., 2005].

CNH share an inhibitory action on neuro-hormonal and immunological systems, and on some growth factors. In particular, the pivotal role of CNH (especially ANP) in modulating the immune response has been reviewed [Kapoun et al., 2004]. The first evidence for a role of CNH in the immune system was given by the fact that peptide hormones and their receptors are expressed in various immune organs. Furthermore, several studies indicated that the CNH system in immune cells underlies specific regulatory mechanisms by affecting the innate as well as the adaptive immune response. In particular, ANP increases phagocytotic activity and production of reactive oxygen species of phagocytes. ANP affects the induced innate immune response by regulating the activation of macrophages at various stages. It also reduces production of pro-inflammatory mediators by inhibition of iNOS and COX-2 as well as TNF- α synthesis. *ANP also affects TNF- α action, i.e. it interferes with the inflammatory effects of TNF- α on the endothelium. α -induced endothelial permeability and adhesion and attraction of inflammatory cells.*

The cited effects on the cardiovascular system and body fluid and electrolyte homeostasis can be explained at least in part by the inhibition of control systems, including the sympathetic nervous system, the renin-angiotensin-aldosterone system (RAAS), the vasopressin/antidiuretic hormone system, the endothelin system, cytokines and growth factors [Ventura et al., 2002] [Vatta et al., 1996] [Vatta et al., 1997] [Fermepin

et al., 2000] [Rocca et al., 2001] [Tsukagoshi et al., 2001] [Kierner et al., 2002b] [Weber et al., 2003] [Kapoun et al., 2004] [Vollmar, 2005]. The endocrine action, shared by plasma ANP and BNP, can be enhanced by natriuretic peptides produced locally in target tissues (paracrine action). Indeed, endothelial cells synthesize CNP, which in turn exerts a paracrine action on vessels [Woodard et al., 2002] [Chen and Burnett] [Morishige et al., 2000] [Takeuchi et al., 2003] [Yasuda et al., 2002] [Qian et al., 2002]. Moreover, renal tubular cells produce urodilatin, another member of the peptide natriuretic family, which has powerful diuretic and natriuretic properties [Vesely, 2003a]. Genes for natriuretic peptides (including ANP, BNP and CNP) are also expressed in the central nervous system, where they likely act as neurotransmitters and/or neuromodulators [Vatta et al., 1996] [Vatta et al., 1997] [Fermepin et al., 2000] [Vesely, 2003a] [Imura et al., 1992] [Langub et al., 1995]. In particular, it was demonstrated that intranasal ANP acts as central nervous inhibitor of the hypothalamus-pituitary-adrenal stress system in humans [Perras et al., 2004]. Finally, co-expression of CNH and of their receptors was observed in rat thymus cells and macrophages [Vollmar et al., 1995] [Vollmar and Schultz, 1995], suggesting that CNH may have immunomodulatory and anti-inflammatory functions in mammals [Vollmar and AK, 2001].

A detailed review [Waschek, 2004] has highlighted a possible major role for CNH in the development of certain systems, in particular skeleton, brain, and vessels. This review cites studies showing severe skeletal defects and impaired recovery after vascular and renal injury in CNH transgenic and knockout (KO) mice [108, 205]. In addition, CNH may have a role in the regulation of proliferation, survival, and neurite outgrowth of cultured neuronal and/or glial cells [108, 205].

Changes in plasma ANP are also correlated with alcohol-associated psychological variables [108, 205]. Acute administration of alcohol stimulates the release of ANP independently of volume-loading effects. Patients whose ANP levels fell markedly during abstinence also reported more intense and frequent craving as well as more anxiety

[108, 205]. Several reports have shown that CNH stimulate the synthesis and release of testosterone in a dose-dependent manner in isolated and purified normal Leydig cells [?]. It has been suggested that this effect on normal Leydig cell steroidogenesis does not involve classical mechanisms of cAMP-mediated regulation of steroidogenic activity by gonadotropins [Pandey, 2005]. The stimulated levels of testosterone production by ANP, BNP, and gonadotropins were comparable, whereas CNP has been found to be a weak stimulator of testosterone production in Leydig cells [Pandey, 2005]. Moreover, testicular cells contain immunoreactive ANP-like materials and a high density of natriuretic peptide receptor-A (NRP-A) [Pandey, 2005]. These findings suggest that CNH play paracrine and/or autocrine roles in testis and testicular cells. Furthermore, the presence of ANP and its receptors has been reported in ovarian cells, too. Increasing evidence strongly support that CNH are present and probably locally synthesized in ovarian cells of different mammalian species and also play an important physiological role in stimulating estradiol synthesis and secretion in the female gonad [Pandey, 2005] [Gutkowska et al., 1993] [Vollmer et al., 1988] [Ivanova et al., 2003]. However, further studies are necessary in order to clarify completely the role played by CNH in the regulation of gonadal function and also to assess the inter-relationship between heart endocrine function and gonadal function in humans.

The huge amount of data reported above strongly supports the hypothesis that CNH are active components of the body integrative network that includes nervous, endocrine and immune systems. According to this hypothesis, the heart can no longer be seen as a passive automaton driven by nervous, endocrine or hemodynamic inputs, but as a leading actor on the stage. Thus, CNH, together with other neuro-hormonal factors, regulate cardiovascular hemodynamics and body fluid and electrolyte homeostasis, and probably modulate inflammatory response in some districts, including the cardiovascular one. This hypothesis implies that there are two counteracting systems in the body: one has sodium-retaining, vasoconstrictive, thrombophylic, pro-inflammatory and hypertrophic actions, while the second one promotes natriuresis

and vasodilatation, and inhibits thrombosis, inflammation and hypertrophy. CNH are the main effectors of the latter system, and work in concert with NO, some prostaglandins, and other vasodilator peptides (such as bradykinin) [Chatterjee et al., 1999] [Ruskoaho et al., 1997] [Gyurko et al., 2000a] [Tanaka et al., 2005] [Booz, 2005]. Under physiological conditions, the effects of these two systems are well balanced via feedback mechanisms, and result in a beat-to-beat regulation of cardiac output and blood pressure in response to endogenous and exogenous stimuli. In patients with HF, the action of the first system is predominant, as a compensatory mechanism, initially, that progressively leads to detrimental effects.

The knowledge so far accumulated regarding CNH suggests that a continuous and intense information exchange flows from the endocrine heart system to nervous and immunological systems and to other organs (including kidney, endocrine glands, liver, adipose tissue, immunocompetent cells) and vice versa (Fig. 3.16). From a pathophysiological point of view, the close link between the CNH system and counter-regulatory systems could explain the increase in circulating levels of CNH in some non-cardiac-related clinical conditions. Increased or decreased BNP levels were frequently reported in acute and chronic respiratory diseases [Kruger et al., 2004a] [Pruszczyk et al., 2003] [Kucher et al., 2003b] [ten Wolde et al., 2003b] [Ando et al., 1996] [Maeder et al., 2003] [Nagaya et al., 2000a] [Leuchte et al., 2004b] [Leuchte et al., 2004a], some endocrine and metabolic diseases [Kohno et al., 1993] [Bernstein et al., 1997] [Parlapiano et al., 1998] [Schultz et al., 2004] [Jensen et al., 1997] [Sugawara et al., 1988] [Fujio et al., 1989] [Opocher et al., 1990] [Lapinski et al., 1991] [Cappuccio et al., 1989] [Straub et al., 1996] [Bhalla et al., 2004b], liver cirrhosis [Villa et al., 1995] [Salo et al., 1996] [Henriksen et al., 2003], renal failure [Vesely, 2003a] [Henriksen et al., 2003], septic shock, chronic inflammatory diseases [McCullough et al., 2004b] [Castillo et al., 2004b] [Witthaut et al., 2003] [Takemura et al., 1998] [Palladini et al., 2003b], subarachnoid hemorrhage [McGirt et al., 2004] [Fukui et al., 2003] [Kurokawa et al., 1996] [Fukui et al., 2004], and some paraneoplastic syndromes

[Mimura et al., 1996] [Marchioli and SL, 1997] [Johnson et al., 1997]. In addition, any myocardial damage leading to the release of sarcoplasmic constituents (including CNH) in extracellular fluid, for instance that due to cardiotoxic agents [Hayakawa et al., 2001a] [Suzuki et al., 1998] [Nousiainen et al., 1998b] [Nousiainen et al., 1999b] [Okumura et al., 2000], cardiac trauma or ischemic necrosis [Mukoyama et al., 1991b] [Morita et al., 1993], also causes an increase in plasma concentration of CNH.

Furthermore, the inter-relationships between the CNH system and pro-inflammatory cytokines suggest that cardiac hormones play an important role in mechanisms responsible for cardiac and vascular adaptation, maladaptation and remodeling in response to various physiological and pathological stimuli [Tenhunen et al., 2005] [Walther et al., 2003] [Glembotski et al., 1993] [Mukoyama et al., 1991b]. Elevated BNP levels in extra-cardiac diseases reveal an endocrine heart response to a “cardiovascular stress” (Fig. 3.17). Indeed, studies reported that plasma BNP concentration is an independent risk factor for mortality (cardiac and/or total) in pulmonary embolism [Kruger et al., 2004a] [Kucher et al., 2003b] [ten Wolde et al., 2003b] and hypertension [Nagaya et al., 2000a], renal failure [Clerico and Emdin, 2004] [Vesely, 2003a] [Henriksen et al., 2003], septic shock [McCullough et al., 2004b], amyloidosis [Palladini et al., 2003b], and diabetes mellitus [Bhalla et al., 2004b] (see Chapter 6 for more details). According to this hypothesis, a BNP assay should be considered as a marker of cardiac stress (Fig. 3.17).

In conclusion, CNH share a powerful action on the cardiovascular system, including diuretic, natriuretic and vasodilator effects and an inhibitory action on ventricular myocyte contraction, as well as on remodeling and inflammatory processes of myocardium and smooth muscle cells. Furthermore, CNH exert a protective effect on endothelial function by decreasing shear stress, modulating coagulation and fibrinolysis pathways, and inhibiting platelet activation. They can also inhibit the vascular remodeling process as well as coronary resteno-

sis post-angioplasty. These effects can be explained, at least in part, by the inhibition of control systems, including the sympathetic nervous system, the RAAS, the vasopressin/antidiuretic hormone system, the endothelin system, cytokines and growth factors. Finally, the endocrine action of ANP and BNP is potentiated at the periphery (target tissues) by the paracrine action of other members of the peptide natriuretic family, such as CNP (in the vascular tissue) and urodilatin (in renal tissue). Finally, some experimental studies performed in KO mice suggest a distinct pathophysiological role for BNP in respect to ANP [LaPointe, 2005]. While BNP KO mice are no different from control mice with regard to blood pressure, urine volume, and urinary electrolyte excretion, they have more extensive ventricular fibrosis, accompanied by increased transforming growth factor- β 3 (TGF- β 3) and collagen mRNA [LaPointe, 2005]. These data suggest that BNP may function more as an autocrine/paracrine inhibitor of cell growth in the heart; while ANP may be considered as a traditional circulating hormone with pronounced diuretic, natriuretic, and antihypertensive effects.

1.5 Natriuretic Peptide Receptors and Intracellular Second Messenger Signaling

Cardiac natriuretic hormones share their biological action by means of specific receptors (NPR), which are present within the cell membranes of target tissues. Three different subtypes of NPRs have so far been identified in mammalian tissues [Pandey, 2005] [Anand-Srivastava and GJ, 1993] [Kuhn, 2004].

1.5.1 Genes Encoding for NPRs

NPR1 is the gene coding the NPR-A receptor (natriuretic peptide receptor A/guanylate cyclase A) and it is located on 1q21-q22 spanning 15,534 bp with 22 exons. The relative mRNA of 3,805 bp leads to a protein of 1,061 amino acids. The NPR-B receptor (natriuretic peptide receptor B/guanylate cyclase B) is codified by the gene NPR2. This gene of 17,303 bp is on chromosome 9 (9p21-p12) and it is organized in 22 exons, which can give two types of mRNA. NPR2 Ia is an mRNA of 3,482 bp that has a 71 nucleotide insertion relative to isoform b, which results in a different, and shorter (995 aa), carboxy-terminus that may disrupt the guanylyl cyclase activity. NPR2 Ib (3,411 bp, 1,047 aa) does not include the alternate exon found in isoform a, and thus isoform b contains a longer carboxy-terminus. The natriuretic peptide receptor C gene, also named NPR3, is on 5p14-p13 and spans 74,698 bp (8 exons), giving an mRNA of 1,753 bp that is translated into a protein of 540 amino acids.

1.5.2 Biological Function of NPRs

NPR-A and NPR-B are generally considered to mediate all known biological actions throughout the guanylate cyclase (GC) intracellular domain, while the third member of the natriuretic peptide receptor family, the NPR-C receptor, does not have a GC domain (Figs. 3.18, 3.19 and 3.20). The GC receptors for ANP/BNP (NPR-GC-A) and CNP (NPR-GC-B) belong to a family of seven isoforms of transmembrane enzymes (from GC-A to GC-G), which all convert guanosine triphosphate into the second messenger cyclic 3',5'-guanosine monophosphate (cGMP) [Anand-Srivastava and GJ, 1993].

Although partly homologous to soluble GC, the receptor for NO, the membrane GCs share a different and unique topology. The single

transmembrane span domain divides the protein structure into an extracellular ligand binding domain and an intracellular region consisting of a protein kinase-homology domain, an amphipathic helical or hinge region, and a cyclase-homology domain [Kuhn, 2004] (Figs. 3.18, 3.19 and 3.20). The cyclase homology domain represents the catalytic cGMP synthesizing domain. The function of the intracellular region consisting of a protein kinase-homology domain is incompletely understood. Although it probably binds ATP and contains many residues conserved in the catalytic domain of protein kinases, kinase activity has not been detected [Kuhn, 2004]. It represses the enzyme activity of the catalytic cGMP-synthesizing domain and at the same time is necessary for its ligand-dependent activation [Mimura et al., 1996]. The coiled-coil hinge region is involved in receptor dimerization, which is also essential for the activation of the catalytic domain [Kuhn, 2004].

The cGMP produced modulates the activity of specific downstream regulatory proteins, such as cGMP-regulated phosphodiesterases, ion channels and cGMP-dependent protein kinases type I (PKG I) and type II (PKG II) (Fig. 3.20). These proteins should be considered to be third messengers, which are differentially expressed in different cell types, ultimately modifying cellular functions [Lohmann et al., 1997] [Pfeifer et al., 1999]. This specific action of CNH on target tissues depends essentially on two different mechanisms.

The physiological expression of NPR-A and NPR-B differs quite significantly in human tissues (Fig. 3.21). NPR-A is found in abundance in larger, conduit blood vessels, whereas the NPR-B is found predominantly in the central nervous system [Ahluwalia et al., 2004a]. Both receptors have been localized in adrenal glands and kidney [Ahluwalia et al., 2004a]. On the other hand, several studies indicate that phosphorylation of the kinase homology domain is a critical event in the regulation of NPRs [Potter and Hunter, 1998a] [Potter and Hunter, 1998b] [Potter, 1998a].

The affinity for ANP, BNP and CNP also varies greatly among the

different NPRs. ANP shows a greater affinity for NPR-A and NPR-C, and CNP for NPR-B, while BNP shows a lower affinity for all NPRs compared to the other two peptides (Fig. 3.21). Activation of the GC-linked NPRs is incompletely understood [Fan et al., 2005a]. NPR-A and NPR B are homo-oligomers in the absence and presence of their respective ligands, indicating that receptor activation does not simply result from ligand-dependent dimerization [Chinkers and EM, 1992]. However, ANP binding does cause a conformational change of each monomer closer together [Fan et al., 2005a] [Chinkers and EM, 1992] [Huo et al., 1999] [Labrecque et al., 1999] [Labrecque et al., 2001]. The stoichiometry of the ligand-receptor complex is 1:2 [Ogawa et al., 2004]. Initial in vitro data suggested that direct phosphorylation of NPR-A by protein kinase C mediated its “desensitization” (i.e., the process by which an activated receptor is turned off) [Duda and RK, 1990]. However, subsequent studies conducted in live cells indicated that desensitization in response to prolonged natriuretic peptide exposure or activators of protein kinase C results in a net loss of phosphate from NPR-A and NPR-B [Potter, 1998a] [Potter and DL, 1992] [Potter and DL, 1994] [Foster and DL, 1998] [Joubert et al., 2001].

Although ligand-dependent internalization and degradation of NPR-A has been intensely studied by several groups for many years, a consensus understanding of the importance of this process in the regulation of NPRs has not emerged [Joubert et al., 2001]. Early studies conducted on PC-12 pheochromocytoma cells suggested that both NPR-A and NPR-C internalize ANP and that both receptors are recycled back to the cell surface [Pandey et al., 1986]. Other studies, using Leydig, Cos, and 293 cell lines, have reported that ANP binding to NPR-A stimulates its internalization, which results in the majority of the receptors being degraded with a smaller portion being recycled to the plasma membrane [Pandey et al., 1986] [Pandey, 1993] [Pandey et al., 2000] [Pandey, 2001]. In contrast, other studies performed in cultured glomerular mesangial and renomedullary interstitial cells from the rat or Chinese hamster ovary cells reported that NPR-A is a constitutively membrane resident protein that neither undergoes endocytosis nor me-

diates lysosomal hydrolysis of ANP [Koh et al., 1992a] [Vieira et al., 2001a]. A more study using 293T cells suggested that NPR-A and NPR B are neither internalized nor degraded in response to receptor occupation [Chinkers and EM, 1992]. Furthermore, this study did not support the hypothesis that down-regulation is responsible for NPR desensitization observed in response to various physiological or pathological stimuli [Joubert et al., 2001]. Further studies are necessary to clarify whether or not ANP binding to NPR-A stimulates its internalization, and whether this process is tissue- and/or species-specific.

It is generally thought that the NPR-C is not linked to GC and so serves as a clearance receptor [Clerico and Emdin, 2004] [de Lemos et al., 2003b] [Ruskoaho, 2003a]. NPR-C is present in higher concentration than NPR-A or NPR-B in several tissues (especially vascular tissue), and it is known constitutively to internalize CNH [Fan et al., 2005a] (Fig.3.22). However, studies have found that CNH interact with NPR-C to suppress the cAMP concentration by inhibition of adenylyl cyclase [MB, 2005] [Drewett et al., 1992]. Specific binding to NPR-C increases inositol triphosphate and diacylglycerol concentrations by activating phospholipase C activity or inhibits DNA synthesis stimulated by endothelin, platelet-derived growth factor and phorbol ester by inhibiting MAPK activity, as reviewed [MB, 2005]. The NPR-C mediated inhibition of adenylyl cyclase is mediated through Gi (inhibitory guanine nucleotide regulatory) proteins. According to this hypothesis, NPR-C, which is present in large amounts, especially on the endothelial cell wall, may mediate some paracrine effects of CNP on vascular tissue [Ahluwalia et al., 2004a] [MB, 2005]. However, further studies are necessary to elucidate the possible role of NPR C receptors as modulators of CNH action and/or degradation in peripheral tissues.

1.6 Metabolic Pathways and Circulating Levels of CNH

Atrial natriuretic peptide and BNP are secreted directly from the heart. In the circulation, CNHs are metabolized via two principal mechanisms: degradation by a membrane-bound endopeptidase (NEP 24.11) and receptor-mediated cellular uptake via NPR-C [Goetze, 2004a] (Fig. 3.22). Some biological characteristics of ANP, BNP and CNP (as well as of their precursors) are summarized in Table 3.3.

1.6.1 ANP Metabolism

Atrial natriuretic peptides are a family of peptides derived from a common precursor, called preproANP, which in humans contains 151 amino acids and has a signal peptide sequence at its amino-terminal end (Fig. 3.11). The pro-hormone is stored in secretion granules of cardiomyocytes as a 126-amino-acid peptide, proANP 1-126, which is produced by cleavage of the signal peptide. When appropriate signals for hormone release are given, proANP 1-126 is further split by some proteases (especially the serine protease corin) [D'Souza et al., 2004] into N-terminal fragment NT-proANP and the COOH-terminal peptide ANP, which is generally considered to be the biologically active hormone, because it contains the cysteine ring (Figs. 3.1 and 3.11).

Studies from the group of Vesely et al. suggested that the NT-proANP can be metabolized in vivo in three peptide hormones with blood pressure-lowering, natriuretic, diuretic and/or kaliuretic properties [Vesely, 2003a]. These peptide hormones, numbered by their amino acid sequences, beginning at the N-terminal end of the proANP pro-hormone, include: 1) the peptide proANP 1-30, also called long-acting natriuretic peptide (LANP); 2) the peptide proANP 31-67 with vessel dilator properties; 3) the peptide proANP 79-98 with kaliuretic properties

[Kapoun et al., 2004]. However, these three peptides do not bind to the same NPRs of CNHs, because they do not have the cysteine ring. Further studies are necessary to confirm and elucidate the biological action of these putative peptide hormones, as well as their in vivo metabolism.

There is some evidence that ANP is secreted according to a pulsatile pattern in humans [Haak et al., 1990] [Haak et al., 1992] [Nugent et al., 1994] [Pedersen et al., 1999] [Bentzen et al., 2003]. Upon secretion, ANP is rapidly distributed and degraded (the metabolic clearance rate of ANP is on average about 2,000 ml/min in healthy subjects) with a plasma half-life of about 4-6 minutes in healthy adult subjects. In humans, about 50% of the ANP secreted into the right atrium is extracted by the peripheral tissues during the first pass throughout the body [Pilo et al., 1998] [Iervasi et al., 1998] [Clerico and Iervasi, 1995] [Iervasi et al., 1997]. Furthermore, circulating ANP represents only a small fraction of the total body pool (no more than 1/15) in normal subjects and plasma ANP concentration shows rapid and wide fluctuations in healthy subjects, even at rest in the recumbent position [Pilo et al., 1998] [Iervasi et al., 1998] [Clerico and Iervasi, 1995] [Iervasi et al., 1997]. The turnover data suggest that circulating levels of ANP may not represent a close estimate of their disposal, and therefore of the activity of the CNH system, as implicitly accepted in physiological or clinical studies in which only the plasma concentration of the hormone is measured, without an estimation of turnover rate. However, it was demonstrated that ANP clearance mechanisms are constant in the presence of rapid and large changes in endogenous ANP plasma levels induced by atrial and/or ventricular pacing, thus indicating that, at least for studies lasting only a few hours, changes in ANP circulating levels may provide a reliable estimate of production rate variations [Iervasi et al., 1997].

1.6.2 BNP Metabolism

The biological action, metabolic pathways, and turnover parameters of BNP are not as well known as those of ANP [Goetze, 2004a]. However, it is commonly believed that the BNP turnover is less rapid than that of ANP with a plasma half-life of about 13-20 minutes; indeed, circulating levels of BNP are more stable than those of ANP in adult healthy subjects (Fig. 3.23). Bentzen et al. [Bentzen et al., 2003] analyzed the secretion pattern of ANP and BNP in 12 patients with chronic HF and in 12 healthy adult subjects. ANP and BNP in plasma were determined by radioimmunoassay (RIA) at 2 min intervals during a 2-h period and were subsequently analyzed for pulsatile behavior using the method of Fourier transformation. All patients and healthy subjects had significant rhythmic oscillations in plasma ANP levels, and 11 patients with HF and 10 healthy subjects had significant rhythmic oscillations in plasma BNP levels [Bentzen et al., 2003]. The amplitude of the main frequency was considerably higher in patients than in healthy subjects, but the main frequency did not differ significantly between patients and healthy subjects for either ANP or BNP. Patients with HF demonstrated pulsatile secretion of ANP and BNP with a much higher absolute amplitude, but with the same main frequency as healthy subjects [Bentzen et al., 2003]. Finally, rhythmic oscillations in plasma ANP levels of healthy subjects showed significantly higher mean amplitude, but not frequency, than those of BNP [Bentzen et al., 2003].

A very small amount of immunoreactive BNP has been found in urine [Totsune et al., 1996] [Ng et al., 2004], but the precise mechanism of renal excretion has not yet been fully clarified. In contrast to BNP, the biologically active peptide, other proBNP-derived inactive fragments also circulate in plasma. These fragments are commonly referred to as “N-terminal proBNP” (NT-proBNP), but the molecular heterogeneity also includes the intact precursor, particularly in patients with HF [Goetze, 2004a] [Shimizu et al., 2003b]. Cardiac secretion of proBNP

and its N-terminal fragments has been demonstrated by blood sampling from the coronary sinus [Hunt et al., 1997]. Some data suggest that the major part of proBNP produced in myocytes is apparently processed prior to release [Goetze, 2004a]; however, intact proBNP peptide was also found in plasma of patients with HF as well as healthy adult subjects [Goetze, 2004a] [Hunt et al., 1997] [Goetze et al., 2005].

A study, employing a new method for the total and equimolar assay of all proBNP-related peptides (i.e., intact proBNP precursor plus NT-proBNP concentrations), found comparable peripheral concentrations of BNP (measured by immunoradiometric assay) and proBNP-related peptides in patients with HF [Goetze et al., 2005]. Moreover, the BNP concentration (median 125 pmol/l) was higher than that of total proBNP (103 pmol/l) in the coronary sinus, suggesting that the cardiac secretion of these two peptides could be different [Goetze et al., 2005]. Alternatively, this finding could also reflect some difference in peripheral elimination of peptides because total proBNP concentration is significantly higher in the pulmonary artery than the aortic root in patients with right ventricular failure [Goetze et al., 2004b].

While NEP enzymes are mainly involved in natriuretic peptide inactivation in vivo, the degradation of BNP seen in vitro is most likely due to other enzymes, such as peptidyl arginine aldehyde proteases, kallikrein, and serine proteases [Belenky et al., 2004b]. However, the role of these enzymes in the degradation of BNP in vivo is unclear.

A study reported that both the BNP and total proBNP concentrations were increased more than 2-fold in the coronary sinus compared to the inferior caval vein (BNP-32: median 125 pmol/l, range 21-993 vs median 52 pmol/l, range 7-705; proBNP: median 103 pmol/l, range 16-691 vs 47 pmol/l, 8-500) [Goetze et al., 2005]. These findings are in accordance with previous studies suggesting that the cardiac gradient for BNP secretion (as estimated by the difference between BNP concentration in coronary sinus and inferior caval vein) ranges from 1.6-fold

to 2.9-fold [Shimizu et al., 2003b] [Mukoyama et al., 1991a] [Mizuno et al., 2001] [Kalra et al., 2003a]. Taking these studies as a whole, ANP and BNP share a similar peripheral extraction value (of about 30-50%). Further studies are necessary to elucidate the metabolism of BNP and in particular the predominant form of the circulating BNP-related peptides.

1.7 CNH Genes and Cardiovascular Diseases

Since CNH have a potent diuretic antihypertensive action, and the impaired action of the peptides may cause hypertension, their genes may be candidates for cardiovascular disease, especially arterial hypertension. Furthermore, transgenic animals (especially mice), overexpressing CNH or knockout for ANP/BNP genes or their specific receptors, have been used to evaluate the pathophysiological role of the CNH system in cardiovascular diseases [Nakayama, 2005].

In transgenic mice with overexpression of ANP and BNP in liver, plasma ANP and BNP levels are from 10- to 100-fold higher than in control mice, with a blood pressure of 20-25 mmHg lower. These mice also have lighter hearts, but with the same cardiac output and rate, than controls [Nakayama, 2005] [Steinhilber et al., 1990a] [Ogawa et al., 1994a] [John et al., 1995a]. The BNP-overexpressing mice show the same hemodynamic changes; on the other hand, ANP KO mice develop NaCl-sensitive hypertension [Nakayama, 2005]. Transgenic mice overexpressing the NPRA gene have also been created; these animals have a lower blood pressure than wild-type mice [Nakayama, 2005]. The corresponding KO mice show an increase in blood pressure compared with controls (on average 10 mmHg in heterozygous and 30 mmHg in homozygous animals), which is not affected by NaCl intake [Ogawa et al., 1994a] [John et al., 1995a]. These data suggest a different pathophysiological mechanism for hypertension between KO mice for the ANP gene and its specific receptor; this difference does

not yet have an explanation [Nakayama, 2005]. NPRC heterozygous KO mice do not show blood pressure variation, whereas homozygous mice show on average a decrease in blood pressure of about 8 mmHg [Nakayama, 2005].

The function of natriuretic peptides was also studied after induction of myocardial infarction in KO mice lacking the natriuretic peptide receptor guanylyl cyclase-A, the receptor for ANP and BNP [Nakanishi et al., 2005]. KO and wild-type mice were subjected to left coronary artery ligation and then followed-up for 4 weeks. KO mice showed significantly higher mortality because of a higher incidence of acute HF, which was associated with diminished water and sodium excretion and with higher cardiac levels of mRNAs encoding ANP, BNP, TGF- β 1, and type I collagen. By 4 weeks after infarction, left ventricular remodeling, including myocardial hypertrophy and fibrosis, and impairment of left ventricular systolic function were significantly more severe in KO than wild-type mice [Nakanishi et al., 2005]. These data confirm that the CNH system has powerful anti-remodeling properties on ventricular cardiomyocytes.

In years, molecular genetic techniques have been introduced in etiological studies of polygenetic diseases, in linkage studies, in sib-pair linkage studies of various candidate genes, and in related studies [Nakayama, 2005] [Clerico, 2003]. The association between some abnormalities in genes, coding for the CNH and their receptors, and some cardiovascular (in particular hypertension) and metabolic (such as diabetes mellitus) diseases has been tested in a large number of clinical studies (see also Chapter 6 for more details). To date, the results obtained are conflicting and seem to depend strictly on the ethnic population of the study. The restriction fragment length polymorphism for the enzyme HpaII, located in intron 2 of NPPA (polymorphism also called Sma I), was reported to be more common in hypertensive African-Americans than in normotensive black controls [Rutledge et al., 1995]; these data were then confirmed in two [Ramasawmy et al., 1993] [Schorr et al., 1997], but not a third [Widecka et al., 1998], Caucasian populations.

Furthermore, another study found that the HpaII polymorphism was not associated with hypertension in the Chinese population of Hong Kong [Cheung et al., 1999].

Regarding other NPPA polymorphisms, Japanese studies reported that both G1837A and T2238C polymorphisms are associated with essential hypertension [Rahmutula et al., 2001], while only a marginally significant association was found with an ANP polymorphism located in the 5'-untranslated region (C664G) [Kato et al., 2000].

Several allelic variants have also been described for genes coding for CNH receptors (see the review by Nakayama [Nakayama, 2005] for a more detailed discussion of this topic). The clearance receptor for natriuretic peptides (NPR-C) is highly expressed in adipose tissue, and its bi-allelic (A/C) polymorphism was detected at position -55 in the conserved promoter element named P1. This variant of the NPR-C P1 promoter is associated with lower ANP levels and higher systolic blood pressure and mean blood pressure in obese hypertensives: the C(-55) variant, in the presence of increased adiposity, might reduce plasma ANP through increased NPR-C-mediated ANP clearance, contributing to higher blood pressure [Sarzani et al., 1999].

In the Japanese population an insertion/deletion (GCTGAGCC) polymorphism has been identified in the 5'-flanking region of the NPRA gene that is associated with essential hypertension and left ventricular hypertrophy [Nakayama et al., 2000a]. Another insertion/deletion polymorphism is on the 3'-untranslated region of the NPRA gene, on exon 22, and it seems to be associated with familial hypertension [Lucarelli et al., 2001]. However, these data should be confirmed in larger studies, including other ethnic populations.

3.9 An Integrated Neuro-Hormonal System Regulates Vascular Function

Endothelial cells release an array of vasoactive mediators that alter

the tone and growth of the underlying smooth muscle and regulate the reactivity of circulating white blood cells, erythrocytes and platelets. These endogenous factors are usually called endothelium-derived vasorelaxant mediators [Ahluwalia and AJ, 2005]. Moreover, it appears that alterations in the capacity of the endothelium to release some mediators in response to pathophysiological stimuli (the so-called endothelium dysfunction) are a major precipitating factor in many cardiovascular diseases. Perhaps the most important of these paracrine mediators are prostacyclin (PGI₂) and nitric oxide (NO). Moreover, a third endothelium-derived vasorelaxant mediator has been described [Ahluwalia and AJ, 2005]. This is termed endothelium-derived hyperpolarizing factor (EDHF) because it elicits a characteristic smooth muscle hyperpolarization and relaxation. Much attention has focused on identifying EDHF(s), with diverse candidates, including cytochrome P450 metabolites, KC ions, anandamide and hydrogen peroxide [Ahluwalia and AJ, 2005]. However, the role of each of these as EDHF remains unsubstantiated.

There is now compelling evidence that CNH (and especially CNP) act as EDHFs in some vascular beds [Ahluwalia and AJ, 2005] [Han and Hasin, 2003] [Houben et al., 2005] [Scotland et al., 2005]. Indeed, numerous studies have demonstrated that ANP, BNP and CNP bind to NPR-A and NPR-B receptors on vascular smooth muscle cells (either freshly isolated or in culture), stimulate cGMP accumulation, and cause a dose dependent vasodilation [Han and Hasin, 2003] [Houben et al., 2005] [Scotland et al., 2005]. This increase in cGMP causes vasodilatation by reducing intracellular calcium levels, as occur when cGMP accumulation is stimulated by NO and its analogs.

It is theoretically conceivable that ANP and BNP act like hormones in vascular tissue by reaching the smooth muscle cells from the circulation after secretion by the heart, while CNP shows a paracrine action, being secreted by endothelial cells [Woodard et al., 2002] [Chen and Burnett] [Yasuda et al., 2002] [Qian et al., 2002] (Fig. 3.15). However, Casco et al. [Casco et al., 2002] demonstrated the existence of a com-

plete CNH system (including the production and secretion of ANP, BNP and CNP) in atherosclerotic human coronary vessels by means of in situ hybridization and immunocytochemistry methods. In particular, the expression of mRNAs of ANP, BNP and CNP, measured by RT-PCR, tended to be increased in macroscopically diseased arteries compared to normal vessels, although only the values for BNP expression were significantly different [Casco et al., 2002]. This study suggests that the CNH system is involved in the pathobiology of intimal plaque formation as well as in vascular remodeling in humans. Some studies indicated that there are complex interactions even among CNH themselves. Nazario et al. [Nazario et al., 1995] reported that ANP and BNP can stimulate CNP production through a guanylate cyclase receptor on endothelial cells. As a result, vasodilatory, and anti-mitogenic effects of ANP and BNP in the vasculature could occur in part through CNP production and subsequent action if these interactions occur in vivo. In other words, ANP/BNP and CNP paracrine system should share a synergic action on vascular tissues.

Several studies have demonstrated complex interactions between CNH and the other endothelium-derived vasorelaxant mediators [Ahluwalia and AJ, 2005]. Indeed, evidence from cellular, animal, and human studies suggests that all CNH are able to stimulate NO production by endothelial NO synthase (eNOS); this effect is probably mediated by clearance receptor NPR-C [Houben et al., 2005]. Stimulation of this NPR-C receptor results in decreased cAMP levels by adenyl cyclase inhibition through an inhibitory guanine nucleotide-regulating protein [Houben et al., 2005]. Furthermore, ANP treatment increases renal and cardiac NO synthesis in rats [Los Angeles Costa et al., 2004]. On the other hand, NO, released from endothelial cells, negatively modulates ANP secretion from atrial myocytes, induced by mechanical stretch in perfused rat heart preparation [Ruskoaho H, 1997]. Furthermore, ANP expression is markedly upregulated in eNOS $-/-$ mice, and exogenous ANP restores ventricular relaxation in wild-type mice treated with NOS inhibitors [Gyurko et al., 2000b]. These data suggest that the CNH and NO systems are linked by a negative feedback

mechanism. Finally, CNH (and especially CNP) mimic many of the anti-atherogenic actions of PGI₂ and NO [Ahluwalia and AJ, 2005]. This gives rise to the possibility that CNP might compensate for the loss of these mediators in cardiovascular pathologies to restore the vasodilator capacity of the endothelium, in addition to its anti-adhesive and anti-aggregatory influences.

CNH also strongly interact with the effectors of counter-regulatory systems at the vascular tissue level [McGrath and de Bold AJ, 2005] [Clerico and Emdin, 2004] [Clerico, 2002a] [de Lemos et al., 2003b] [Ruskoaho, 2003a] [Ventura et al., 2002] [Vatta et al., 1996] [Vatta et al., 1997] [Fermepin et al., 2000] [Rocca et al., 2001] [Tsukagoshi et al., 2001] [Kierner et al., 2002b] [Weber et al., 2003] [Kapoun et al., 2004] [Vollmar, 2005]. In particular, interactions between CNH and ET-1 also appear to be important physiologically; indeed, the vascular effects of CNH are directly opposite to those of ET-1 [Ahluwalia and AJ, 2005] [Han and Hasin, 2003]; in particular, ET-1-induced vasoconstriction and myocyte hypertrophy is inhibited by CNH. While CNP has little natriuretic and diuretic action compared to ANP or BNP, it is capable of modulating the vascular effects of the local RAAS by opposing potent vasoconstriction to angiotensin II [Han and Hasin, 2003]. CNP not only functionally antagonizes ET-1 and angiotensin II, but it also directly modulates ET-1 [Kohno et al., 1992] and angiotensin II [Davidson et al., 1996a] synthesis. On the other hand, ET-1 induces an increase in the number of endothelial cells that secrete CNP [Evans et al., 2002]. Therefore, the parallel production and activity of vasodilator CNP and vasoconstrictors such as ET-1 and angiotensin II allows for tight local regulation of these vasoactive peptides and thus blood flow [Ahluwalia and AJ, 2005] [Han and Hasin, 2003] [Evans et al., 2002].

Furthermore, the inter-relationships between the CNH system and pro-inflammatory cytokines suggest that cardiac hormones play an important role in mechanisms responsible for cardiac and vascular adaptation, maladaptation and remodeling in response to various physiolog-

ical and pathological stimuli [Tenhunen et al., 2005] [Walther et al., 2003] [Glembotski et al., 1993] [Mukoyama et al., 1991b]. The identification of CNP as an EDHF, combined with its expression in endothelial cells, indicates that CNP is suited to modulate the activity of circulating cells, particularly leukocytes and platelets. Moreover, inflammatory stimuli such as IL-1b, TNF and lipopolysaccharide [Suga et al., 1993a] stimulate the release of CNP from isolated endothelial cells. As a result, modulation of the biological activity of CNP is likely to have a profound influence on the development of an inflammatory response. Certainly, an anti-atherogenic activity of CNP fits with the cytoprotective, anti-inflammatory actions of NO and PGI₂, the other major endothelium-derived vasorelaxants [Ahluwalia and AJ, 2005] [Han and Hasin, 2003] [Houben et al., 2005] [Scotland et al., 2005] [Suga et al., 1993a].

From a clinical point of view, it is important to note that exogenous application of CNP in situations where endothelial NO production is compromised might be therapeutic in disorders that are associated with endothelial dysfunction. For example, overexpression of CNP by adenoviral-gene delivery in veins dramatically reduces the luminal narrowing (neointimal hyperplasia) that develops when it is grafted to the carotid artery, thereby retaining patency of the graft [Ohno et al., 2002]. CNH, including CNP, also suppress the production of pro-inflammatory cyclooxygenase 2 metabolites in isolated cells [Vollmar and AK, 2001] [Kierner et al., 2002a]. Other studies demonstrated a direct effect of CNP on immune-cell recruitment in vivo [Ahluwalia and AJ, 2005] [Scotland et al., 2005]. Therefore, like NO, endothelial CNP (like ANP and BNP) exerts a protective anti-inflammatory effect [Vollmar et al., 1995] [Vollmar and Schultz, 1995] [Vollmar and AK, 2001] [Ahluwalia and AJ, 2005] [Scotland et al., 2005] [Kierner et al., 2005]. This inhibitory effect of CNH on leukocytes indicates that these peptides modulate the expression of adhesion molecules on either the endothelium or leukocytes.

Several data support the thesis that CNH (and especially CNP) are

important, endogenous, anti-atherogenic mediators. CNP is a potent inhibitor of vascular smooth muscle migration and proliferation that is stimulated by oxidized LDL [Kohno et al., 1992]. CNP also inhibits the proliferation of vascular smooth muscle [Furuya et al., 1991], and enhances endothelial cell regeneration in vitro and in vivo [Ohno et al., 2002]. The observation that CNP alters leukocyte-endothelial interactions indicates that it might also affect platelet function. In accordance with this, thrombus formation is suppressed significantly in the presence of CNP [Suga et al., 1993a], which indicates that inhibition of coagulation might contribute to the vasoprotective properties of this peptide. Observations that CNP blocks platelet aggregation, induced by thrombin, confirm that endothelium-derived CNP also exerts an anti-thrombotic effect [Ahluwalia and AJ, 2005].

All the above-mentioned studies demonstrate that CNH (and especially CNP) exert a protective effect on endothelial function by decreasing shear stress, modulating coagulation and fibrinolysis pathways, and inhibiting platelet activation (Fig. 3.15). They can also inhibit the vascular remodeling process as well as coronary restenosis post-angioplasty [Ma et al., 2004] [Chen and Burnett] [Morishige et al., 2000] [Takeuchi et al., 2003] [Yasuda et al., 2002] [Qian et al., 2002] [Nakanishi et al., 2005] [Ahluwalia and AJ, 2005] [Ohno et al., 2002] [Kierner et al., 2005]. These vasoprotective actions should be considered as a result of complex inter-relationships between the CNH system and both the synergic (including NO, PGI₂, and other endothelium-derived vasoactive mediators) and the counter-regulatory systems (including endothelins, RAAS, cytokines, and growth factors).

1.8 Summary and Conclusion

Natriuretic peptides (including ANP, BNP, CNP, DNP and urodilatin) constitute a family of peptide hormones and neurotransmitters, sharing a similar peptide chain, characterized by a cysteine bridge (Fig.

3.1). The physiological relevance of these peptides is well demonstrated by their presence since the first dawning of life, from unicellular to pluricellular organisms, including plants and all animals. Furthermore, their genes have been repeatedly doubled during evolution, starting from an ancestral gene, thus suggesting that these peptides are indispensable for life (Fig. 3.6).

CNH have powerful physiological effects on the cardiovascular system, body fluid, and electrolyte homeostasis. These effects can be explained at least in part by the inhibition of counter-regulatory systems, including the sympathetic nervous system, RAAS, the vaso pressin/antidiuretic hormone system, the endothelin system, cytokines and growth factors. The endocrine action shared by plasma ANP and BNP can be enhanced by natriuretic peptides produced locally in target tissues (paracrine action). Indeed, endothelial cells synthesize CNP, which exerts a paracrine action on vessels. Moreover, renal tubular cells produce urodilatin, another member of the peptide natriuretic family, which shows powerful diuretic and natriuretic properties. Genes for natriuretic peptides (including ANP, BNP and CNP) are also expressed in the central nervous system, where they likely act as neurotransmitters and/or neuromodulators. Finally, co-expression of CNH and their receptors was observed in immunocompetent cells, suggesting that CNH may have immunomodulatory and anti-inflammatory functions in mammals. Furthermore, CNH are expressed in almost all the body tissues as well as their specific receptors, including organs and tissues not discussed in this chapter, such as gut [Bosc et al., 2000], skeletal [Vollmar and AK, 2001] and ocular [Rollin et al., 2004] tissues. In all tissues, CNH could also act as a local mediator or paracrine effector of tissue-specific functions.

These data, taken as a whole, strongly suggest that natriuretic peptides constitute a family sharing endocrine, paracrine and autocrine actions and neurotransmitter and immunomodulator functions. Therefore, it can be hypothesized that the CNH system is closely related to the other regulatory systems (nervous, endocrine and immunological) in

a biological hierarchical network (Fig. 3.16) [Barabasi and ZN, 2004] [?].

1.9 Therapeutics of Natriuretic Peptides

Measurement of serum BNP levels is used in the clinic as a diagnostic indicator for heart failure, and synthetic forms of both ANP and BNP have been approved in some countries for the treatment of heart failure [Gardner, 2003]. The extent of their usefulness, however, has come under question due to their limited renal actions, and trials are underway to determine the most effective use of these peptides. In this section, we will explore the history of both synthetic ANP and BNP as therapeutic agents.

1.9.1 Synthetic ANP (Anaritide and Carperitide)

Studies revealed that the mature form of ANP is a 28-amino-acid peptide and that smaller versions are degradation products that maintain various levels of activity. The most widely studied of these is the 25-amino acid peptide lacking the first three amino-terminal residues. This peptide is referred to as ANF IV and its synthetic form is called anaritide. Since the activities of the 25-amino acid and mature 28-amino acid peptide were similar, many studies were conducted with the smaller peptide. Studies by Cody and colleagues indicated that infusion of anaritide in healthy male volunteers resulted in natriuresis, diuresis, and reduction in systolic blood pressure; however, in seven patients with congestive heart failure, the changes in urine volume and sodium excretion were minimal [Cody et al., 1986b]. Saito and colleagues observed a similar lack of diuresis and natriuresis, when congestive heart failure patients were infused with the mature form of ANP [Saito et al., 1987b].

Meanwhile, others acknowledged the renal hyporesponsiveness to anaritide in congestive heart failure patients, but indicated that the renal parameters did show a statistically significant increase in larger patient samples [Fifer et al., 1990]. In Japan, clinical studies on the effectiveness of mature ANP continued; and in 1995, synthetic full length ANP (carperitide) was approved for the treatment of acute decompensated heart failure. In the United States, clinical use of BNP, not ANP, was explored for the treatment of heart failure due to its larger renal responsiveness, and possibly due to unique patient opportunities.

Investigations were also initiated to study the effectiveness of ANP in the treatment of human renal disease. Specifically, trials were conducted to evaluate the ability of anaritide infusion to reduce the need for dialysis in patients with acute tubular necrosis. The initial study with 53 patients suggested a positive outcome for patients receiving anaritide because they had increased creatinine clearance and a decreased need for dialysis [Rahman et al., 1994]. This led to the formation of a multicenter placebo-controlled clinical trial in 504 patients with acute tubular necrosis. While 24-h infusion of anaritide did not improve the overall survival of the patients without dialysis, it appeared that a subset of patients might have benefited [Allgren et al., 1997]. Thus, a second trial was conducted in patients with oliguric acute renal failure. However, this 222 patient trial indicated no statistically significant benefit of anaritide in dialysis-free survival [Lewis et al., 2000]. Both trials remarked on the severe hypotension that often occurred as a result of the anaritide infusion. In fact, it is this severe hypotension that appears to be limiting the utility of anaritide or nesiritide as a therapy for either heart failure or renal disease. The authors stated in their discussion, it is possible that if this hypotension could have been avoided, anaritide would have been efficacious [Lewis et al., 2000]. Anaritide was also investigated for its ability to prevent radiocontrast-induced nephropathy. However, in a 247 person clinical trial anaritide along with hydration was no more effective at preventing radiocontrast-induced nephropathy than hydration alone [Kurnik et al., 1998].

Finally, in 2004, studies conducted in Sweden compared the ability of the loop diuretic, furosemide, or mature ANP (1-28) to increase GFR, renal blood flow, and reduce renal oxygen consumption in patients with acute renal failure. They concluded that furosemide was a more effective agent [Sward et al., 2005]. Therefore, despite its potent natriuretic and diuretic effects in normal, healthy subjects, clinical studies conducted to date indicate little or no therapeutic benefit of ANP analogs in the successful treatment of renal disease.

1.9.2 Synthetic BNP (Nesiritide)

Given the natriuretic effects of ANP, the related peptide BNP, was assumed to elicit a similar response. McGregor and colleagues demonstrated that administration of porcine BNP resulted in a natriuretic response and an increase in urinary excretion of cGMP [McGregor et al., 1990]. Yoshimura and colleagues reported the same response in healthy volunteers to infusion of human BNP [Yoshimura et al., 1991].

Patients with congestive heart failure also responded to infusion of BNP. The effectiveness of 24-h infusion of nesiritide to patients with congestive heart failure was examined in a multicenter, placebo-controlled trial. The peptide resulted in a reduction of both preload and afterload resulting in an increase in stroke volume and cardiac output [Mills et al., 1999]. The results of a second multicenter trial, called the Vasodilation in the Management of Acute Congestive Heart Failure (VMAC) study, compared the effects of the addition of nitroglycerin or nesiritide versus placebo to standard therapy. The group treated with nesiritide had improved dyspnea after 3 h treatment, while there was no difference in the other groups. The nitroglycerin group reported more adverse effects than the nesiritide group. Additionally, patients receiving nesiritide had less adverse cardiovascular effects at either the 0.015 or 0.03mcg/kg/min infusion rate compared to patients receiving dobutamine as determined by the 246-patient PRECEDENT Trial

[deLissovoy et al., 2003].

With the approval of the first new intravenous compound for the treatment of heart failure in many years, use of nesiritide was immediate. After approval, the number of patients treated with nesiritide was larger than any clinical trial and with the larger sample population came some unpleasant findings. Initially, Wang and colleagues reported in 2004 that nesiritide does not improve renal function in patients with chronic heart failure [Wang et al., 2004b], but more damaging were two meta-analysis studies by Sackner-Bernstein and colleagues indicating that nesiritide worsened renal function and increased the likelihood of death [Sackner-Bernstein et al., 2005a] [Sackner-Bernstein et al., 2005b].

The results of a 75-person study (BNP-CARDS study), however, suggest nesiritide has no detrimental effect on renal function, when cohorts of similar baseline renal function were compared [Witteles et al., 2007]. The number of persons in this study was small, however, so a more definitive conclusion on whether nesiritide impairs renal function will have to wait until the result of more detailed, larger studies are released. Several such studies are currently in progress. One is a clinical trial enlisting at least 1,900 patients throughout Europe and Latin America - the ETNA (Evaluating Treatment with Nesiritide in Acute Decompensated Heart Failure) trial. This trial was scheduled to begin in 2006 to study the efficacy of nesiritide on treatment of acutely decompensated heart failure. Results from the trial are not yet available. The second study involving about 900 patients, called FUSION II, was conducted to determine the safety and efficacy of outpatient administration of nesiritide to patients with heart failure. Preliminary analysis indicates that nesiritide did not induce renal complications or increase patient mortality [Cleland et al., 2007].

Finally, there is the ASCEND HF trial (Acute Study of Clinical Effectiveness of Nesiritide in Decompensated Heart Failure). This trial is scheduled to compare the effects of nesiritide treatment versus placebo

for a minimum of 24 h up to a maximum of 7 days in 7,000 heart failure patients. Meanwhile, other therapeutic applications of nesiritide have also been investigated. Given that nesiritide was often reported to decrease pulmonary capillary wedge pressure, Michaels and colleagues tested its effectiveness in pulmonary hypertension, however, they found no effect for a 30 min infusion [Michaels et al., 2005]. Chen and colleagues have investigated the effectiveness of subcutaneous injections of nesiritide. Their most recent paper on effects in a dog heart failure pacing model suggest that subcutaneous injection of nesiritide reduces both preload and afterload but has no effect on cardiac output [Chen et al., 2006].

1.10 BNP and NT-proBNP

1.10.1 Differences in Physiology

BNP is a hormonally active natriuretic peptide that is mainly released from the cardio-myocytes in the left ventricular wall. In reaction to stretch and tension of the myocardial wall the pro-hormone proBNP splits into BNP and the hormonally inactive remnant N-terminal proBNP (NT-proBNP) by proteolytic cleavage Fig. 4. [Pfister and Schneider, 2004] This process occurs under influence of integrins, structures at the Z-disc of sarcomeres, that measure stretch of these sarcomeres [Liang et al., 2000, Pyle and Solaro, 2004] after which both peptides will be secreted in equimolar amounts into the circulation.

Circulating BNP acts as an antagonist of the renin angiotensine aldosterone system, and protects the body from plasma overload by inducing diuresis, natriuresis, vascular dilatation and inhibition of the sympathetic nervous system. [Sudoh et al., 1988] The half life of BNP is around 20 minutes and the half life of NT-proBNP is around 120 minutes. BNP is known to be cleared from the blood by natriuretic




Figure 4: Secretion of BNP and NtproBNP

peptide clearance receptors, by neuro endopeptidases and by the kidneys. Little is known on the exact clearance mechanism of NT-proBNP, although it has been suggested that the kidneys play a major role in this clearance. [Hall, 2005] Absolute values of BNP are significantly lower than values of NT-proBNP, despite equimolar secretion. The reference ranges for BNP and NT-proBNP vary depending on the assay that is used and the nature of the control population. In general, the suggested normal range for circulating BNP is 0.5-30 pg/ml and for circulating NT-proBNP the suggested normal range is 68-112 pg/ml. [Cowie et al., 2003] These natriuretic peptides may be beneficial in clinical practice since plasma levels of BNP and NT-proBNP are elevated in patients with HF and are related to the severity of the disease. [Mukoyama et al., 1990]

1.10.2 BNP and NT-proBNP in clinical practice

BNP and NT-proBNP plasma levels are promising tools in the daily management of suspected or established HF. Most studies on the use of BNP and NT-proBNP in clinical practice addressed their diagnostic properties. However, an increasing amount of evidence is available on the prognostic value of BNP and NT-proBNP and some studies provided hopeful results for the benefits of NT-proBNP guided medical treatment.

Diagnosis

Recent trials provided strong evidence that BNP and NT-proBNP are powerful diagnostic tools in exclusion and diagnosis of HF. The Breathing Not Properly study showed, by means of receiver operating characteristics analyses, that a BNP value of 100 pg/ml was the optimal value to differentiate patients with dyspnoea caused by HF from patients with dyspnoea due to pulmonary pathology at the emergency department Fig. 5. [Maisel et al., 2002b]. This value of 100 pg/ml also discriminated non-systolic HF (LVEF \leq 45%) from non-HF patients at the emergency department. It has also been suggested that BNP could be used to discriminate systolic from diastolic HF. Although non-systolic HF patients had significantly lower BNP plasma levels than systolic HF patients (LVEF \leq 45%), BNP only had modest added value in differentiating non-systolic from systolic HF. In another study, a BNP value of 100 pg/ml added significant value to the diagnosis of HF on top of clinical judgement. [McCullough et al., 2002]

An international pooled analysis of 1256 patients provided cut off values for NT-proBNP in an emergency department setting. An age independent cut point of 300 pg/ml had a negative predictive value of 98%. Additionally, an optimal strategy to identify acute HF was to use age stratified cut off points of 450, 900 and 1800 pg/ml for ages \leq 50, 50-75, and \geq 75 respectively which yielded 90% sensitivity and 84%

specificity for acute HF Fig. 6. [Januzzi et al., 2006, JL et al., 2005] Furthermore, BNP and NT-proBNP seem useful as diagnostic tools in primary care (where most patients with suspected HF are encountered and where only limited diagnostic tools are available) and as such are recommended in recent guidelines. [Swedberg et al., 2005]

The added value of these natriuretic peptides on top of established diagnostic tools, including symptoms and signs, has not been properly studied, in particular in relevant subgroups, and currently large studies are underway addressing this issue. Discharge diagnoses have been instrumental in providing estimates and time trends in prevalence and incidence of HF. However, previous studies in the Netherlands, Sweden and in the United States showed that respectively 20%, 18% and 33% of the patients that were given the discharge diagnosis 'HF' at close examination did not have HF at all. [Ingelsson et al., 2005, Goff et al., 2000] It is unknown whether the established BNP cut off value of 100 pg/ml can also be used at discharge after admission for HF.

Prognosis

The prognostic value of BNP and NT-proBNP is well established in several groups of patients. An early study on 85 patients with chronic HF revealed that BNP is a strong independent predictor of mortality. [Tsutamoto et al., 1997] Another study confirmed these results in a larger research population of 452 systolic HF patients (LVEF \leq 35%). In this study BNP was found to be a strong independent predictor of sudden death during a follow up period of 3 years. [Berger et al., 2002] Furthermore, NT-proBNP was a predictor of sudden death in this study population. A substudy of the COPENHAGEN trial (n=1011) revealed that NT-proBNP was consistently associated with an increased risk for all-cause mortality and hospitalisation for HF in patients with severe HF (LVEF \leq 25%). [Hartmann et al., 2004] Another study on 142 patients with advanced HF also reported that NT-proBNP was an independent predictor of all cause mortality. [Gardner et al., 2003b]




Figure 5: ROC curves for BNP in the diagnosis of heart failure at the emergency department.




Figure 6: ROC curves for NTproBNP in the diagnosis of heart failure at the emergency department.

Guidance of treatment

BNP and NT-proBNP are influenced by drugs that are prescribed to HF patients like diuretics, [Tsutamoto et al., 2004] beta blockers, [Richards et al., 1999] ACE inhibitors or angiotensine II receptor blockers [Latini et al., 2002b] and therefore these natriuretic peptides could possibly be used to guide medical treatment. A small study by the Australia-New Zealand Heart Failure Group including 69 patients with symptomatic HF provided evidence of the possible benefit of a NT-proBNP guided approach to therapy. Half of the patients received therapy guided by plasma NT-proBNP, therapy in the remaining patients was guided by clinical monitoring at the same frequency, but with the physician blinded to the NT-proBNP result. Clinical monitoring was based on scores assigned to 10 symptoms or signs of HF used in the Framingham criteria for HF. The study found significantly lower mortality, fewer hospitalisations and episodes of decompensated HF in the NT-proBNP-guided therapy group (target 1680 pg/ml). [Troughton et al., 2000] Larger studies are underway that are about to provide firmer evidence as to whether or not BNP and NT-proBNP can be used as a marker in the monitoring of treatment of HF patients. [Buckley et al., 1999]

1.10.3 Variables influencing BNP and NT-proBNP levels: potential limitations?

Although natriuretic peptide levels are of value in the diagnosis and prognosis of HF patients, several clinical conditions other than HF influence BNP and NT-proBNP plasma levels as well. These influences may be a disadvantage for the use of BNP and NT-proBNP in clinical practice of HF since it may lead to biased interpretations of the test results.

Cardiac variables: BNP and NT-proBNP are also elevated in patients

with acute coronary syndrome. After acute myocardial infarction, levels of BNP rise rapidly during the first 24 hours and then tend to stabilize, [deLemos et al., 2001] and in patients with a Q-wave infarction, a peak in NT-proBNP levels was found after 12-48 hours. [Talwar et al., 2000] In patients with unstable angina pectoris, BNP levels were found to be four times higher compared to patients with stable angina pectoris. [Kikuta et al., 1996] Moreover, atrial fibrillation resulted in increased BNP levels in patients without, but not in patients with HF. [Knudsen et al., 2005] Right ventricular failure due to acute pulmonary embolism can also be determined by BNP. [Tulevski et al., 2002] Furthermore, hypertensive patients have higher BNP and NT-proBNP levels compared to non-hypertensive subjects. [Boomsma and van den Meiracker, 2001]

Non-cardiac variables: A few studies in relatively small study populations without HF showed that anaemia causes elevated BNP levels [Tsuji et al., 2004, Willis et al., 2005, Wold et al., 2005] and in a study on a small group of HF patients anaemia was also related to increased NT-proBNP levels. [Wu et al., 2005] However, besides that these studies were limited in sample size, they only investigated one of the two peptides and the effect of anaemia on NT-proBNP was investigated in HF patients only. Furthermore, anaemia is often caused by renal dysfunction, but this co-morbidity has not been investigated in detail in these studies. Since both BNP and NT-proBNP are known to be elevated in case of renal dysfunction, [Luchner et al., 2005] and because renal function and HF are interrelated, a study investigating the effect of anaemia and renal function on both BNP and NT-proBNP in HF patients is needed. An additional variable that is related to both BNP and NT-proBNP is obesity. In several large studies lower natriuretic peptide levels were associated with higher body mass indexes. [Das et al., 2005, Krauser et al., 2005, Mehra et al., 2004b, Wang et al., 2004c] As far as diabetes is concerned, results are conflicting between BNP and NT-proBNP; BNP levels did not differ between patients with or without diabetes, [Wu et al., 2004b] but NT-proBNP levels seem to be higher in diabetic patients compared to non diabetics. [Magnusson

et al., 2004] Furthermore, ascitic cirrhosis, hyperaldosteronism, hypercortisolism, carcinoma, subarachnoid hemorrhage, [Pfister and Schneider, 2004] lung cancer, tuberculosis and pulmonary embolism [Arden et al., 1995] are clinical conditions with reported elevated natriuretic peptide levels.

Patient related variables: Studies showed that both BNP and NT-proBNP levels are influenced by biological variation, with the biological variation of BNP being higher compared to NT-proBNP (up to 44% and up to 35% respectively). [Bruins et al., 2004b, Wu et al., 2003b] Both BNP and NT-proBNP increase with advancing age and are higher in females compared to males in healthy subjects. [Raymond et al., 2003]

Previous research shows that BNP is related to maximal exercise performance. [Kruger et al., 2002] Moreover, the influence of moderate physical activity (75% of the maximum) on BNP levels was investigated in 10 healthy subjects, 10 HF patients with NYHA class I-II and in 10 HF patients with NYHA class III-IV. A significant increase in BNP levels was observed directly after exercise. [McNairy et al., 2002] However, it is not known whether B-type natriuretic peptide levels also reflect sub-maximal functional capacity during daily activities and whether they are related to quality of life.

2 Clinical Considerations and Applications in Cardiac Diseases

2.1 Circulating Levels of Cardiac Natriuretic Hormones

2.1.1 Physiological Considerations and Clinical Interpretation

2.1.2 Influence of Age and Gender

The circulating levels of CNH are regulated or modified by several physiological factors (such as circadian variations, age, gender, exercise, body posture, and water immersion), eating habits (especially sodium intake), clinical conditions (Table 5.1), and drugs (including corticosteroids, sex steroid hormones, thyroid hormones, diuretics, angiotensin-converting enzyme [ACE] inhibitors, and adrenergic agonists and antagonists) [Bold et al., 2001] [Clerico, 2002b] [Ruskoaho, 2003b] [de Lemos et al., 2003a] [Clerico and M, 2004] [Goetze, 2004b].

The wide variations of circulating levels of CNH in adult healthy subjects in relation to aging and gender could have a particular clinical relevance [Clerico et al., 2002] [Redfield et al., 2002] [Vasan et al., 2002] [Wang et al., 2002] (Figs. 5.1 and 5.2, Table 5.2). Indeed, Vasan et al. [Vasan et al., 2002] demonstrated that the diagnostic accuracy of CNH assay for community screening is gender-dependent.

In order to explain these variations, the possible influence of sex steroid hormones on the CNH system, as well as the modification of the cardiovascular system with aging, should be taken into account [Friesinger, 1999] [Bold et al., 1996] [de Bold, 1999] [Maffei et al., 2001]. According

to these mechanisms, the higher CNH values of women during the fertile adult period could be explained by the physiological stimulation of female sex steroid hormones. In particular, the BNP concentration is on average 36% higher in women than in men aged less than 50 years [Clerico et al., 2002] (Figs. 5.1 and 5.2, Table 5.2). The increase in CNH with aging may be due to the decline in myocardial function and other organs (including kidney), typical of senescence [Sayama et al., 1999]. In this case, the CNH assay may be considered as a biochemical marker of increased risk of cardiac morbidity in old age [Ueda et al., 2003]. Moreover, the increase in CNH with aging may be due to a decrease in their clearance rate. Indeed, an age modulation of maximum binding capacity of clearance (C-type) receptors for CNH was reported in platelets of elderly persons [Giannessi et al., 2001].

All CNH derive from pre-pro-hormones (i.e., preproANP and preproBNP), containing a signal peptide sequence at the amino-terminal end. The pro-hormones (i.e., proANP and proBNP) are produced by cleavage of signal peptide, and then are further split into inactive longer NH₂-terminal fragments (i.e., NT-proANP or NT proBNP), and a biologically active shorter COOH-terminal peptide (i.e., ANP or BNP), which are secreted in the blood in equimolar amounts (Figs. 3.11 and 3.12). However, ANP and BNP have a shorter plasma half-life and consequently lower plasma concentration, compared to NT-proANP and NT-proBNP (Table 4.1) [Bold et al., 2001] [Clerico, 2002b] [Ruskoaho, 2003b] [de Lemos et al., 2003a] [Clerico and M, 2004] [Goetze, 2004b] [Clerico et al., 2002] [Clerico et al., 2000b].

Studies on structure-activity relationships have shown the importance for the binding to the specific receptors of the central ring structure of CNH, formed by a disulfide bridge between the two cysteine residues. For this reason, only ANP and BNP, which present the disulfide bridge in the peptide chain, share the typical hormonal activity of CNH, while the NT-proANP and NT-proBNP do not [Bold et al., 2001] [Clerico, 2002b] [Ruskoaho, 2003b] [de Lemos et al., 2003a] [Clerico and M, 2004] [Goetze, 2004b] [Clerico et al., 2002].

Theoretically, setting up an immunoassay for NT-proANP and NT-proBNP should be easier because their plasma concentrations are higher than ANP and BNP [Clerico et al., 2000b]. On the other hand, NT-proANP and NT-proBNP immunoassays may be affected by several analytical problems, mainly concerning the different assay specificities; consequently, very different results are produced by different methods with a large bias [Clerico, 2002b] [Clerico and M, 2004] [Goetze, 2004b] [Clerico et al., 2000b] (Table 4.1). The different analytical performance might affect the diagnostic accuracy of the assays, in discriminating between subjects with or without cardiac disease [Clerico, 2002b] [Clerico and M, 2004] [Goetze, 2004b].

The respective advantages of measuring biologically active peptide hormones (ANP and BNP), or inactive peptides (NT-proANP and NT-proBNP) are summarized in Table. The assay of the inactive propeptides better fits the definition of disease marker than the assay of circulating levels of ANP or BNP, which, on the other hand, may be considered a more reliable index of the activation status of the CNH system. Considering the biochemical and physiological characteristics of the different peptides, it is conceivable that ANP is a better marker of acute overload and/or rapid cardiovascular hemodynamic changes than BNP and, especially, than NT-proANP or NT-proBNP [Clerico, 2002b] [Clerico and M, 2004]. For example, circulating levels of ANP are known to be more affected by body position and decreased to a greater extent by a hemodialysis session in patients with chronic renal failure than those of BNP, while plasma NT-proANP is unchanged [Clerico et al., 2001].

2.1.3 Resistance to the Biological Action of CNH

Patients with chronic HF show increased CNH plasma levels compared to normal subjects (Table 3.1, Fig. 3.13). These findings have been defined the “endocrine paradox” in HF [Goetze, 2004b], i.e., extremely

high circulating levels of hormones with powerful natriuretic activity in patients with congestive HF, who show physical signs of fluid retention and vasoconstriction due to a relatively poor biological activity of the CNH system. A blunted natriuretic response after pharmacological doses of ANP and BNP has been observed in experimental models and in patients with chronic HF, suggesting a resistance to the biological effects of CNH, principally natriuresis [Clerico and G, 1995a] [Charloux et al., 2003] [Clerico et al., 2000a] [Clerico and G, 1995b] [Cody et al., 1986a] [Saito et al., 1987a] [Komeichi et al., 1995] [Zeidel, 2000]. This resistance syndrome was also demonstrated by in vivo turnover studies using radioactive tracers in patients with HF [Iervasi et al., 1995] [Clerico et al., 2005a].

Resistance to the biological action of CNH could, theoretically, have three different causes (Table 5.3). First, circulating CNH could be, at least in part, inactive. Furthermore, a great fraction of CNH could be inactivated by plasma and tissue proteases before they bind to specific receptors. These two conditions account for all possible mechanisms acting at the pre-receptor level. Second, down-regulation of specific receptors could explain a reduced CNH activity. Finally, some mechanisms can act at postreceptor level, counteracting the biological effects of CNH.

A resistance to the biological action of CNH may be theoretically due to an increase in degradation (turnover) of circulating biologically active peptides. CNH are degraded in vivo and in vitro by several types of proteolytic enzymes, including serin-proteases, peptidyl arginine aldehyde proteases, kallikrein like proteases, and neutral endopeptidases (NEP) [Clerico and M, 2004] [Goetze, 2004b] [Clerico et al., 2000b] [Apple et al., 2005] [Panteghini and A, 2004] [Belenky et al., 2004a] [Shimizu et al., 2002] [Shimizu et al., 2003a].

Individual differences in the ability of heart tissue to mature the precursor of CNH peptides, or of peripheral tissues to degrade them, may help to explain why there are some differences in the clinical presenta-

tion among patients with HF with similar clinical severity and ventricular function [Goetze, 2004b]. However, further studies are necessary to confirm this hypothesis. From a clinical point of view, it is important to note that some drugs sharing an inhibitory action on both NEP and ACE (so called vasopeptidase inhibitors) may have some beneficial effects in patients with arterial hypertension and/or HF because the administration of these drugs can potentiate the biological activity of CNH system by increasing the concentration of biologically active peptides [Trindade and JL, 2001] [Sagnella, 2002] [Dawson and AD, 2002] [Floras, 2002].

It is important to note that renal function can affect the biological action of CNH in different ways. CNH are small peptides freely filtered by renal glomerulus; the kidneys are probably responsible for about 50% of metabolic clearance rate of plasma ANP and BNP and in this way renal diseases can affect the circulating levels of CNH. Indeed, a decreased renal function greatly increases the plasma CNH concentration and consequently more peptide hormones are available for other target tissues (such as brain, vascular tissue, adrenal gland and so on) [Clerico and M, 2004]. However, luminal perfusion with ANP has been shown to reduce sodium efflux from the inner medullar collecting duct, suggesting that this hormone has also luminal sites of action. As a consequence, a reduction in the filtration can potentially induce renal hypo-responsiveness to CNH. To date, however, ANP has been detected only on tubular basolateral membranes [Charloux et al., 2003].

Some studies suggest that the resistance to biological effects of CNH in HF may be due, in part, to variations in the relative amount of the three different types of natriuretic peptide-specific receptors. In particular, there could be an upregulation of type C receptors (NPR-C) with a parallel down regulation of type A and B receptors (NPR-A and NPR-B) [Andreassi et al., 2001] [Tsunoda et al., 1988] [Tsutamato et al., 1993] [Mukkaddam-Daher et al., 1996] [Kuhn et al., 2004]. NPR-A and NPR-B mediate all known hormonal actions of CNH, therefore their down-regulation should induce a deactivation of the CNH sys-

tem. The upregulation of NPR-C receptors that strongly contribute to the clearance of biologically active peptides could further increase the resistance to CNH in patients with HF [Andreassi et al., 2001].

These findings are well in accordance with the results of in vivo kinetic studies obtained using radioactive tracers in patients with HF [Iervasi et al., 1995] [Clerico et al., 2005a]. Moreover, a study confirmed that mRNA expression levels of ANP, BNP and the NPR-C receptor were markedly increased in human failing hearts [Mukkaddam-Daher et al., 1996]. Reversal of cardiomyocyte hypertrophy during left ventricular assist device support was accompanied by normalization of ANP, BNP and NPR-C mRNA levels and a significant recovery of responsiveness to ANP [Kuhn et al., 2004]. However, [Fan et al., 2004] found that neither NPR-A nor NPR-B were internalized or degraded in response to natriuretic peptide binding in cultured cells.

Another well-characterized deactivation mechanism is the process by which an activated receptor is turned off, commonly referred to as “desensitization”. Phosphorylation of the intracellular kinase homology domain of NPR-A and NPR-B is required for hormone-dependent activation of the receptor, while dephosphorylation at this site causes desensitization. Deactivation of the CNH system via desensitization of NPR-A and NPR-B can occur in response to various pathophysiological stimuli [Fan et al., 2004] [Bryan and LR, 2002] [Potter and T, 1998] .

A peripheral resistance to the biological effects of CNH may play an important role in other clinical conditions, besides HF. For example, NPR-C is also present on cellular membranes of adipose tissue. It was suggested that the increase in NPR-C receptors observed in obese subjects can in turn increase the peripheral degradation of CNH and consequently blunt the action of the CNH system [Dessi-Fulgheri et al., 1998] [Sarzani et al., 2004]. Indeed, studies have documented decreased circulating levels of CNH (especially BNP) in obese subjects, compared to nonobese subjects matched for age and gender [Dessi-Fulgheri et al.,

1998] [Wang et al., 2004a]. This reduced activity of the CNH system may increase the risk of developing arterial hypertension and other cardiovascular diseases due to the non-contrasted and therefore prevailing effects of the counter regulatory system with sodium-retentive and vasoconstrictive properties [Sarzani et al., 2004] [Mehra et al., 2004a] [Wang et al., 2004a]

However, studies found that the NPR-C receptor could be coupled to a G-protein that inhibits cyclic AMP synthesis. These receptors, which are present in great amount especially on the endothelial cell wall, may mediate some paracrine effects of CNP on vascular tissue [Ahluwalia et al., 2004b] [Drewett et al., 1990] [Anand-Srivastava et al., 1988]. Therefore, further studies will be necessary to elucidate the possible role of NPR-C receptors as modulators of CNH action and/or degradation in peripheral tissues.

A large number of studies demonstrated that the activation of the neuro-hormonal system accelerates the left ventricular functional impairment in patients with HF [Bold et al., 2001] [Clerico, 2002b] [Ruskoaho, 2003b] [de Lemos et al., 2003a] [Clerico and M, 2004] [Goetze, 2004b] [Packer, 1992] [Benedict, 1994] [Emdin et al., 2004]. Drugs that contrast the detrimental effects of the neuro-hormonal system activation play a key role for the current pharmacological treatment of HF. Some of these, such as ACE inhibitors, angiotensin II receptor blockers, -blockers, and spironolactone decrease the circulating levels of CNH [Clerico and M, 2004] [Richards et al., 2002] [Latini et al., 2002a] [Richards and RW, 2004] [Bettencourt, 2004] [Cowie and GF, 2002], “normalize” their kinetics, and increase their biological activity. Furthermore, they enhance the natriuretic effect of ANP or BNP analogs administered to patients. In other words, the treatment with this type of pharmacological agents decreases the systemic resistance to the biological effects of CNH [Clerico and G, 1995a] [Clerico et al., 2000a].

Patients with HF show a progressive and parallel increase in CNH levels and in some neuro-hormones and cytokines. This increase can

be closely related to disease severity, as assessed by functional NYHA class (Table 3.1, Fig. 3.13). Plasma BNP values, normalized by mean values found in healthy subjects, are significantly higher than other normalized neuro-hormone and cytokine values in HF (Fig. 5.3, Table 5.4) [Emdin et al., 2004].

On average, the response of the CNH system to the increasing challenge of disease severity may not be linear (Fig. 5.4). The curve reported in Figure 5.4 suggests that the CNH system responds with a sharp increase in BNP plasma concentration in the early phase of HF (NYHA class I-II patients), followed, with the clinical progression of the disease, by a blunted increase (NYHA class III), and finally by a plateau (NYHA class IV) (Table 3.1, Fig. 5.4). These findings are consistent with the results from *in vivo* kinetic studies, indicating that ANP turnover (i.e., metabolic clearance rate and production rate) and natriuresis are both increased in patients in the early phase of HF (NYHA class I), as compared to patients with congestive HF [Clerico and G, 1995a] [Clerico et al., 2000a]. This suggests that resistance to the biological action of CNH is characteristic only of the congestive stage of HF. During the asymptomatic, early phase of the disease, the CNH system is able to compensate for the overactivity of the counter regulatory system (Fig. 5.4).

2.1.4 Diagnostic Accuracy of CNH Assay in Plasma from Patients with Cardiac Diseases

The CNH system activation is modulated not only by hemodynamic factors, but also by the activity of the counter-regulatory neuro-hormonal system. Consequently, it is likely that very small changes in hemodynamics, not assessable by echocardiographic examination, may produce significant (and measurable) variations in plasma concentrations of CNH. Moreover, studies have indicated that very small changes in some neuro-hormones and cytokines can produce wider variations in

BNP circulating levels [Emdin et al., 2004] (Fig. 5.3).

It is well known that changes in hemodynamic parameters (such as left ventricular ejection fraction, EF) and plasma CNH levels (expressed in a log scale) are closely related in patients with cardiovascular diseases (Fig. 5.5) [Clerico, 2002b] [Ruskoaho, 2003b] [de Lemos et al., 2003a] [Clerico and M, 2004] [Hammerer-Lercher et al., 2001] [Packer, 1992] [Benedict, 1994] [Emdin et al., 2004] [Richards et al., 2002] [Latini et al., 2002a] [Richards and RW, 2004] [Bettencourt, 2004] [Cowie and GF, 2002]. However, correlations between plasma CNH levels and echocardiographically measured parameters, such as left ventricular EF, myocardial mass and chamber volumes, are usually less close in the general population (large community-based sample, including healthy subjects with or without individuals with asymptomatic myocardial dysfunction) [Redfield et al., 2002] [Vasan et al., 2002] [Nakamura et al., 2002] [Kawai et al., 2004], as well shown by data reported in Figure 5.6.

It should be emphasized that the recognition of HF syndrome is not equivalent to the clinical diagnosis of cardiomyopathy or to the assessment of left ventricular dysfunction, these latter terms describing possible structural reasons for the development of HF. Instead, HF is a clinical syndrome that is characterized by specific symptoms (namely dyspnea and fatigue) and signs (namely fluid retention). There is no diagnostic test for HF, because it is largely a clinical diagnosis based on a careful history and physical examination [bib, 2001]. Therefore, there are no objective criteria for the identification and/or clinical stratification of patients with suspected HF; several “reference (gold) standards” [Bossuyt, 2003], based on clinical, laboratory and instrumental examinations, can be used to evaluate the diagnostic accuracy of the CNH assay. For this evaluation, some studies took into consideration only echocardiographic assessment of ventricular dysfunction, while in others either results of echocardiography or all clinical data have been used for the diagnosis of HF [Clerico and M, 2004].

Unfortunately, using echocardiography as the unique reference standard may lead to misinterpretations when evaluating the diagnostic accuracy of the CNH assay. A systematic review of clinical studies in patient populations with a prevalence of HF ranging from 3.8% to 51% to determine the diagnostic accuracy of BNP assays found that the sensitivity (ranging from 90 to 97%) was much less variable (by 4.4-fold) than the specificity (ranging from 53 to 84%) [Clerico and M, 2004].

Furthermore, in a meta-analysis [Doust et al., 2004], diagnostic accuracy of BNP assays greatly varied according not only to the group of patients studied, but also to the reference standard used (left ventricular EF $\leq 40\%$ or $\leq 55\%$, diagnosis of diastolic dysfunction, diagnosis of systolic + diastolic dysfunction, or integrative clinical criteria). These data indicate that the choice of a suitable and accurate reference standard for evaluation of the diagnostic accuracy of the BNP assay in patients with HF may be a problem that is actually underestimated in the literature. For a proper definition of diagnostic accuracy, tested individuals should be grouped into those with and without disease, by means of an independent clinical judgment, considering both the CNH assay and echocardiographic data.

Careful echocardiographic examinations usually show slightly better or even similar diagnostic accuracy than the BNP assay in patients with cardiac diseases [Logeart et al., 2002] [Dokainish et al., 2004b]. On the other hand, the CNH assay may have some advantages compared to echocardiographic examinations alone in specific clinical settings. For example, Williams et al. suggested that in patients with chronic HF, the NT-proBNP assay reflects functional cardiac impairment and decreased exercise capacity (measured by peak exercise oxygen consumption) better than the left ventricular EF [Williams et al., 2004].

Several studies indicate that BNP and NT-proBNP are powerful and independent risk markers of cardiovascular events (especially mortality) not only in patients with HF, but also in those with acute coronary

syndrome, as reported in and systematic reviews [Clerico and M, 2004] [Galvani et al., 2004] [Jernberg et al., 2004b] [Wiviott et al., 2004]. Some studies also suggested that the cardiovascular risk increases progressively to CNH concentration [Jernberg et al., 2004b] [Wiviott et al., 2004] [Kellett, 2004b]; that is, there is no threshold that actually identifies patients with null risk.

Diagnostic sensitivity of BNP/NT-proBNP assays in detecting left ventricular systolic dysfunction could be suboptimal in asymptomatic or low-risk individuals, especially in women [Vasan et al., 2002]. Specificity of BNP assays in patients with HF ranged from 53 to 84% and positive predictive values from 3 to 85% in several studies. These data indicate that CNH assays can produce a relatively large number of false-positive results. Consequently, many individuals, actually without HF (about 15-60% of those with positive CNH tests), might undergo expensive and/or harmful investigations to rule out the disease, or even be inappropriately labeled as cardiac patients [Clerico and M, 2004].

Several false-positive results can be observed in patients with various clinical diagnoses, as reported in Table 5.1. In these patients, increased plasma BNP may predict, even in the presence of a normal echocardiographic examination, an increased risk of mortality or major cardiac events, including pulmonary embolism [Kruger et al., 2004b] [Kucher et al., 2003a] [ten Wolde et al., 2003a] and hypertension [Nagaya et al., 2000b], renal failure [Vesely, 2003b] [McCullough et al., 2004a], septic shock [Castillo et al., 2004a], some chronic inflammatory diseases (such as amyloidosis and sarcoidosis) [Palladini et al., 2003a] [Yasutake et al., 2005], and diabetes mellitus [Bhalla et al., 2004a].

On the other hand, false-negative results could be found in patients on treatment with anti-adrenergic agents, diuretics and/or ACE inhibitors, all drugs that can reduce CNH levels [Clerico and M, 2004]. As shown in Figure 5.7, a large number of patients with only mild HF (NYHA classes I and II) may have values slightly above or even under the 99th percentile of distribution values of BNP concentration

in healthy subjects. In these patients, successful treatment and consequent improvement in cardiac function and exercise capacity, and reduction in filling pressure and cardiac volumes, is usually associated with a marked fall in CNH levels: thus, a larger number of patients could have BNP values within the reference range values [Clerico and M, 2004] [Takeda et al., 2004].

However, at matched echocardiographic alterations, patients in whom BNP levels drop in response to therapy have a reduced rate of major cardiac events or mortality, compared to untreated hypertensive patients, who could have similar echocardiographic abnormalities. This represents the rationale for using the CNH assay for therapy decision making and for monitoring HF patients [Clerico and M, 2004] [Richards et al., 2002] [Latini et al., 2002a] [Richards and RW, 2004] [Bettencourt, 2004] [Cowie and GF, 2002].

In populations with a higher prevalence of cardiac diseases, including only individuals with a clinical suspicion of HF, the diagnostic sensitivity of BNP can improve up to 95%, or even more, as long as appropriate cut-off values are selected [Clerico and M, 2004] [Doust et al., 2004]. In this case, a strategy called “SnNout”, which maximizes test sensitivity, could be used to rule out the presence of HF [Sackett and RB, 2002]. Furthermore, CNH assay also shows high (>95%) negative predictive values [Clerico and M, 2004] [Prontera et al., 2003], which can help to confirm the absence of HF. This is the rationale for choosing the CNH assay as the first step for an algorithm for the diagnosis of HF [Remme and K, 2001] [bib, 2003]. Such a clinical strategy has proved successful in some studies evaluating the cost-effectiveness of using plasma BNP measurements for screening of cardiac dysfunction in the general population [Nielsen et al., 2003] [Sim et al., 2003] [Ng et al., 2003b].

2.1.5 Biological Variation of Plasma BNP: a Problem or a Clinical Resource?

The variability of measured plasma concentrations of many substances is due to three different sources: pre-analytical, analytical and inherent biological variation. The latter is usually described as a random variation around a homeostatic setting point, and defined as the intra-individual or within-subject biological variation [Fraser, 2004].

In order to achieve a correct interpretation of serial test results that are collected for follow-up or tailored treatment of HF patients, several studies [Pagani et al., 2003] [d’Eril et al., 2003] [Wu et al., 2003a] [Wu and A, 2004] [Bruins et al., 2004a] evaluated the biological variation of BNP and its related peptides, in both healthy subjects and cardiac patients. Due to secretory bursts and its rapid turnover (half-life about 15-20 min) [Bold et al., 2001] [Vanderheyden et al., 2004], it is not surprising that intra-individual biological variations of plasma BNP levels were found to be very large, in both healthy subjects and patients with heart failure (ranging from 30 to 50%) [Pagani et al., 2003] [Wu et al., 2003a] [Wu and A, 2004] [Bruins et al., 2004a]. According to this, only a decrease of more than 50% or a more than 2-fold increase in plasma BNP should be assumed to be statistically significant in an individual patient.

In contrast with this assumption, a clinical trial [Takeda et al., 2004] has suggested that a BNP decrease inferior to this calculated reference change could be clinically relevant in patients with heart failure. In this study, only the group of patients treated with the β -blocker agent carvedilol, who respond on average with a decrease of only 38% in plasma BNP, showed a clinical improvement [Takeda et al., 2004].

Furthermore, several studies have demonstrated that cardiovascular risk (mortality or major cardiovascular events) increases continuously and progressively throughout the whole range of BNP concentrations

in patients with cardiovascular diseases [Clerico and M, 2004] [Jernberg et al., 2004b] [Wiviott et al., 2004] [Kellett, 2004b].

In order to explain these conflicting findings, it should be taken into account that BNP secretion is closely regulated by specific pathophysiological mechanisms. Thus, changes in plasma hormone levels cannot be interpreted as random variations around a setting point, but as strictly determined by the activation level of the counter-regulatory system and by changes in hemodynamics. The clinician should look at the intra-individual variation in BNP as a mirror of variation in neuro-endocrine network activity. According to this hypothesis, it was suggested to consider all changes in BNP concentration as potentially clinically relevant, even when narrower than the calculated intra-individual biological variation [Clerico et al., 2005b]. In other words, BNP variations should be interpreted and considered by physicians, as the variability of heart rhythm and blood pressure, by taking into account clinical history and examination, comprehensive of the response to specific treatments, as well as of laboratory and instrumental test findings.

2.2 CNH Assay as Diagnostic and Prognostic Tool in Cardiac Diseases

In this second part of the Chapter 5, we will review the clinical relevance of CNH assay (especially of BNP and NT-proBNP). There has been an explosion of clinical researches and trials concerning the routine use of CNH (especially BNP) assay in the diagnosis and risk stratification of patients with cardiovascular diseases in the first years of the new century . It is clearly impossible to cover all these scientific contributes. In order to reduce the number of references without reducing the efficacy and objectiveness, we have used systematic reviews and meta-analyses, when available. First, we will review the clinical

relevance of CNH assay in the screening and classification of patients with cardiac dysfunction, divided according to the severity of disease and age, or associated to some particular clinical conditions (such as myocardial infarction or treatment with cardiotoxic drugs) . Moreover, we will discuss the diagnostic accuracy and clinical relevance of assay in coronary artery disease , where more BNP and NT-proBNP are increasingly frequently assayed with (almost in part) unexpected results. Finally, the diagnostic accuracy of CNH assay will be compared to that of other clinical tests, such as ECG, echocardiogram and chest radiogram . The last part of the Chapter is dedicated to prognostic relevance of CNH assay in cardiac diseases and general population , to relevance of measurement of CNH in tailoring the therapy , and to its cost-effectiveness in patients with HF.

2.2.1 Use of CNH Assay in the Screening and Classification of Patients with Cardiac Dysfunction

The diagnosis of HF can often be difficult, mainly in primary care settings, where patients may present with non-specific symptoms and signs, such as dyspnea, fatigue, and ankle swelling [bib, 2001] [Remme and K, 2001] [bib, 2003]. In several population-based studies, fewer than 40% of patients with a suspected diagnosis of HF in primary care had this diagnosis confirmed by more specific and accurate clinical investigations, which are often expensive, timeconsuming and demanding for the patient [bib, 2001] [Remme and K, 2001] [bib, 2003] [Remes et al., 1991] [Fox and andWood DA et al, 2001]. As a result, a relatively simple and inexpensive biochemical test (such as the CNH assay) may be very useful to confirm the clinical suspicion of HF in this clinical setting [Clerico and M, 2004] [Apple et al., 2005] [Panteghini and A, 2004].

2.2.2 Diagnostic Accuracy of CNH Assay in Asymptomatic,

Mild Ventricular Systolic Dysfunction

Patients with asymptomatic left ventricular systolic dysfunction are likely to have lower plasma BNP than those with overt HF [Clerico, 2002b] [Ruskoaho, 2003b] [de Lemos et al., 2003a] [Clerico and M, 2004] [Apple et al., 2005] [Panteghini and A, 2004] [Latini et al., 2002a] [Richards and RW, 2004] [Bettencourt, 2004] [Cowie and GF, 2002] [Nakamura et al., 2002] [Doust et al., 2004], as shown in (Fig. 3.13.)

Two large studies [Vasan et al., 2002] [Nakamura et al., 2002] evaluated the diagnostic accuracy of the CNH assay as a screening method in a general population. The first study analyzed the Framingham Heart Study cohort (3,177 individuals) using BNP and NT-proANP in the evaluation of left ventricular hypertrophy and systolic dysfunction in a community population [Vasan et al., 2002].

The presence of the disease was evaluated by using echocardiographic findings (the prevalence of left ventricular systolic dysfunction was 9.3% in the 1,470 men and 2.5% in the 1,707 women tested, respectively). The area under the curve (AUC) of receiver operating characteristic (ROC) analysis for CNH assay for identifying both left ventricular hypertrophy and systolic dysfunction was on average about 0.75, with a good specificity (assumed 95% both for men and women) and negative predictive value (NPV, on average ranging from 92% to 97% in men, and from 91% to 98% in women), but a poor sensitivity (i.e., ranging from 27% to 28% in men, and from 13% to 40% in women) and positive predictive value (PPV, from 22% to 38% in men, and from 5% to 40% in women), using gender-related BNP cut-off values [Vasan et al., 2002].

The aim of the second study was to examine the validity of plasma BNP measurement for detection of various cardiac abnormalities in a

rural Japanese population (1,098 subjects, 693 men and 405 women), with a low prevalence of coronary heart disease and left ventricular systolic dysfunction (i.e., only 37 participants, corresponding to 3.0%, showed an EF \leq 30%). The diagnosis was carried out by two independent cardiologists based on a medical questionnaire, chest radiogram, electrocardiogram (ECG), and echocardiographic report. The optimal threshold for identification of disease was a BNP concentration of 50 ng/l (14.4 pmol/l), with an area under the ROC curve of 0.970, a sensitivity of 89.7%, a specificity of 95.7%, PPV of 44.3%, and NPV of 99.6%, respectively. [Nakamura et al., 2002]

The conclusions of these two studies, though similar in aim, as well as in clinical and experimental protocols, were strongly conflicting. The Japanese study suggested that the BNP assay is a very efficient and cost-effective mass screening technique for identifying patients with various cardiac abnormalities regardless of etiology and degree of left ventricular systolic dysfunction [Nakamura et al., 2002], while the Framingham study suggested only limited usefulness of the CNH assay as a mass screening tool for this clinical condition, especially in women [Vasan et al., 2002].

This discrepancy may be due to the different gold standard used for the diagnosis of heart failure adopted in the two studies, as discussed in a previous paragraph. However, these two studies, taken as a whole, indicate that the CNH assay may have only a limited usefulness as a screening method for HF in a general population, owing to the poor sensitivity and PPV. However, both studies also found good specificity and NPV, thus suggesting that the CNH assay may be used to rule out HF in an asymptomatic individual.

2.2.3 Diagnostic Accuracy of CNH Assay in Patients with Suspected HF

Many studies [Nielsen et al., 2003] [Omeland et al., 1996b] [Hobbs et al., 2002] [Cowie et al., 1997] [Maisel et al., 2002a] [Maisel et al., 2003] [Fisher et al., 2001] [Apple et al., 2003] [Bay et al., 2003] [McLean et al., 2003] [Richards et al., 1998] [and Yoshimura M and et al, 1994] [Nakagawa et al., 1995] [Vasan et al., 1995] [Bonow and JE, 1992] [Bonow and JE, 1992] [Suzuki et al., 2000] [Maisel et al., 2001] [Lubien et al., 2002] [Vanderheyden and and Verstrecken S et al, 2004] [Hayakawa et al., 2001b] [Ono et al., 2001] have suggested that the CNH assay could be useful as a screening method and/or for the differential diagnosis in patients suspected of HF in different clinical settings:

- randomly selected high-risk community populations
- primary-care patients with a new diagnosis of HF
- patients with acute dyspnea in the emergency department
- consecutive unselected hospital inpatients
- patients admitted to the intensive care unit.

The studies concerning the diagnostic accuracy of the CNH assay in patients with HF have also been analyzed by two systematic reviews [Clerico and M, 2004] [Doust et al., 2004]; unfortunately, these studies showed quite heterogeneous data, thus introducing some bias in the statistical analysis. In particular, even some high quality studies were not designed with the primary goal of evaluating the diagnostic accuracy of the CNH assay. Indeed, this aim was considered only at a post-hoc analysis and retrospectively assessed in blood samples collected for different original purposes, some years before the actual evaluation of diagnostic accuracy. This may introduce a significant

bias, although its true clinical relevance is difficult to assess [Clerico and M, 2004].

The second most important cause of heterogeneous data is the different reference (gold) standard used to evaluate the diagnostic accuracy of the CNH assay. In some studies, the patients studied were stratified and grouped according to clinical severity, as described by functional classification (usually NYHA classification). In other studies, only echocardiographic measurements were used as “gold standard” for the accuracy of the CNH assay for the diagnosis of left ventricular dysfunction (and not for the clinical diagnosis of HF) [Clerico and M, 2004] [Doust et al., 2004].

Furthermore, comparison of studies concerning the diagnostic accuracy of CNH assay is also difficult because different populations were enrolled and different immunoassays were used. Indeed, diagnostic accuracy (especially predictive values) is strictly dependent on disease prevalence (pre-test probability), which greatly varies according to the clinical setting considered (i.e., screening for general population, outpatients seen by a general practitioner, or in primary care, emergency department, coronary care unit, and so on). In particular, the prevalence of HF in the populations studied varied from less than 5% (in studies of screening of asymptomatic population) to more than 50% (in studies including patients referred to hospital with suspicion of HF) [Clerico and M, 2004].

Finally, data reported in the literature suggested that diagnostic accuracy may vary significantly in relation with the specific cardiac peptide measured and/or immunoassay used. Unfortunately, some studies do not clearly indicate the type of immunoassay used to measure CNH, while the majority do not report the assay performance (and often even the reference values) evaluated in their own laboratory.[Clerico and M, 2004]

Taking all studies as a whole, a meta-analysis showed that the odds

ratio for diagnostic accuracy of BNP assay in different groups of patients with suspected HF is highly significant. In particular, the pooled diagnostic odds ratio, when clinical criteria were used as gold standard for HF, was 30.9 (95% confidence interval 27.0-35.4), while it fell to 11.9 (8.4-16.1) when a value 40% of left ventricular EF, estimated by echocardiography, was used as reference standard [Doust et al., 2004].

In a metaanalysis, including data of three studies comparing the diagnostic accuracy of BNP assay in patients with diastolic dysfunction, the pooled odds ratio was 28.3 (95% confidence interval 2.66-300.5). However, a bias may affect these studies, as suggested by the significant test of heterogeneity [Doust et al., 2004].

Finally, [Wright et al., 2003] evaluated the effect of NT-proBNP assay on the clinical diagnostic accuracy of HF in primary care by means of a prospective, randomized controlled trial in 305 patients. Each patient was randomized in two groups, one in which the general practitioner had at their disposal the NT-proBNP assay results (NT-proBNP assay group), while the other did not (control group). The diagnostic accuracy improved by 21% in the NT-proBNP assay group and by 8% in the control group ($p = 0.002$). This study indicates that NT-proBNP measurement significantly improves the clinical diagnostic accuracy of HF in general practice [Wright et al., 2003].

2.2.4 Diagnostic Accuracy of CNH Assay in Patients with Acute Myocardial Infarction

Circulating levels of CNH increase after acute myocardial infarction (AMI); the extent of the increase is related to the size of the infarct [Horio et al., 1993] [and Yasue H and Yoshimura M et al, 1993] [Arakawa et al., 1994] [Uusimaa et al., 1999]. Patients with smaller infarcts tend to have a monophasic increase in plasma BNP, peaking at 20 hours after the onset of symptoms; on the other hand, those with larger

infarcts, lower EF, and clinical signs of HF may present a further peak at 5 days after admission [and Yasue H and Yoshimura M et al, 1993]. Other studies are less convincing regarding the ability of the CNH assay to identify patients with significant myocardial dysfunction after AMI [Omeland et al., 1996a] [Panteghini et al., 2003]. These conflicting results could be due to the differences in sample collection time, type of CNH (ANP, BNP, or NT-proBNP) measured, type of assay, and inclusion criteria adopted. However, persisting elevation of CNH levels at 1 or 2 months after AMI usually suggests a high risk of adverse remodeling and subsequent HF [Clerico and M, 2004].

The diagnostic accuracy of the BNP assay in patients with myocardial infarction was evaluated in a meta analysis [Doust et al., 2004] taking into account only two studies [Bettencourt et al., 2000] [Choy et al., 1994a], fitting the inclusion criteria of this analysis; the pooled odds ratio was 9.4 (95% confidence interval 4.5-19.4).

2.2.5 Diagnostic Accuracy of CNH Assay in Elderly People

Heart failure is primarily a disease of old age; chronic HF increases in prevalence with aging from 1% in people aged ≤ 65 years to 5% in those ≥ 65 years of age, and this clinical condition is the first cause of morbidity and mortality in older people [bib, 2001] [Remme and K, 2001] [bib, 2003] [Baruch et al., 2004a]. A study demonstrated that elderly patients present with more advanced HF, as evidenced by their higher morbidity and mortality rate along with greater neurohormonal activation [Baruch et al., 2004a]. According to these findings, elderly people should be considered to be a population with high risk for developing HF and so the BNP/NT-proBNP assay may be useful as a screening test for HF in older age. Indeed, several studies reported that the BNP/NT-proBNP assay could be clinically useful in elderly people suspected to have HF [Ueda et al., 2003] [Baruch et al., 2004a] [Hutcheon et al., 2002] [Hedberg et al., 2004a] [Heidenreich et al., 2004]

[Ray et al., 2004] [Valle et al., 2005].

Two studies compared the diagnostic accuracy of the BNP assay and that of ECG in elderly people screened for HF. A prospective cohort study specifically evaluated the diagnostic accuracy for HF of BNP assay in 299 consecutive patients (mean age 79 years, 65% women) attending day-hospital over a period of 13 months. This study suggested that both BNP assay and ECG were sensitive in detecting left ventricular systolic dysfunction, but lacked specificity (but the combination of the two tests improved diagnostic accuracy) [Hutcheon et al., 2002].

The other study reported that both the ECG and the plasma concentration of BNP were highly efficient in excluding left ventricular systolic dysfunction in 407 75-year-old subjects. However, compared with the BNP, the ECG yields a lower number of false-positive cases. Therefore, this study suggested that in screening for left ventricular systolic dysfunction in elderly people, the BNP assay has a diagnostic value in addition to the ECG, but only in individuals with abnormal ECG [Hedberg et al., 2004a].

Another study indicated that the BNP assay may be particularly useful in elderly patients, especially in differentiating cardiogenic pulmonary edema from respiratory causes of dyspnea [Ray et al., 2004]. Screening of populations with more than 1% prevalence of HF (such as people with age more than 60 years) with BNP followed by echocardiography should provide a health benefit at a cost that is comparable to or less than other accepted health interventions [Heidenreich et al., 2004]. Finally, the NT-proBNP assay was demonstrated to be useful for detecting HF among people living in elderly nursing homes [Valle et al., 2005].

2.2.6 Detection of Drug Cardiotoxicity by means of BNP Assay

Several studies demonstrated that the BNP assay may also be a marker of chemotherapy cardiotoxicity [Suzuki and Yamazaki T et al, 1998] [Nousiainen et al., 1998a] [Nousiainen et al., 1999a] [Okumura and Yoshida T et al, 2000] [Hayakawa et al., 2001c]. In particular, two studies ([Pichon et al., 2005] and [Sandri et al., 2005]) also suggested that BNP/NT-proBNP assay is a predictive marker of cardiac dysfunction in patients affected by aggressive malignancies and treated with high-dose chemotherapy. The acute release of circulating levels of troponin should be only a mirror of the death of myocytes, while the persistent increase in BNP, after several days or weeks from the administration of cardiotoxic drug, should be specifically related to ventricular remodeling and myocardial dysfunction. Further studies are necessary to confirm these data, and to evaluate the respective diagnostic accuracy and clinical relevance of troponin and BNP assays in the follow-up of patients treated with potentially cardiotoxic drugs.

2.2.7 Diagnostic Accuracy of CNH Assay in Coronary Artery Disease

Two very studies suggested that the BNP assay could be a reliable marker of ischemia in patients with coronary artery disease [Foote et al., 2004] [Sabatine et al., 2004]. It could be hypothesized that exercise-induced ischemia results in increased wall stress and triggers release of CNH from myocytes. According to this hypothesis, Foote et al. [Foote et al., 2004] measured NT proBNP and BNP in blood samples from a group of normal volunteers, and two groups of patients, one with and the other without coronary artery disease, before and after maximal exercise. Post-exercise increases in NT-proBNP and BNP were approximately 4-fold higher in the ischemic group than

in the nonischemic group; while in volunteers, the increase was almost identical to that of the non-ischemic patient group. At equal specificity to the ECG (58.8%), the sensitivities of the BNP/NT-proBNP assay in detecting ischemia were 90 and 80%, respectively; in contrast, the sensitivity of the exercise ECG was only 37.5%. In the study by Sabatine et al. [Sabatine et al., 2004], transient myocardial ischemia was associated with an immediate rise in circulating BNP levels, and the magnitude of the rise was proportional to the severity of ischemia. These findings demonstrate an important link between the severity of an acute ischemic insult and the circulating levels of BNP. However, further studies are necessary to evaluate the relevance of the BNP/NT-proBNP assay.

2.3 Comparison between the Diagnostic Accuracy of CNH Assay and that of Other Tests and Clinical Investigations

Signs and symptoms correlate poorly with the presence of HF, for this reason diagnosis relies on clinical judgment based on a combination of history, physical examination and appropriate investigation [Cowie and GF, 2002] [Remme and K, 2001] [bib, 2003] [Cardarelli and TG, 2003].

Cowie et al. [Cowie et al., 1997] reported that ROC curves for BNP (AUC = 0.96), ANP (0.93), and NT-proANP (0.89) were better than that for cardiothoracic ratio on chest radiogram (0.79) in screening for patients likely to have HF and requiring further clinical assessment. Nielsen et al. [Nielsen et al., 2003] found that BNP assay showed a diagnostic accuracy better than ECG in a random sample of 1,257 community subjects. Another study suggested that BNP assay together with the presence of major ECG abnormalities and history reduced by a factor of six (in comparison to consideration of BNP assay in isola-

tion) the number of subjects requiring echocardiography to detect one case of myocardial dysfunction in a large population screening (1,360 patients tested) [Ng et al., 2003a].

Several studies have compared the diagnostic accuracy of BNP assay and ECG in the elderly population. In 75-year-old subjects both the ECG and the plasma concentration of BNP are highly efficient in excluding left ventricular systolic dysfunction, as suggested [Talwar et al., 1999]. In another study, several types of structural heart disease, in particular valvular heart disease, could be identified exclusively by BNP testing, suggesting that BNP measurement can make a significant contribution to screening for CHF precursors when used in combination with ECG in elderly populations (856 subjects enrolled, with age 65 years) [Nakamura et al., 2005]. However, compared with the BNP assay, the ECG yielded a lower number of false positive cases in another study [Hedberg et al., 2004b]. In screening for left ventricular systolic dysfunction, the BNP has a diagnostic value in addition to the ECG, but only in individuals with abnormal ECG [Hedberg et al., 2004b].

NT-proBNP alone was a better predictor of left ventricular dysfunction than any other single or combination of factors, while the ECG had a poor predictive value for left ventricular systolic dysfunction in identifying patients with left ventricular systolic dysfunction in a high-risk population (243 patients, 129 men, median age 73 years, range 20-94) [Talwar et al., 1999].

Another study [Ogawa et al., 2002] compared the diagnostic accuracy of BNP assay, ECG, chest radiography and echocardiography; the results of this study confirmed that sensitivity, specificity and accuracy of BNP assay and echocardiography were significantly better than those of ECG and chest radiography [Ogawa et al., 2002].

A huge number of studies have compared the diagnostic accuracy of BNP assay and echocardiography in patients with acute or chronic

HF [Clerico and M, 2004] [Latini et al., 2002a] [Richards and RW, 2004] [Bettencourt, 2004] [Cowie and GF, 2002] [Doust et al., 2004] [Struthers, 1993]. According to the guidelines for the diagnosis of heart failure proposed by the Task Force on Heart Failure of the European Society of Cardiology [Remme and K, 2001], echocardiography is recommended as the most practical tool to demonstrate cardiac dysfunction. However, a epidemiological study [Thomas et al., 2004] suggested that the diagnosis of HF might be better defined in terms of symptoms, elevated neuro-hormones and impaired cardiac workload. In more than 50% of the clinical studies, the echocardiographic investigation has been considered as the gold standard (more exactly the reference standard method) for the evaluation of diagnostic accuracy of CNH assay in patients with heart failure [Clerico and M, 2004] [Doust et al., 2004]. When we use the echocardiography investigation as reference method, we assume that BNP assay cannot theoretically have a better diagnostic accuracy than it. Indeed, conflicting results were found when different clinical settings (patient's selection) and/or gold standard were used [Clerico and M, 2004] [Doust et al., 2004].

Choy et al. [Choy et al., 1994b] showed that in post-AMI patients plasma BNP is superior to all clinical indices of left ventricular systolic dysfunction (EF \leq 40%), including signs and symptoms and a clinical score (Peel Index). Dokainish et al. [Dokainish et al., 2004a] found that BNP assay and comprehensive Doppler echocardiography have similar diagnostic accuracy for congestive HF, although echo-Doppler trended toward higher specificity than BNP for congestive HF. Moreover, serial BNP measurements during the treatment of acute HF provide incremental prognostic information over clinical presentation and repetitive echocardiographic examination [Gackowski et al., 2004].

Mak et al. suggested that both BNP assay and a complete echocardiographic examination do not have adequate discriminatory power to be used in isolation in the evaluation of left ventricular diastolic function. Therefore, all available information, including systolic function, chamber dimensions and all Doppler variables must be considered in

the analysis of individual patients [Mak et al., 2004].

Steg et al. assessed the diagnostic performance of BNP testing and echocardiographic assessment of left ventricular systolic function, separately and combined, for the identification of congestive HF in 1,586 patients presenting to the emergency department with acute dyspnea. The proportions of patients who were correctly classified were 67% for BNP alone, 55% for EF alone, 82% for the two variables together, and 97.3% when clinical, ECG, and chest radiograph data were added. This study suggested that BNP measurement was superior to twodimensional echocardiographic determination of left ventricular EF in identifying congestive HF, regardless of the threshold value. The two methods combined have marked additive diagnostic value [Steg et al., 2005].

Some studies indicated that BNP/NT-proBNP levels do not accurately predict serial hemodynamic changes and consequently do not obviate the need for pulmonary artery catheterization in patients requiring invasive hemodynamic monitoring [Forfia et al., 2005] [O'Neill et al., 2005] [Parsonage et al., 2005].

In conclusion, BNP assay shows significantly higher predictive characteristics than ECG and chest radiography, and a cost-benefit value significantly greater than that of echocardiography [Nielsen et al., 2003] [Ogawa et al., 2002]. BNP measurement may exclude normal heart with high probability owing to its high degree of sensitivity and NPV when used in screening high-risk populations, therefore reducing the echocardiographic diagnostic burden; this is the rationale for considering the BNP assay in the first step of an algorithm for the differential diagnosis of heart failure [Shimizu et al., 2002] [Cowie and GF, 2002] [Remme and K, 2001] [bib, 2003] [Cardarelli and TG, 2003] [Struthers, 1993].

2.4 Use of CNH Assay as Prognostic Marker in Cardiovascular Diseases

Several well-designed and conducted studies suggested that the CNH assay may be useful as a prognostic marker mainly in two clinical conditions: HF and acute coronary artery syndromes (ACS), as also demonstrated by systematic reviews [Clerico and M, 2004] [Galvani et al., 2004] [Parsonage et al., 2005] [Rodeheffer, 2004].

2.4.1 Prognosis in HF

The prognostic role of CNH assay (especially NT-proANP, BNP and NT-proBNP) in patients with HF is well demonstrated by a huge number of studies [Ueda et al., 2003] [Valle et al., 2005] [Doust et al., 2005] [Benedict et al., 1996] [Wallen et al., 1997] [Tsutomoto et al., 1999] [Stanek et al., 2001] [Anand et al., 2003] [Gardner et al., 2003a] [Baruch et al., 2004b] [Kellett, 2004a] [Kirk et al., 2004] [Berger et al., 2005a] [Hulsmann et al., 2005] [Segawa et al., 2005] [Stanton and andWijeyesundera HC et al, 2005].

In all these studies, CNH concentrations were always found to be independent risk markers for morbidity (increased future major cardiovascular events and/or hospitalization) and/or mortality in patients with acute or chronic HF [Clerico and M, 2004]. In some studies CNH levels were stronger predictors of mortality and/or major cardiovascular events than left ventricular EF, NYHA class, and/or presence of diabetes or hypertension, as well as sex and age in patients with chronic HF [Wallen et al., 1997] [Tsutomoto et al., 1999] [Gardner et al., 2003a] [Kellett, 2004a] [Kirk et al., 2004] [Berger et al., 2005a] [Segawa et al., 2005] [Stanton and andWijeyesundera HC et al, 2005]. A continuous relationship was generally found between BNP levels and mortality [Kellett, 2004a], thus suggesting that it is not possible to observe a

threshold for the risk. On average, a systematic analysis of the most important studies suggested an odds ratio of about 2 for the risk of mortality in patients with BNP values above the cut-off [Clerico and M, 2004].

In general, BNP assay was found to be the most powerful indicator for poor outcome compared to other neuro effectors and hormones, at least in the largest studies [Clerico and M, 2004] [Latini et al., 2004] [Berger et al., 2005b]. However, more data on the clinical relevance of combination of two or more neuro-hormones in risk stratification of HF are necessary.

2.4.2 Prognosis in ACS

Acute coronary artery syndrome encompasses a continuum of cardiac ischemic events, ranging from unstable angina pectoris, with no biochemical evidence of myocardial necrosis, to ST-elevation AMI [bib, 2000] [bib, 2002]. The prognosis of patients with ACS varies widely, and several clinical, electrocardiographic, and biochemical markers have been used to identify high-risk individuals in need of aggressive intervention [bib, 2002].

Several studies reported that CNH assay (in particular BNP and NT-proBNP) provides valuable prognostic information in patients with ACS [Richards et al., 1998] [Omland et al., 1996a] [Omland et al., 2002] [Arakawa et al., 1996] [Darbar et al., 1996] [Cirilley and M, 2001] [McDonagh et al., 2001] [Richards et al., 2001] [Inoue et al., 2002] [Sabatine et al., 2002] [Jernberg et al., 2002]. A meta-analysis confirmed the powerful prognostic value of BNP/NT-proBNP in patients with ACS for death both in the short term (≤ 50 days, mean odds ratio 3.38, CI 95% 2.44-4.68) and long term (≥ 10 months, mean odds ratio 4.31, 3.77-4.94) [Galvani et al., 2004].

Some studies suggest that BNP levels improve a simple risk score in patients with unstable angina and non-ST-elevation myocardial infarction [Wylie et al., 2004] as well as the risk assessment performance obtained with the cTnI and hs-CRP assays in patients with STsegment elevation myocardial infarction [Jernberg et al., 2004a]. These data suggest the utility of a “multimarker strategy” or biomarker profile to characterize patients with ACS [Wiviott et al., 2004].

A study reported that there may be clear differences among the profiles of individual natriuretic peptide levels in the 2 years after AMI. Similar profiles were found with BNP and NT-proBNP plasma concentrations, while those of NT-proANP and CNP differed significantly in a cohort of 236 patients with AMI complicated by clinical, radiologic or echocardiographic evidence of left ventricular dysfunction. Moreover, a single measurement of plasma natriuretic peptide levels during the hospital admission provides limited prognostic information, while NT-proBNP measured by an ELISA method in the 30 days after AMI identifies a cohort of patients at increased risk of adverse outcome thereafter [Squire et al., 2005].

Two studies reported that in patients with clinically stable, angiographically documented coronary artery disease, plasma BNP [Omland et al., 2005] or NT-proBNP [Kragelund et al., 2005] levels are independently related to long term survival in a multivariate model. These studies suggested that BNP and NT-proBNP are markers of long term mortality even in patients with stable coronary disease and add prognostic information above and beyond that provided by conventional cardiovascular risk factors and the degree of left ventricular systolic dysfunction [Omland et al., 2005] [Kragelund et al., 2005].

In order to explain these clinical findings, it is important to note that experimental studies in animals reported that myocardial ischemia or even hypoxia per se could induce the synthesis/secretion of CNH (in particular BNP) from the intact heart in vivo as well as ventricular cells in culture [Hama et al., 1995] [Toth et al., 1994]. Furthermore,

these experimental data are also in accordance with clinical studies indicating that transient myocardial ischemia in patients with stable coronary artery disease is associated with an immediate rise in circulating BNP levels, and that the magnitude of rise is proportional to the severity of ischemia [Foote et al., 2004] [Sabatine et al., 2004] [Marumoto et al., 1995].

2.4.3 Prognostic Relevance of CNH Assay in the General Population

While measurement of plasma BNP concentration has been shown to be useful in the diagnosis of HF (especially acute HF), its role as a screening test for detection of preclinical ventricular remodeling or dysfunction in the general population has not been established [Clerico and M, 2004] [Marumoto et al., 1995] [Dyrbye and MM, 2003]. However, some studies evaluated the prognostic relevance of CNH assay in the general population, especially in high-risk populations, such as elderly people [Ueda et al., 2003] [Benedict et al., 1996] [Arakawa et al., 1996] [Toth et al., 1994] [Goetze et al., 2004a] [Marumoto et al., 1995] [Dyrbye and MM, 2003] [andVasan RS, 2003]. These studies demonstrated that CNH (and especially BNP/NT-proBNP) levels are a sensitive and accurate biochemical marker of an increased risk of cardiac morbidity and total mortality in very elderly persons [Ueda et al., 2003] [Valle et al., 2005] [Wallen et al., 1997] [McDonagh et al., 2001] [andVasan RS, 2003] [Davis et al., 1992]. However, in a community-based study (mean age 56 years) there were no significant trends of increasing incidence of hypertension across BNP categories in men or women, while higher plasma BNP levels were associated with increased risk of BP progression in men but not women [Knight et al., 1999]. The differences between the results of this study [Knight et al., 1999] and those of others [Ueda et al., 2003] [Rodeheffer, 2004] [McDonagh et al., 2001] [andVasan RS, 2003] [Davis et al., 1992] may be due to the different age and gender of subjects studied. Probably, the prognostic power of BNP

assay increases progressively with age and may be gender-dependent.

2.5 CNH Assay in Management of Patients with HF

2.5.1 Clinical Relevance of CNH Assay in Tailoring the Therapy of HF

Medical therapy for HF is based on improving the symptoms and signs of fluid retention (change in dyspnea, edemas, and body weight are the usual markers of response to treatment) and titrating the dosage of drugs (such as diuretics, ACE inhibitors, β -blockers, and spironolactone) following the evidence from randomized clinical trials [Cowie and GF, 2002] [bib, 2001] [Remme and K, 2001] [bib, 2003]. There is no specific surrogate end-point for treating patients with HF that can be used to fine-tune therapy [Cowie and GF, 2002] [bib, 2001] [Remme and K, 2001] [bib, 2003].

Many authors have suggested that the results of CNH assay (especially BNP/NTproBNP assay) may be useful in monitoring and tailoring the medical therapy in HF patients, and in providing a practical objective indicator of optimal therapy [Richards et al., 2002] [Latini et al., 2002a] [Richards and RW, 2004] [Bettencourt, 2004] [Cowie and GF, 2002] [bib, 2001] [Remme and K, 2001] [bib, 2003] [McCullough et al., 2004a] [Takeda et al., 2004] [Cardarelli and TG, 2003] [Murdoch et al., 1999] [Troughton and andYandle TG et al, 2000] [Kawai et al., 2001a] [Maisel, 2001] [Nicholls et al., 2001] [Troughton et al., 2001] [McGeoch et al., 2002] [Mueller and P, 2002] [Hobbs, 2003] [Richards, 2003] [Maisel et al., 2004] [Morimoto et al., 2004] [Wu et al., 2004a] [Mueller et al., 2004], including patients subjected to cardiac transplantation [Kirchhoff et al., 2004].

CNH usually respond to effective treatment with drugs [Clerico and M, 2004] [Richards et al., 2002] [Latini et al., 2002a] [Richards and RW, 2004] [Bettencourt, 2004] [Cowie and GF, 2002] or left ventricular assist device [Kemperman et al., 2004] [Thompson et al., 2005] with a prompt reduction of their circulating levels. Indeed, ACE inhibitors, valsartan, diuretics, nitrates, and endothelin receptor antagonists have been shown to reduce plasma CNH levels in parallel with hemodynamic and clinical improvement [Latini et al., 2002a] [Richards and RW, 2004] [Murdoch et al., 1999] [McGeoch et al., 2002] [Davidson et al., 1996b] [Nishikimi et al., 1996] [Missouris et al., 1998] [Hara et al., 2000a] [Tsutamoto et al., 2001] [Johnson et al., 2002] [Cotter et al., 2004].

More variable effects on plasma CNH levels have been reported after therapy with β -blockers [Takeda et al., 2004] [Bouissou et al., 1989] [Colantonio et al., 1991] [Yoshimura et al., 1994] [Sanderson et al., 1995] [Luchner et al., 1998] [Fujimura et al., 2000] [Hara et al., 2000b] [Hirooka et al., 2001] [Kawai et al., 2001b] [Hara et al., 2002] [andAndreasson K and et al, 2002] [andYu CM andYip G et al, 2003] [Konishi et al., 2003] [van den Meiracker et al., 2003] [andYoshikawa T and et al, 2004]. Some authors suggested that these variable effects may be at least in part attributable to different specificities or to ancillary properties of β -blockers [Latini et al., 2002a]. Ohta et al. [Ohta et al., 2000] reported that both high and low doses of carvedilol have the effect of increasing plasma ANP and BNP levels in rats. This effect was closely related to the upregulation of ANP and BNP mRNA expression, and the down-regulation of NPR-C mRNA expression in the heart [Ohta et al., 2000]. According to these data, we could assume that an acute administration of β -blockers causes an early rise in plasma CNH, while sustained treatment, significantly improving cardiac function and clinical conditions, induces a significant fall in hormone levels [Takeda et al., 2004] [Hara et al., 2000b] [Hara et al., 2002] [andYu CM andYip G et al, 2003] [Konishi et al., 2003].

Despite this huge number of studies suggesting the clinical relevance of monitoring patients with HF by means of CNH assay, two stud-

ies [Murdoch et al., 1999] [Troughton and Yandle TG et al, 2000] have been designed specifically to evaluate the clinical use of CNH assay in monitoring and tailoring the medical therapy in patients with HF. Murdoch et al. [Murdoch et al., 1999] studied 20 patients with mild to moderate chronic HF and receiving stable conventional therapy, who were randomly assigned to titration of ACE inhibitor dosage, according to serial measurement of plasma BNP or to optimal empirical ACE-inhibitor therapy for 8 weeks. Only the BNP-driven approach was associated with a significant reduction in plasma BNP concentration throughout the duration of the study and a significantly greater suppression when compared with empiric therapy after 4 weeks (mean reduction in BNP group -42.1%, 95% CI -58.2, -19.7; mean reduction in empiric therapy group 12.0%, 95% CI -31.8, 13.8; $P = 0.03$) [Murdoch et al., 1999].

Troughton et al. studied 69 patients with impaired systolic function ($EF \leq 40\%$) and symptomatic HF (NYHA class II-IV), who were randomized to receive treatment guided by either plasma NT proBNP concentration or standardized clinical assessment. During the follow-up (minimum 6 months, median 9.5 months), there were fewer total cardiovascular events (death, hospital admission, or HF decompensation) in the NTproBNP-guided group than in the clinical group (19 vs 54, $p = 0.02$). At 6 months, 27% of patients in the NT-proBNP-guided group and 53% in the clinical group had experienced a first cardiovascular event ($p = 0.034$). Changes in left ventricular function, quality of life, renal function, and adverse events were similar in both groups [Troughton and Yandle TG et al, 2000].

Morimoto et al. conducted a cost-effectiveness analysis of regular BNP measurement in the outpatient setting. The target population was symptomatic CHF patients aged 35-85 years, discharged from the hospital. Intervention was BNP measurement once every 3 months (BNP group) or no BNP measurement (clinical group). The baseline analysis during the 9-month period after hospitalization suggested that the introduction of BNP measurement in heart failure management is not

only cost-effective by reducing hospitalization, but also improves the outcome of patients, as assessed by (quality-adjusted life year) analysis [Morimoto et al., 2004].

A randomized clinical trial compared the titration of β -blocker therapy with bisoprolol according to plasma levels of BNP with empiric clinical therapy based on signs and symptoms. Forty-one patients with heart failure were randomized into a clinical trial. The clinical group had β -blocker dosage increased according to standard care, whereas the BNP group had β -blocker dosage up-titrated according to plasma BNP levels plus standard care. The primary outcome was mean β -blocker dose achieved after 3 months. BNP-guided up-titration of β -blocker in ambulatory patients with heart failure did not result significantly different doses of β -blocker at the end of 3 months. However, 45% of patients in the clinical group were on the maximum dose of β -blocker vs. only 19% of patients in the BNP group, although left ventricular ejection fraction was significantly improved in both groups by 7.3%. The slightly lower doses in the BNP group were possibly better tolerated than the doses achieved in the clinical group. Furthermore, a trend toward better quality of life was seen in the BNP group [da Silva et al., 2005].

Accurate diagnosis of clinical deterioration in heart failure can be difficult [bib, 2001] [Remme and K, 2001]. To prevent development into overt congestion, which often requires hospitalization, early diagnosis is of paramount importance. There is a need for objective measurements to aid early diagnosis in a setting where symptoms may be non-specific and abnormalities on physical examination often subtle and minor [Lewin et al., 2005]. Heart failure guidelines recommend the use of weight gain monitoring to help in this task, with the added advantage that patient self-care is encouraged [bib, 2001] [Remme and K, 2001]. It is advised that an increase of 2 Kg over stable body weight over a period of 48-72 h should initiate contact with medical or nursing personnel [bib, 2001] [Remme and K, 2001] [Lewin et al., 2005].

However, Lewin et al. suggested that neither weight gain nor increase in BNP are adequately sensitive as a screen for clinical deterioration in patients with established heart failure (34 clinically stable and other 43 with clinical deterioration of heart failure status). In particular, weight gain is very insensitive, though an increase of 2 Kg demonstrates high specificity for clinical deterioration. On the other hand, BNP change appears to provide better sensitivity than weight change, but it has poor specificity in an established heart failure population [Lewin et al., 2005].

2.5.2 Cost-Effectiveness of CNH Assay in Management of Patients with Acute or Chronic HF

Nielsen et al. [Nielsen et al., 2003] sought to assess the cost-effectiveness of using plasma BNP as a pre-echocardiographic screening test for left ventricular systolic dysfunction in the general population. Screening high-risk subjects by BNP before echocardiography could reduce the cost per detected case of left ventricular systolic dysfunction by 26% for the cost ratio of 1/20 (BNP/echocardiogram). Greater reduced costs (up to 50%) can be predicted for the group of low-risk subjects [Nielsen et al., 2003]. Other studies reported similar results [Sim et al., 2003] [Heidenreich et al., 2004], including a study on old people living in nursing homes [Valle et al., 2005].

Mueller et al. conducted a prospective, randomized, controlled study of 452 patients who presented to the emergency department with acute dyspnea: 225 patients were randomly assigned to a diagnostic strategy involving the measurement of BNP, and 227 were assessed in a standard manner. This study indicated that BNP assay improved the evaluation and treatment of patients with acute dyspnea and thereby reduced the time to discharge and the total cost of treatment in the emergency department [Mueller et al., 2004].

Other studies suggested that BNP testing can also reduce the cost for hospitalization in patients with heart failure [Morimoto et al., 2004] [Wu et al., 2004a], including old people living in nursing homes [Valle et al., 2005]. However, the cost-effectiveness analysis strongly depends on the relative cost of the BNP test compared to that of echocardiograms and/or hospitalization, as well as on the prevalence of HF in the population screened. Unfortunately, these parameters can vary considerably among departments, countries, and health-care systems; so that each laboratory/clinical department should analyze the cost-effectiveness in its own economical framework. Furthermore, cost-effectiveness analysis is also dependent on the sensitivity of BNP assay for detecting HF. Cost-effectiveness will improve if more specific assays are used: this would decrease the number of subjects with false-positive results, and consequently the number of further useless investigations. However, further larger and randomized clinical trials are necessary to confirm the clinical relevance and cost-effectiveness of BNP assay in the follow-up of patients with HF.

In conclusion, several experimental and clinical studies strongly suggest that the circulating CNH should be better considered as an index of activation of the neuro-endocrine system, rather than a marker of myocardial dysfunction [Clerico et al., 2006]. The activation or deactivation of the CNH system is almost always the resultant of one or more physiological or pathological changes. For this reason, the results of CNH assays must be interpreted by taking into account clinical history and examination, as well as all laboratory and instrumental tests. Of course, the great number of pathophysiological mechanisms that can affect the CNH system render sometimes difficult for clinicians to recognize the cause(s) of variations in its activity. CNH assays should be considered as an intellectual spur for the search of pathophysiological mechanisms that can satisfactorily explain the measured variations in hormone concentrations. On the other hand, CNH measurements add a complementary information to other instrumental and investigative tests [Emdin et al., 2005].

References

- Myocardial infarction redefined - a consensus document of the joint european society of cardiology/american college of cardiology committee for the redefinition of myocardial infarction. *j am coll cardiol*. 2000.
- American heart association task force on practice guidelines. evaluation and management of chronic heart failure in the adult. acc/aha practice guidelines. 2001.
- Acc/aha 2002 guideline update for the management of patients with unstable angina and non-st-segment elevation myocardial infarction. a report of the american college of cardiology/american heart association task force on practice guidelines (committee on the management of patients with unstable angina). 2002.
- Clinical guideline 5. chronic heart failure. management of chronic heart failure in adults in primary and secondary care. 2003.
- author = Kiefer and Wiedemann year =, 2004, title = Neuroendocrine pathways of addictive behaviour., journal = Addict Biol, 9, 12, 205.*
- Ahluwalia and Hobbs AJ. Endothelium-derived c-type natriuretic peptide volume = , more than just a hyperpolarized factor. *Trends Pharmacol Sci*, 26:162–167, 2005.
- Ahluwalia, MacAllister, and Hobbs AJ. Vascular actions of natriuretic peptides. cyclic gmp-dependent and -independent mechanisms. *Basic Res Cardiol*, 99:83–89, 2004a.
- Ahluwalia, MacAllister RJ, and Hobbs AJ. Vascular actions of natriuretic peptides. cyclic gmp-dependent and -independent mechanisms. *Basic Res Cardiol*, 99:83–89, 2004b.
- RL Allgren, TC Marbury, SN Rahman, LS Weisberg, AZ Fenves, RA Lafayette, RM Sweet, FC Genter, BR Kurnik, JD Conger, and

- MH Sayegh. Anaritide in acute tubular necrosis; auriculin anaritide acute renal failure study group. *N Engl J Med*, 336:828–834, 1997.
- FA Almeida, M Suzuki, and T Maack. Atrial natriuretic factor increases hematocrit and decreases plasma volume in nephrectomized rats. *Life Sci*, 39:1193–1199, 1986.
- F Ammarguella, I Larouche, and EL Schiffrin. Myocardial fibrosis in doca-salt hypertensive rats: effect of endothelin et(a) receptor antagonism. *Circulation*, 103:319–324, 2001.
- Anand, Fisher LD, and Chiang YT et al. Val-heft investigators. changes in brain natriuretic peptide and norepinephrine over time and mortality and morbidity in the valsartan heart failure trial (val-heft). *Circulation*, 107:1278–1283, 2003.
- Anand-Srivastava and Trachte GJ. Atrial natriuretic factor receptors and signal transduction mechanisms. *Pharmacol Rev*, 45:455–497, 1993.
- Anand-Srivastava, Sairam MR, and Cantin M. Ring-deleted analogs of atrial natriuretic factor inhibit adenylate cyclase/camp system. possible coupling of clearance atrial natriuretic factor receptors to adenylate cyclase/camp signal transduction system. *J Biol Chem*, 265:8566–8572, 1988.
- Persson and Andreasson K and Kahan T et al. Neurohormonal activation in heart failure after acute myocardial infarction treated with beta-receptor antagonists. *Eur J Heart Fail*, 4:73–82, 2002.
- Ando, Ogawa, and Yamaki. Plasma concentrations of atrial, brain, and c-type natriuretic peptides and endothelin-1 in patients with chronic respiratory diseases. *Chest*, 110:462–468, 1996.
- Andreassi, Del Ry S, and Palmieri C et al. Up-regulation of ‘clearance’ receptors in patients with chronic heart failure: a possible explanation for the resistance to biological effects of cardiac natriuretic hormones. *Eur J Heart Fail*, 3:407–414, 2001.

- Freitag and Vasan RS. Screening for left ventricular systolic dysfunction: the use of b-type natriuretic peptide. *Heart Fail Monit*, 4:38–44, 2003.
- Morita and Yasue H and Yoshimura M et al. Increased plasma levels of brain natriuretic peptide in patients with acute myocardial infarction. *Circulation*, 88:82–91, 1993.
- Yoshizawa and Yoshikawa T and Nakamura I et al. Brain natriuretic peptide response is heterogeneous during beta-blocker therapy for congestive heart failure. *J Card Fail*, 10:310–315, 2004.
- Yasue and Yoshimura M and Sumida H et al. Localization and mechanism of secretion of b-type natriuretic peptide in comparison with those of a-type natriuretic peptide in normal subjects and patients with heart failure. *Circulation*, 90:195–203, 1994.
- Fung and Yu CM and Yip G et al. Effect of beta blockade (carvedilol or metoprolol) on activation of the renin-angiotensin-aldosterone system and natriuretic peptides in chronic heart failure. *Am J Cardiol*, 92:406–410, 2003.
- LK Antos and LR Potter. Adenine nucleotides decrease the apparent K_m of endogenous natriuretic peptide receptors for gtp. *Am J Physiol Endocrinol Metab*, 293:E1756–1763, 2007.
- LK Antos, SE Abbey-Hosch, DR Flora, and LR Potter. Atp-independent activation of natriuretic peptide receptors. *J Biol Chem*, 280:26928–26932, 2005.
- Apple, Trinity E, and Steen J et al. Bnp test utilization for chf in community hospital practice. *Clin Chim Acta*, 328:191–193, 2003.
- Apple, Panteghini M, and Ravkilde J et al. Quality specifications for b-type natriuretic peptide assays. *Clin Chem*, 51:486–493, 2005.

- Arakawa, Nakamura M, Aoki H, and Hiramori K. Relationship between plasma level of brain natriuretic peptide and myocardial infarct size. *Cardiology*, 85:334–340, 1994.
- Arakawa, Nakamura M, Aoki H, and Hiramori K. Plasma brain natriuretic peptide concentrations predict survival after acute myocardial infarction. *J Am Coll Cardiol*, 27:1656–1661, 1996.
- Arden, Viars, and Weiss. *Genomics*, 26:385–389, 1995.
- Barabasi and Oltvai ZN. Network biologyvolume = , understanding the cell’s functional organ- ization. *Nat Rev Genet*, 5:101–113, 2004.
- RW Barbee, BD Perry, RN Re, JP Murgo, and LJ Field. Hemodynamics in transgenic mice with overexpression of atrial natriuretic factor. *Circ Res*, 74:747–751, 1994.
- CF Bartels, H Bukulmez, P Padayatti, DK Rhee, C van Ravenswaaij-Arts, RM Pauli, S Mundlos, D Chitayat, LY Shih, LI Al-Gazali, S Kant, T Cole, J Morton, V Cormier-Daire, L Faivre, M Lees, J Kirk, GR Mortier, J Leroy, B Zabel, CA Kim, Y Crow, NE Braverman, F van-den Akker, and MLA Warman. Mutations in the transmembrane natriuretic peptide receptor npr-b impair skeletal growth and cause acromesomelic dysplasia, type maroteaux. *Am J Hum Genet*, 75:27–34, 2004.
- Baruch, Glazer RD, and Aknay N et al. Morbidity, mortality, physiologic and functional parameters in elderly and non-elderly patients in the valsartan heart failure trial (valheft). *Am Heart J*, 148:951–957, 2004a.
- Baruch, Glazer RD, and Aknay N et al. Morbidity, mortality, physiologic and functional parameters in elderly and non-elderly patients in the valsartan heart failure trial (valheft). *Am Heart J*, 148:951–957, 2004b.

- Bay, Kirk V, and Parner J et al. Nt-probnp: a new diagnostic screening tool to differentiate between patients with normal and reduced left ventricular systolic function. *Heart*, 89:150–154, 2003.
- Belenky, Smith A, and Zhang B et al. The effect of class-specific protease inhibitors on the stabilization of b-type natriuretic peptide in human plasma. *Clin Chim Acta*, 340:163–172, 2004a.
- Belenky, Smith, and Zhang. The effect of class-specific protease inhibitors on the stabilization of b-type natriuretic peptide in human plasma. *Clin Chim Acta*, 340:163–172, 2004b.
- Benedict. Neurohormonal aspects of congestive heart failure. (*Review*) *Cardiol Clin*, 12:9–23, 1994.
- Benedict, Shelton B, and Johnstone DE et al. Prognostic significance of plasma norepinephrine in patients with asymptomatic left ventricular dysfunction. solvd investigators. *Circulation*, 94:690–697, 1996.
- BD Bennett, GL Bennett, RV Vitangcol, JR Jewett, J Burnier, W Henzel, and DG Lowe. Extracellular domain-igg fusion proteins for three human natriuretic peptide receptors; hormone pharmacology and application to solid phase screening of synthetic peptide antisera. *J Biol Chem*, 266:23060–23067, 1991.
- Bentzen, Pedersen, and Pedersen HB. Abnormal rhythmic oscillations of atrial natriuretic peptide and brain natriuretic peptide in heart failure. *Clin Sci*, 104:303–312, 2003.
- Berger, Huelsmann M, and Strecker K et al. Neurohormonal risk stratification for sudden death and death owing to progressive heart failure in chronic heart failure. *Eur J Clin Invest*, 35:24–31, 2005a.
- Berger, Huelsmann M, and Strecker K et al. Neurohormonal risk stratification for sudden death and death owing to progressive heart failure in chronic heart failure. *Eur J Clin Invest*, 35:24–31, 2005b.

- R Berger, M Huelsman, K Strecker, A Bojic, P Moser, B Stanek, and R Pacher. B-type natriuretic peptide predicts sudden death in patients with chronic heart failure. *Circulation*, 105:2392–7, 2002.
- Bernstein, Midtbo, and Urdal. Serum n-terminal pro-atrial natriuretic factor 1-98 before and during thyroxine replacement therapy in severe hypothyroidism. *Thyroid*, 7:415–419, 1997.
- Bettencourt. Nt-probnp and bnp: biomarkers for heart failure management. (Review) *Eur J Heart Fail*, 6:359–363, 2004.
- Bettencourt, Ferreira A, and Pardal-Oliveira N et al. Clinical significance of brain natriuretic peptide in patients with postmyocardial infarction. *Clin Cardiol*, 23:921–927, 2000.
- Bhalla, Chiang A, and Epshteyn VA et al. Prognostic role of b-type natriuretic peptide levels in patients with type 2 diabetes mellitus. *J Am Coll Cardiol*, 44:1047–1052, 2004a.
- Bhalla, Chiang, and Epshteyn VA. *title = Prognostic role of B-type natriuretic peptide levels in patients with type, 2 diabetes mellitus.*, 44:1047–1052, 2004b.
- GE Bilder, TL Schofield, and EH Blaine. Release of atrial natriuretic factor; effects of repetitive stretch and temperature. *Am J Physiol*, 251:F817–F821, 1986.
- De Bold, Bruneau BG, and Kuroski de Bold ML. Mechanical and neuroendocrine regulation of the endocrine heart. (Review) *Cardiovasc Res*, 31:7–18, 1996.
- De Bold, Ma KKY, and Zhang Y et al. The physiological and pathophysiological modulation of the endocrine function of the heart. (Review) *Can J Physiol Pharmacol*, 79:705–714, 2001.
- Bonow and Udelson JE. Left ventricular diastolic dysfunction as a cause of congestive heart failure. *Ann Intern Med*, 117:502–510, 1992.

- F Boomsma and AH van den Meiracker. Plasma a- and b-type natriuretic peptides: physiology, methodology and clinical use. *Cardio-vasc Res*, 51:442–9, 2001.
- Booz. Putting the brakes on cardiac hypertrophyvolume = , exploiting the no-cgmp count- er-regulatory system. *Hypertension*, 45:341–346, 2005.
- Gonzalez Bosc, Majowicz, and Vidal NA. Effects of atrial natriuretic peptide in the gut. *Peptides*, 21:875–887, 2000.
- Bossuyt. Study design and quality of evidence. in: Price cp, christenson rh (eds) evidence-based laboratory medicine. from principles to outcome. *AACC Press, Washington DC, pp 75-92*, 2003.
- Bouissou, Galen FX, and Richalet JP et al. Effects of propranolol and pindolol on plasma anp levels in humans at rest and during exercise. *Am J Physiol*, 257:R259–264, 1989.
- J Brown, Q Chen, and G Hong. An autocrine system for c-type natriuretic peptide within rat carotid neointima during arterial repair. *Am J Physiol*, 272:H2919–H2931, 1997.
- Bruins, Fokkema MR, and Romer JW et al. High intraindividual variation of b-type natriuretic peptide (bnp) and amino-terminal probnp in patients with stable chronic heart failure. *Clin Chem*, 50:2052–2058, 2004a.
- S Bruins, MR Fokkema, JW Romer, MJ Dejongste, FP van der Dijs, JM van den Ouweland, and FA Muskiet. High intraindividual variation of b-type natriuretic peptide (bnp) and amino-terminal probnp in patients with stable chronic heart failure. *Clin Chem*, 50:2052–8, 2004b.
- Bryan and Potter LR. The atrial natriuretic peptide receptor (npr-a/gc-a) is dephosphorylated by distinct microcystin-sensitive and magnesium-dependent protein phosphatases. *J Biol Chem*, 277:16041–16047, 2002.

- PM Bryan, D Smirnov, A Smolenski, S Feil, R Feil, F Hofmann, S Lohmann, and LR Potter. A sensitive method for determining the phosphorylation status of natriuretic peptide receptors: cgk- α does not regulate npr-a. *Biochemistry*, 45:1295–1303, 2006.
- MG Buckley, NJ Marcus, and MH Yacoub. Cardiac peptide stability, aprotinin and room temperature: importance for assessing cardiac function in clinical practice. *Clin Sci (Lond)*, 97:689–95, 1999.
- B Burczynska, T Duda, and RK Sharma. Atp signaling site in the arm domain of atrial natriuretic factor receptor guanylate cyclase. *Mol Cell Biochem*, 301:93–107, 2007.
- JC Burnett, PC Kao, DC Hu, DW Heser, D Heublein, JP Granger, TJ Opgenorth, and GS Reeder. Atrial natriuretic peptide elevation in congestive heart failure in the human. *Science*, 231:1145– 1147, 1986.
- Cappuccio, Markandu, and Buckley MG. Raised plasma levels of atrial natriuretic peptides in addison’s disease. *J Endocrinol Invest*, 12: 205–207, 1989.
- Cardarelli and Lumicao TG. B-type natriuretic peptide: a review of its diagnostic, prognostic, and therapeutic monitoring value in heart failure for primary care physicians. (Review) *J Am Board Fam Pract*, 16:327–333, 2003.
- Casco, Veinot, and Kuroski de Bold ML. Natriuretic peptide system gene expression in human coronary arteries. *J Histochem Cytochem*, 50:799–809, 2002.
- Castillo, Zagler A, Carrillo-Jimenez R, and Hennekens CH. Brain natriuretic peptide: a potential marker for mortality in septic shock. *Int J Infect Dis*, 8:271–274, 2004a.
- Castillo, Zagler, Carrillo-Jimenez, and Hennekens CH. Brain natriuretic peptide volume = , a potential marker for mortality in septic shock. *Int J Infect Dis*, 8:271–274, 2004b.

- MS Chang, DG Lowe, M Lewis, R Hellmiss, E Chen, and DV Goeddel. Differential activation by atrial and brain natriuretic peptides of two different receptor guanylate cyclases. *Nature*, 341:68–72, 1989.
- CJ Charles, TC Prickett, EA Espiner, MT Rademaker, AM Richards, and TG Yandle. Regional sampling and the effects of experimental heart failure in sheep: differential responses in a, b and c-type natriuretic peptides. *Peptides*, 27:62–68, 2006.
- Charloux, Piquard F, and Doutreleau S et al. Mechanisms of renal hyporesponsiveness to anp in heart failure. (Review) *Eur J Clin Invest*, 33:769–778, 2003.
- Chatterjee, Hawksworthy, and McLay JS. Cytokine-stimulated nitric oxide production in the human renal proximal tubule and its modulation by natriuretic peptides volume = , a novel immunomodulatory mechanism? *Exp Nephrol*, 7:438–448, 1999.
- Chen and Burnett.
- HH Chen, JA Schirger, A Cataliotti, and JC Burnett. Intact acute cardiorenal and humoral responsiveness following chronic subcutaneous administration of the cardiac peptide bnp in experimental heart failure. *Eur J Heart Fail*, 8:681–686, 2006.
- Cheung, Leung, and Shiu. HpaII polymorphism in the atrial natriuretic peptide gene and hypertension. *Am J Hypertens*, 12:524–527, 1999.
- H Chikuda, F Kugimiya, K Hoshi, T Ikeda, T Ogasawara, T Shimoaka, H Kawano, S Kamekura, A Tsuchida, N Yokoi, K Nakamura, K Komeda, UI Chung, and H Kawaguchi. Cyclic gmp-dependent protein kinase ii is a molecular switch from proliferation to hypertrophic differentiation of chondrocytes. *Genes Dev*, 18:2418–2429, 2004.
- Chinkers and Wilson EM. Ligand-independent oligomerization of natriuretic peptide receptors. identification of heteromeric receptors

- and a dominant negative mutant. *J Biol Chem*, 267:18589–18597, 1992.
- Choy, Darbar D, and Lang CC et al. Detection of left ventricular dysfunction after acute myocardial infarction: comparison of clinical, echocardiographic, and neurohormonal methods. *Br Heart J*, 72:16–22, 1994a.
- Choy, Darbar D, and Lang CC et al. Detection of left ventricular dysfunction after acute myocardial infarction: comparison of clinical, echocardiographic, and neurohormonal methods. *Br Heart J*, 72:16–22, 1994b.
- TH Chun, H Itoh, Y Ogawa, N Tamura, K Takaya, T Igaki, J Yamashita, K Doi, M Inoue, K Masatsugu, R Korenaga, J Ando, and K Nakao. Shear stress augments expression of c-type natriuretic peptide and adrenomedullin. *Hypertension*, 29:1296–1302, 1997.
- H Chusho, N Tamura, Y Ogawa, A Yasoda, M Suda, T Miyazawa, K Nakamura, K Nakao, T Kurihara, Y Komatsu, H Itoh, K Tanaka, Y Saito, and M Katsuki. Dwarfism and early death in mice lacking c-type natriuretic peptide. *Proc Natl Acad Sci U S A*, 98:4016–4021, 2001.
- AL Clavell, AJ Stingo, CM Wei, DM Heublein, and JC Burnett. C-type natriuretic peptide: a selective cardiovascular peptide. *Am J Physiol*, 264:R290–295, 1993.
- JG Cleland, AP Coletta, and AL Clark. Clinical trials update from the american college of cardiology 2007: Alpha, everest, fusion ii, validd, parr-2, remodel, spice, courage, coach, remadhe, pro-bnp for the evaluation of dyspnoea and this-diet. *Eur J Heart Fail*, 9:740–745, 2007.
- Clerico. Pathophysiological and clinical relevance of circulating levels of cardiac natriuretic hormones volume = , is their assay merely a marker of cardiac disease? *Clin Chem Lab Med*, 40:752–760, 2002a.

- Clerico. Pathophysiological and clinical relevance of circulating levels of cardiac natriuretic hormones: is their assay merely a marker of cardiac disease? (*Opinion Article*) *Clin Chem Lab Med*, 40:752–760, 2002b.
- Clerico. Increasing impact of laboratory medicine in clinical cardiology. *Clin Chem Lab Med*, 41:871–883, 2003.
- Clerico and Emdin. Diagnostic accuracy and prognostic relevance of the measurement of the cardiac natriuretic peptides volume = , a review. *Clin Chem*, 50:33–50, 2004.
- Clerico and Iervasi G. Alterations in metabolic clearance of atrial natriuretic peptides in heart failure: how do they relate to the resistance to atrial natriuretic peptides? (*Review*) *J Card Fail*, 1:323–328, 1995a.
- Clerico and Iervasi G. Alterations in metabolic clearance of atrial natriuretic peptides in heart failure: how do they relate to the resistance to atrial natriuretic peptides? *J Card Fail*, 1:323–328, 1995b.
- Clerico and Iervasi. Alterations in metabolic clearance of atrial natriuretic peptides in heart failure volume = , how do they relate to the resistance to atrial natriuretic peptides? *J Card Fail*, 1:323–328, 1995.
- Clerico and Emdin M. Diagnostic accuracy and prognostic relevance of the measurement of the cardiac natriuretic peptides: a review. (*Review*) *Clin Chem*, 50:33–50, 2004.
- Clerico, Iervasi G, and Pilo A. Turnover studies on cardiac natriuretic peptides: methodological, pathophysiological and therapeutical considerations. (*Review*) *Curr Drug Metab*, 1:85–105, 2000a.
- Clerico, Del Ry S, and Giannessi D. Measurement of natriuretic cardiac hormones (anp, bnp, and related peptides) in clinical practice: the need for a new generation of immunoassay methods. (*Review*) *Clin Chem*, 46:1529–1534, 2000b.

- Clerico, Caprioli R, Del Ry S, and Giannessi D. Clinical relevance of cardiac natriuretic peptides measured by means of competitive and non-competitive immunoassay methods in patients with renal failure on chronic hemodialysis. *J Endocrinol Invest*, 24:24–30, 2001.
- Clerico, Del Ry S, and Maffei S et al. Circulating levels of cardiac natriuretic hormones in healthy adult subjects: effects of aging and sex. *Clin Chem Lab Med*, 40:371–377, 2002.
- Clerico, Iervasi G, and Pilo A. Turnover studies on cardiac natriuretic peptides: methodological, pathophysiological and therapeutical considerations. *Curr Drug Metab*, 1:85–105, 2005a.
- Clerico, Zucchelli GC, Pilo A, and Emdin M. Clinical relevance of biological variation of b-type natriuretic peptide. (*Letter*) *Clin Chem*, 51:925–926, 2005b.
- Clerico, Recchia FA, Passino C, and Emdin M. Cardiac endocrine function is an essential component of the homeostatic regulation network: physiological and clinical implications. *Am J Physiol Heart Circ Physiol*, 290:H17–29, 2006.
- Cody, Atlas SA, and Laragh JH et al. Atrial natriuretic factor in normal subjects and heart failure patients: plasma levels and renal, hormonal and hemodynamic responses to peptide infusion. *J Clin Invest*, 78:1362–1374, 1986a.
- RJ Cody, SA Atlas, JH Laragh, SH Kubo, AB Covit, KS Ryman, A Shaknovich, K Pondolfino, M Clark, MJ Camargo, RM Scarborough, and JA Lewicki. Atrial natriuretic factor in normal subjects and heart failure patients; plasma levels and renal, hormonal, and hemodynamic responses to peptide infusion. *J Clin Invest*, 78:1362–1374, 1986b.
- Colantonio, Casale R, and Desiati P et al. Short-term effects of atenolol and nifedipine on atrial natriuretic peptide, plasma renin activity,

- and plasma aldosterone in patients with essential hypertension. *J Clin Pharmacol*, 31:238–242, 1991.
- Cotter, Kaluski E, and Stangl K et al. The hemodynamic and neuro-hormonal effects of low doses of tezosentan (an endothelin a/b receptor antagonist) in patients with acute heart failure. *Eur J Heart Fail*, 6:601–609, 2004.
- Cowie and Mendez GF. Bnp and congestive heart failure. (*Review*) *Prog Cardiovasc Dis*, 44:293–321, 2002.
- Cowie, Struthers AD, and Wood DA et al. Value of natriuretic peptides in assessment of patients with possible new heart failure in primary care. *Lancet*, 350:1349–1353, 1997.
- MR Cowie, P Jourdain, A Maisel, U Dahlstrom, F Follath, R Isnard, A Luchner, T McDonagh, J Mair, M Nieminen, and G Francis. Clinical applications of b-type natriuretic peptide (bnp) testing. *Eur Heart J*, 24:1710–8, 2003.
- Crilley and Farrer M. Left ventricular remodelling and brain natriuretic peptide after first myocardial infarction. *Heart*, 86:638–642, 2001.
- MG Currie, DM Geller, BR Cole, NR Siegel, KF Fok, SP Adams, SR Eubanks, GR Galluppi, and P Needleman. Purification and sequence analysis of bioactive atrial peptides (atriopeptins). *Science*, 223:67–69, 1984.
- Beck da Silva, de Bold A, and Fraser M et al. Bnp-guided therapy not better than expert’s clinical assessment for beta-blocker titration in patients with heart failure. *Congest Heart Fail*, 11:248–253, 2005.
- Darbar, Davidson NC, and Gillespie N et al. Diagnostic value of b-type natriuretic peptide concentrations in patients with acute myocardial infarction. *Am J Cardiol*, 78:284–287, 1996.

- SR Das, MH Drazner, DL Dries, GL Vega, HG Stanek, SM Abdullah, RM Canham, AK Chung, D Leonard, Jr Fh Wians, and JA deLemos. Impact of body mass and body composition on circulating levels of natriuretic peptides: results from the dallas heart study. *Circulation*, 112:2163–8, 2005.
- Davidson, Barr, and Struthers AD. C-type natriuretic peptide. an endogenous inhibitor of vascular angiotensin-converting enzyme activity. *Circulation*, 93:155–159, 1996a.
- Davidson, Coutie WJ, Webb DJ, and Struthers AD. Hormonal and renal differences between low dose and high dose angiotensin converting enzyme inhibitor treatment in patients with chronic heart failure. *Heart*, 75:576–581, 1996b.
- Davis, Fish LC, and Elahi D et al. Atrial natriuretic peptide levels in the prediction of congestive heart failure risk in frail elderly. *JAMA*, 267:2625–2629, 1992.
- Dawson and Struthers AD. Vasopeptidase inhibitors in heart failure. (Review) *J Renin Angiotensin Aldosterone Syst*, 3:156–159, 2002.
- AJ de Bold, HB Borenstein, AH Veress, and H Sonnenberg. A rapid and potent natriuretic response to intravenous injection of atrial myocardial extract in rats. *Life Sciences*, 28(1):89–94, 1981.
- Kuroski de Bold. Estrogen, natriuretic peptides and the renin-angiotensin system. (Review) *Cardiovasc Res*, 41:524–531, 1999.
- de Lemos, McGuire DK, and Drazner MH. B-type natriuretic peptide in cardiovascular disease. (Review) *Lancet*, 362:316–322, 2003a.
- de Lemos, McGuire, and Drazner MH. B-type natriuretic peptide in cardiovascular disease. *Lancet*, 362:316–322, 2003b.
- S Del-Ry, C Passino, M Maltinti, M Emdin, and D Giannessi. C-type natriuretic peptide plasma levels increase in patients with chronic

- heart failure as a function of clinical severity. *Eur J Heart Fail*, 7: 1145–1148, 2005.
- JA deLemos, DA Morrow, JH Bentley, T Omland, MS Sabatine, CH McCabe, C Hall, CP Cannon, and E Braunwald. The prognostic value of b-type natriuretic peptide in patients with acute coronary syndromes. *N Engl J Med*, 345:1014–21, 2001.
- G deLissovoy, DM Stier, G Ciesla, M Munger, and AJ Burger. Economic implications of nesiritide versus dobutamine in the treatment of patients with acutely decompensated congestive heart failure. *Am J Cardiol*, 92:631–633, 2003.
- Dessi-Fulgheri, Sarzani R, and Rappelli A. The natriuretic peptide system in obesity-related hypertension: new pathophysiological aspects. *J Nephrol*, 11:296–299, 1998.
- DM Dickey, DR Flora, PM Bryan, X Xu, Y Chen, and LR Potter. Differential regulation of membrane guanylyl cyclases in congestive heart failure: natriuretic peptide receptor (npr)-b, not npr-a, is the predominant natriuretic peptide receptor in the failing heart. *Endocrinology*, 148:3518– 3522, 2007.
- Dokainish, Zoghbi WA, and Lakkis NM et al. Comparative accuracy of b-type natriuretic peptide and tissue doppler echocardiography in the diagnosis of congestive heart failure. *Am J Cardiol*, 93:1130–1151, 2004a.
- Dokainish, Zoghbi WA, and Lakkis NM et al. Comparative accuracy of b-type natriuretic peptide and tissue doppler echocardiography in the diagnosis of congestive heart failure. *Am J Cardiol*, 93:1130–1135, 2004b.
- Doust, Glasziou PP, Pietrzak E, and Dobson AJ. A systematic review of the diagnostic accuracy of natriuretic peptides for heart failure. (Review) *Arch Intern Med*, 164:1978–1984, 2004.

- Doust, Pietrzak E, Dobson A, and Glasziou P. How well does b-type natriuretic peptide predict death and cardiac events in patients with heart failure: systematic review. (*Review*) *BMJ*, 330:625–633, 2005.
- Drewett, Ziegler RJ, and Trachte GJ. Neuromodulatory effects of atrial natriuretic factor are independent of guanylate cyclase in adrenergic neuronal pheochromocytoma cells. *J Pharmacol Exp Ther*, 255:497–503, 1990.
- Drewett, Ziegler, and Trachte GJ. Neuromodulatory effects of atrial natriuretic peptides correlate with an inhibition of adenylate cyclase but not an activation of guanylate cyclase. *J Pharmacol Exp Ther*, 260:689–696, 1992.
- Duda and Sharma RK. Regulation of guanylate cyclase activity by atrial natriuretic factor and protein kinase c. *Mol Cell Biochem*, 93:179–184, 1990.
- Dyrbye and Redfield MM. The role of brain natriuretic peptide in population screening. *Heart Fail Rev*, 8:349–354, 2003.
- Melzi d’Eril, Tagnochetti T, and Nauti A et al. Biological variation of n-terminal probrain natriuretic peptide in healthy individuals. *Clin Chem*, 49:1554–1555, 2003.
- D’Souza, Davis, and Baxter GF. Autocrine and paracrine actions of natriuretic peptides in the heart. *Pharmacol Ther*, 101:113–129, 2004.
- BS Edwards, RS Zimmerman, TR Schwab, DM Heublein, and JC Burnett. Atrial stretch, not pressure, is the principal determinant controlling the acute release of atrial natriuretic factor. *Circ Res*, 62: 191–195, 1988.
- Emdin, Passino C, and Prontera C et al. Cardiac natriuretic hormones, neuro-hormones, thyroid hormones and cytokines in normal subjects and patients with heart failure. *Clin Chem Lab Med*, 42:627–636, 2004.

- Emdin, Clerico A, and Clemenza F et al. Recommendations for the clinical use of cardiac natriuretic peptides. *Ital Heart J*, 6:430–446, 2005.
- Evans, Youssef, and Yandle TG. Effects of endothelin-1 on release of adrenomedullin and c-type natriuretic peptide from individual human vascular endothelial cells. *J Endocrinol*, 175:225–232, 2002.
- Fan, Bryan PM, and Antos LK et al. Downregulation does not mediate natriuretic peptide-dependent desensitization of npr-a or npr-b: guanylyl cyclase-linked natriuretic peptide receptors do not internalize. *Mol Pharmacol (printed online Sept 30, 2004)*, 2004.
- Fan, Bryan, and Antos LK. Down-regulation does not mediate natriuretic peptide-dependent desensitization of natriuretic peptide receptor (npr)-a or npr-b. guanylyl cyclase-linked natriuretic peptide receptors do not internalize. *Mol Pharmacol*, 67:1–10, 2005a.
- D Fan, PM Bryan, LK Antos, RJ Potthast, and LR Potter. Down-regulation does not mediate natriuretic peptide-dependent desensitization of natriuretic peptide receptor (npr)-a or npr-b: Guanylyl cyclase-linked natriuretic peptide receptors do not internalize. *Mol Pharmacol*, 67:174–183, 2005b.
- Fermepin, Vatta, and Presas. B-type and c-type natriuretic peptides modify norepinephrine uptake in discrete encephalic nuclei of the rat. *Cell Mol Neurobiol*, 20:763–771, 2000.
- MA Fifer, CR Molina, AC Quiroz, TD Giles, HC Herrmann, IR Descheerder, DL Clement, S Kubo, RJ Cody, and JN Cohn. Hemodynamic and renal effects of atrial natriuretic peptide in congestive heart failure. *Am J Cardiol*, 65:211–216, 1990.
- Fisher, Filzmaier K, and Stiegler H et al. Evaluation of a new, rapid bedside test for qualitative determination of b-type natriuretic peptide. *Clin Chem*, 47:591–594, 2001.

- Floras. Vasopeptidase inhibition: a novel approach to cardiovascular therapy. (Review) *Can J Cardiol*, 18:177–182, 2002.
- JP Fluckiger, B Waeber, G Matsueda, B Delaloye, J Nussberger, and HR Brunner. Effect of atriopeptin iii on hematocrit and volemia of nephrectomized rats. *Am J Physiol*, 251:H880–H883, 1986.
- TG Flynn, ML deBold, and AJ deBold. The amino acid sequence of an atrial peptide with potent diuretic and natriuretic properties. *Biochem Biophys Res Commun*, 117:859–865, 1983.
- Foote, Pearlman JD, and Siegel AH and Yeo KT. Detection of exercise-induced ischemia by changes in b-type natriuretic peptide. *J Am Coll Cardiol*, 44:1980–1987, 2004.
- Forfia, Watkins SP, and Rame JE et al. Relationship between b-type natriuretic peptides and pulmonary capillary wedge pressure in the intensive care unit. *J Am Coll Cardiol*, 45:1667–1671, 2005.
- WG Forssmann, R Richter, and M Meyer. The endocrine heart and natriuretic peptides: histochemistry, cell biology, and functional aspects of the renal urodilatin system. *Histochem Cell Biol*, 110:335–357, 1998.
- Foster and Garber DL. Dual role for adenine nucleotide in the regulation of the atrial natriuretic factor receptor subtypes by angiotensin ii. *J Biol Chem*, 273:16311–16318, 1998.
- Fox and Cowie MR and Wood DA et al. Coronary artery disease as the cause of incident heart failure in the population. *Eur Heart J*, 22: 228–236, 2001.
- Fraser. Inherent biological variation and reference values. (Review) *Clin Chem Lab Med*, 42:758–764, 2004.
- Friesinger. Cardiovascular disease in the elderly. (Review) *Cardiol Clin*, 17:35–49, 1999.

- Fujimura, Yasumura Y, and Ishida Y et al. Improvement in left ventricular function in response to carvedilol is accompanied by attenuation of neurohumoral activation in patients with dilated cardiomyopathy. *J Card Fail*, 6:3–10, 2000.
- Fujio, Ohashi, and Nawata. Cardiovascular, renal and endocrine effects of a- human atrial natriuretic peptide in patients with cushing's syndrome and primary aldosteronism. *J Hypertens*, 7:653–659, 1989.
- Fukui, Nawashiro, and Otani. Focal brain edema and natriuretic peptides in patients with subarachnoid hemorrhage. *Acta Neurochir Suppl*, 86:489–491, 2003.
- Fukui, Katoh, and Tsuzuki. Focal brain edema and natriuretic peptides in patients with subarachnoid hemorrhage. *J Clin Neurosci*, 11:507–511, 2004.
- F Fuller, JG Porter, AE Arfsten, J Miller, JW Schilling, RM Scarborough, JA Lewicki, and DB Schenk. Atrial natriuretic peptide clearance receptor; complete sequence and functional expression of cDNA clones. *J Biol Chem*, 263:9395–9401, 1988.
- Furuya, Yoshida, and Hayashi. C-type natriuretic peptide is a growth inhibitor of rat vascular smooth muscle cells. *Biochem Biophys Res Commun*, 177:927–931, 1991.
- Gackowski, Isnard R, and Golmard JL et al. Comparison of echocardiography and plasma b-type natriuretic peptide for monitoring the response to treatment in acute heart failure. *Eur Heart J*, 25:1788–1796, 2004.
- Galvani, Ferrini D, and Ottani T. Natriuretic peptides for risk stratification of patients with acute coronary syndromes. (Review) *Eur J Heart Fail*, 6:327–333, 2004.
- Gardner, Ozalp F, and Murday AJ et al. N-terminal pro-brain natriuretic peptide. a new gold standard in predicting mortality in

- patients with advanced heart failure. *Eur Heart J*, 24:1735–1743, 2003a.
- DG Gardner. Natriuretic peptides: markers or modulators of cardiac hypertrophy? *Trends Endocrinol Metab*, 14:411–416, 2003.
- RS Gardner, F Ozalp, AJ Murday, SD Robb, and TA McDonagh. N-terminal pro-brain natriuretic peptide a new gold standard in predicting mortality in patients with advanced heart failure. *Eur Heart J*, 24:1735–43, 2003b.
- Giannessi, Andreassi MG, and Del Ry S et al. Possibility of age regulation of the natriuretic peptide c-receptor in human platelets. *J Endocrinol Invest*, 24:8–16, 2001.
- Glembotski, Irons, and Krown KA. Myocardial α -thrombin receptor activation induces hypertrophy and increases atrial natriuretic factor gene expression. *J Biol Chem*, 268:20646–20652, 1993.
- Goetze. Biochemistry of pro-b-type natriuretic peptide-derived peptidesvolume = , the endocrine heart revisited. *Clin Chem*, 9:1503–1510, 2004a.
- Goetze. Biochemistry of pro-b-type natriuretic peptide-derived peptides: the endocrine heart revisited. (Review) *Clin Chem*, 9:1503–1510, 2004b.
- Goetze, Gore A, and Moller CH et al. Acute myocardial hypoxia increases bnp gene expression. *FASEB J*, 18:1928–1930, 2004a.
- Goetze, Videbaek, and Boesgaard. Pro-brain natriuretic peptide as marker of cardiovascular or pulmonary causes of dyspnea in patients with terminal parenchymal lung disease. *J Heart Lung Transplant*, 23:80–87, 2004b.
- Goetze, Rehfeld, and Videbaek. B-type natriuretic peptide and its precursor in cardiac venous blood from failing hearts. *Eur J Heart Fail*, 7:69–74, 2005.

- DC Goff, Jr, DK Pandey, FA Chan, C Ortiz, and MZ Nichaman. Congestive heart failure in the united states: is there more than meets the i(cd code)? the corpus christi heart project. *Arch Intern Med*, 160:197–202, 2000.
- MF Goy, PM Oliver, KE Purdy, JW Knowles, JE Fox, PJ Mohler, X Qian, O Smithies, and N Maeda. Evidence for a novel natriuretic peptide receptor that prefers brain natriuretic peptide over atrial natriuretic peptide. *Biochem J*, 358:379–387, 2001.
- C Grepin, L Dagnino, L Robitaille, L Haberstroh, T Antakly, and M Nemer. A hormone-encoding gene identifies a pathway for cardiac but not skeletal muscle gene transcription. *Mol Cell Biol*, 14:3115–3129, 1994.
- Gutkowska, Tremblay, and Antakly. The atrial natriuretic peptide system in rat ovaries. *Endocrinology*, 132:693–700, 1993.
- Gyurko, Kuhlencordt, Fishman, and Huang OL. Modulation of mouse cardiac function in vivo by enos and anp. *Am J Physiol*, 278: H971–981, 2000a.
- Gyurko, Kuhlencordt, Fishman, and Huang OL. Modulation of mouse cardiac function in vivo by enos and anp. *Am J Physiol*, 278, 2000b.
- Haak, Jungmann, and Schoffling. 24-hour variation in atrial natriuretic peptide. *Lancet*, 335:167–168, 1990.
- Haak, Jungmann, Schoffling, and Usadel KH. Evidence for pulsatile secretion of human atrial natriuretic peptide in healthy subjects. *Exp Clin Endocrinol*, 99:108–109, 1992.
- C Hall. Nt-probnp: the mechanism behind the marker. *J Card Fail*, 11:S81–3, 2005.

- Hama, Itoh H, and Shirakami G et al. Rapid ventricular induction of brain natriuretic peptide gene expression in experimental acute myocardial infarction. *Circulation*, 92:1158–1164, 1995.
- Hammerer-Lercher, Neubauer E, and Muller S et al. Head-to-head comparison of n-terminal pro-brain natriuretic peptide, brain natriuretic peptide and n-terminal pro-atrial natriuretic peptide in diagnosing left ventricular dysfunction. *Clin Chim Acta*, 310:193–197, 2001.
- Han and Hasin. Cardiovascular effects of natriuretic peptides and their interrelation with endothelin-1. *Cardiovasc Drugs Ther*, 17:41–52, 2003.
- Hara, Hamada M, and Shigematsu Y et al. Effect of beta-blocker on left ventricular function and natriuretic peptides in patients with chronic heart failure treated with angiotensin converting enzyme inhibitor. *Jpn Circ J*, 64:365–369, 2000a.
- Hara, Hamada M, and Shigematsu Y et al. Effect of beta-blocker on left ventricular function and natriuretic peptides in patients with chronic heart failure treated with angiotensin converting enzyme inhibitor. *Jpn Circ J*, 64:365–369, 2000b.
- Hara, Hamada M, and Ohtsuka T et al. Comparison of treatment effects of bevantolol and metoprolol on cardiac function and natriuretic peptides in patients with dilated cardiomyopathy. *Heart Vessels*, 17:53–56, 2002.
- Hardt and Sadoshima. Negative regulators of cardiac hypertrophy. *Cardiovasc Res*, 63:500–509, 2004.
- F Hartmann, M Packer, AJ Coats, MB Fowler, H Krum, P Mohacsi, JL Rouleau, M Tendera, A Castaigne, SD Anker, I Amann-Zalan, S Hoersch, and HA Katus. Prognostic impact of plasma n-terminal

- pro-brain natriuretic peptide in severe chronic congestive heart failure: a substudy of the carvedilol prospective randomized cumulative survival (copernicus) trial. *Circulation*, 110:1780–6, 2004.
- Hayakawa, Komada, and Hirayama. Plasma levels of natriuretic peptides in relation to doxorubicin-induced cardiotoxicity and cardiac function in children with cancer. *Med Pediatr Oncol*, 37:4–9, 2001a.
- Hayakawa, Komada Y, and Hirayama M et al. Plasma levels of natriuretic peptides in relation to doxorubicin-induced cardiotoxicity and cardiac function in children with cancer. *Med Pediatr Oncol*, 37:4–9, 2001b.
- Hayakawa, Komada Y, and Hirayama M et al. Plasma levels of natriuretic peptides in relation to doxorubicin-induced cardiotoxicity and cardiac function in children with cancer. *Med Pediatr Oncol*, 37:4–9, 2001c.
- Hayashi, Tsutamoto, and Wada. Intravenous atrial natriuretic peptide prevents left ventricular remodeling in patients with first anterior acute myocardial infarction. *J Am Coll Cardiol*, 37:1820–1826, 2001.
- Hedberg, Lonnberg I, and Jonason T et al. Electrocardiogram and b-type natriuretic peptide as screening tools for left ventricular systolic dysfunction in a population-based sample of 75-year-old men and women. *Am Heart J*, 148:524–529, 2004a.
- Hedberg, Lonnberg I, and Jonason T et al. Electrocardiogram and b-type natriuretic peptide as screening tools for left ventricular systolic dysfunction in a population-based sample of 75-year-old men and women. *Am Heart J*, 148:524–529, 2004b.
- Heidenreich, Gubens MA, and Fonarow GC et al. Cost-effectiveness of screening with b-type natriuretic peptide to identify patients with reduced left ventricular ejection fraction. *J Am Coll Cardiol*, 43: 1019–1026, 2004.

- Henriksen, Gotze, and Fuglsang. Increased circulating pro-brain natriuretic peptide (probnp) and brain natriuretic peptide (bnp) in patients with cirrhosis volume = , relation to cardiovascular dysfunction and severity of disease. *Gut*, 52:1511–1517, 2003.
- JP Henry, OH Gauer, and JL Reeves. Evidence of the atrial location of receptors influencing urine flow. *Circ Res*, 4:85–90, 1956.
- Hirooka, Yasumura Y, and Ishida Y et al. Comparative left ventricular functional and neurohumoral effects of chronic treatment with carvedilol versus metoprolol in patients with dilated cardiomyopathy. *Jpn Circ J*, 65:931–936, 2001.
- Hobbs. Using bnp to diagnose, manage, and treat heart failure. (*Review*) *Cleve Clin J Med*, 70:333–336, 2003.
- Hobbs, Davis RC, and Roalfe AK et al. Reliability of n-terminal pro-brain natriuretic peptide assay in diagnosis of heart failure: cohort study in representative and high risk community populations. *Br Med J*, 324:1498–1500, 2002.
- R Holtwick, M Gotthardt, B Skryabin, M Steinmetz, R Potthast, B Zetsche, RE Hammer, J Herz, and M Kuhn. Smooth muscle-selective deletion of guanylyl cyclase-a prevents the acute but not chronic effects of anp on blood pressure. *Proc Natl Acad Sci U S A*, 99:7142–7147, 2002.
- R Holtwick, M Van-Eickels, BV Skryabin, HA Baba, A Bubikat, F Begrow, MD Schneider, DL Garbers, and M Kuhn. Pressure-independent cardiac hypertrophy in mice with cardiomyocyte-restricted inactivation of the atrial natriuretic peptide receptor guanylyl cyclase-a. *J Clin Invest*, 111:1399–1407, 2003.
- Horio, Shimada K, and Kohno M et al. Serial changes in atrial and brain natriuretic peptides in patients with acute myocardial infarction treated with early coronary angioplasty. *Am Heart J*, 126:293–299, 1993.

- Houben, van der Zander, and de Leeuw PW. Vascular and renal actions of brain natriuretic peptide in man volume = , physiology and pharmacology. *Fundam Clin Pharmacol*, 19:411–419, 2005.
- Hulsmann, Berger R, and Mortl D et al. Incidence of normal values of natriuretic peptides in patients with chronic heart failure and impact on survival: a direct comparison of n-terminal atrial natriuretic peptide, n-terminal brain natriuretic peptide and brain natriuretic peptide. *Eur J Heart Fail*, 7:552–556, 2005.
- Hunt, Richards, and Nicholls MG. Immunoreactive amino-terminal pro-brain natriuretic peptide (nt-probnp) volume = , a new marker of cardiac impairment. *Clin Endocrinol*, 47:287–296, 1997.
- Huo, Abe, and Risono KS. Ligand binding-dependent limited proteolysis of the atrial natriuretic peptide receptor juxtamembrane hinge structure essential for transmembrane signal transduction. *Biochemistry*, 38:16941–16951, 1999.
- Hutcheon, Gillespie ND, Struthers AD, and McMurdo ME. B-type natriuretic peptide in the diagnosis of elderly day hospital patients. *Age Ageing*, 31:295–301, 2002.
- Iervasi, Clerico A, and Berti S et al. Altered tissue degradation and distribution of atrial natriuretic peptide in patients with idiopathic dilated cardiomyopathy and its relationship with clinical severity of the disease and sodium handling. *Circulation*, 91:2018–2027, 1995.
- Iervasi, Clerico, and Pilo. Evidence that anp tissue extraction is not changed by large increases of its plasma levels induced by pacing in humans. *J Clin Endocrinol Metab*, 82:884–888, 1997.
- Iervasi, Clerico, and Pilo. Atrial natriuretic peptide is not degraded from lungs in humans. *J Clin Endocrinol Metab*, 83:2898–2906, 1998.
- Imura, Nakao, and Itoh. The natriuretic peptide system in the brain-volume = , implications in the central control of cardiovascular and neuroendocrine functions. *Front Neuroendocrinol*, 13:217–249, 1992.

- E Ingelsson, J Arnlov, J Sundstrom, and L Lind. The validity of a diagnosis of heart failure in a hospital discharge register. *Eur J Heart Fail*, 7:787–91, 2005.
- Inoue, Sakuma M, and Yaguchi I et al. Early recanalization and plasma brain natriuretic peptide as an indicator of left ventricular function after acute myocardial infarction. *Am Heart J*, 143:790–796, 2002.
- Ivanova, Gregoraszczuk, and Augustowska. Localization of atrial natriuretic peptide in pig granulosa cells isolated from ovarian follicles of various size. *Reprod Biol*, 3:173–181, 2003.
- Y Iwanaga, I Nishi, S Furuichi, T Noguchi, K Sase, and Y Kihara. B-type natriuretic peptide strongly reflects diastolic wall stress in patients with chronic heart failure: comparison between systolic and diastolic heart failure. *Journal of American College of Cardiology*, 47:742–8, 2006.
- JD Jamieson and GE Palade. Specific granules in atrial muscle cells. *J Cell Biol*, 23:151–172, 1964.
- JL Januzzi, KR van, J Lainchbury, A Bayes-Genis, J Ordonez-Llanos, M Santalo-Bel, YM Pinto, and M Richards. Nt-probnp testing for diagnosis and short-term prognosis in acute destabilized heart failure: an international pooled analysis of 1256 patients: the international collaborative of nt-probnp study. *Eur Heart J*, 27:330–7, 2006.
- J Jaubert, F Jaubert, N Martin, LL Washburn, BK Lee, EM Eicher, and JL Guenet. Three new allelic mouse mutations that cause skeletal overgrowth involve the natriuretic peptide receptor c gene (npr3). *Proc Natl Acad Sci U S A*, 96:10278–10283, 1999.
- Jensen, Carstens, Ivarsen, and Pedersen EB. A new, fast and reliable radioimmunoassay of brain natriuretic peptide in human plasma. reference values in healthy subjects and in patients with different diseases. *Scand J Clin Lab Invest*, 57:529–540, 1997.

- Jernberg, Stridsberg M, and Lindahl B. 2002.
- Jernberg, James S, and Lindahl B et al. Nt-probnp in unstable coronary artery disease experiences from the fast, gusto iv and frisc ii trials. *Eur J Heart Fail*, 6:319–325, 2004a.
- Jernberg, James S, and Lindahl B et al. Natriuretic peptides in unstable coronary artery disease. (Review) *Eur Heart J*, 25:1486–1493, 2004b.
- Jr Januzzi JL, CA Camargo, S Anwaruddin, AL Baggish, AA Chen, DG Krauser, R Tung, R Cameron, JT Nagurney, CU Chae, DM Lloyd-Jones, DF Brown, S Foran-Melanson, PM Sluss, E Lee-Lewandrowski, and KB Lewandrowski. The n-terminal pro-bnp investigation of dyspnea in the emergency department (pride) study. *Am J Cardiol*, 95:948–54, 2005.
- John, Kregge, and Oliver PM. Genetic decrease in atrial natriuretic peptide and salt-sensitive hypertension. *Science*, 267:679–681, 1995a.
- SW John, JH Kregge, PM Oliver, JR Hagaman, JB Hodgins, SC Pang, TG Flynn, and O Smithies. Genetic decreases in atrial natriuretic peptide and salt-sensitive hypertension [published erratum appears in science 1995 mar 24;267(5205):1753]. *Science*, 267:679–681, 1995b.
- SW John, AT Veress, U Honrath, CK Chong, L Peng, O Smithies, and H Sonnenberg. Blood pressure and fluid-electrolyte balance in mice with reduced or absent anp. *Am J Physiol*, 271:R109–114, 1996.
- Johnson, Damodaran, and Rushin. Ectopic production and processing of atrial natriuretic peptide in a small cell. *Cancer*, 79:35–44, 1997.
- Johnson, Omland T, and Hall C et al. Neurohormonal activation rapidly decreases after intravenous therapy with diuretics and vasodilators for class iv heart failure. *J Am Coll Cardiol*, 39:1623–1629, 2002.

- Joubert, Labrecque, and De Lean. Reduced activity of the npr-a kinase triggers dephosphorylation and homologous desensitization of the receptor. *Biochemistry*, 40:11096–11105, 2001.
- S Joubert, C Jossart, N McNicoll, and A De-Lean. Atrial natriuretic peptide-dependent photolabeling of a regulatory atp-binding site on the natriuretic peptide receptor-a. *FEBS J*, 272:5572–5583, 2005.
- Kalra, Clague, and Bolger AP. Myocardial production of c-type natriuretic peptide in chronic heart failure. *Circulation*, 107:571–573, 2003a.
- PR Kalra, JR Clague, AP Bolger, SD Anker, PA Poole-Wilson, AD Struthers, and AJ Coats. Myocardial production of c-type natriuretic peptide in chronic heart failure. *Circulation*, 107:571–573, 2003b.
- K Kangawa, Y Tawaragi, S Oikawa, A Mizuno, Y Sakuragawa, H Nakazato, A Fukuda, N Minamino, and H Matsuo. Identification of rat gamma atrial natriuretic polypeptide and characterization of the cdna encoding its precursor. *Nature*, 312:152–155, 1984.
- Kapoun, Liang, and O’Young. B-type natriuretic peptide exerts broad functional opposition to transforming growth factor-beta in primary human cardiac fibroblasts volume = , fibrosis, myofibroblasts conversion, proliferation, and inflammation. *Circ Res*, 94:453–461, 2004.
- Kato, Sugiyama, and Morita. Genetic analysis of the atrial natriuretic peptide gene in essential hypertension. *Clin Sci*, 98:251–258, 2000.
- Kawai, Hata K, and Takaoka H et al. Plasma brain natriuretic peptide as a novel therapeutic indicator in idiopathic dilated cardiomyopathy during beta-blocker therapy: a potential of hormone-guided treatment. *Am Heart J*, 141:925–932, 2001a.
- Kawai, Hata K, and Takaoka H et al. Plasma brain natriuretic peptide as a novel therapeutic indicator in idiopathic dilated cardiomy-

- opathy during beta-blocker therapy: a potential of hormone-guided treatment. *Am Heart J*, 141:925–932, 2001b.
- Kawai, Hata K, and Tanaka K et al. Attenuation of biological compensatory action of cardiac natriuretic peptide system with aging. *Am J Cardiol*, 93:719–723, 2004.
- Kellett. Prediction of in-hospital mortality by brain natriuretic peptide levels and other independent variables in acutely ill patients with suspected heart disease. *Can J Cardiol*, 20:686–690, 2004a.
- Kellett. Prediction of in-hospital mortality by brain natriuretic peptide levels and other independent variables in acutely ill patients with suspected heart disease. *Can J Cardiol*, 20:686–690, 2004b.
- Kemperman, van den Berg M, Kirkels H, and de Jonge N. B-type natriuretic peptide (bnp) and n-terminal probnp in patients with end-stage heart failure supported by a left ventricular assist device. *Clin Chem*, 50:1670–1672, 2004.
- Kierner, Lehner, Hartung, and Vollmar AM. Inhibition of cyclooxygenase-2 by natriuretic peptides. *Endocrinology*, 143:846–852, 2002a.
- Kierner, Weber, and Furst. Inhibition of p38 mapk activation via induction of mkp-1 atrial natriuretic peptide reduces tnfa-induced actin polymerization and endothelial permeability. *Circ Res*, 90:874–881, 2002b.
- Kierner, Furst, and Vollmar AM. Vasoprotective actions of the atrial natriuretic peptide. *Curr Med Chem Cardiovasc Hematol Agents*, 3:11–21, 2005.
- K Kikuta, H Yasue, M Yoshimura, E Morita, H Sumida, H Kato, K Kugiyama, H Ogawa, K Okumura, Y Ogawa, and K Nakao. Increased plasma levels of b-type natriuretic peptide in patients with unstable angina. *Am Heart J*, 132:101–7, 1996.

- Kirchhoff, Gradaus R, and Stypmann J et al. Vasoactive peptides during long-term follow-up of patients after cardiac transplantation. *J Heart Lung Transplant*, 23:284–288, 2004.
- Kirk, Bay M, and Parner J et al. N-terminal probnp and mortality in hospitalised patients with heart failure and preserved vs. reduced systolic function: data from the prospective copenhagen hospital heart failure study (chhf). *Eur J Heart Fail*, 6:335–341, 2004.
- B Kisch. Electron microscopy of the atrium of the heart in guinea pig. *Exp Med Surg*, 14:99–112, 1956.
- I Kishimoto, SK Dubois, and DL Garbers. The heart communicates with the kidney exclusively through the guanylyl cyclase-a receptor: acute handling of sodium and water in response to volume expansion. *Proc Natl Acad Sci U S A*, 93:6215–6219, 1996.
- I Kishimoto, K Rossi, and DL Garbers. A genetic model provides evidence that the receptor for atrial natriuretic peptide (guanylyl cyclase-a) inhibits cardiac ventricular myocyte hypertrophy. *Proc Natl Acad Sci U S A*, 98:2703–2706, 2001.
- Knight, Fish LC, and Kiely DK et al. Atrial natriuretic peptide and the development of congestive heart failure in the oldest old: a seven-year prospective study. *J Am Geriatr Soc*, 47:407–411, 1999.
- JW Knowles, G Esposito, L Mao, JR Hagaman, JE Fox, O Smithies, HA Rockman, and N Maeda. Pressure-independent enhancement of cardiac hypertrophy in natriuretic peptide receptor a- deficient mice. *J Clin Invest*, 107:975–984, 2001.
- CW Knudsen, T Omland, P Clopton, A Westheim, AH Wu, P Duc, J McCord, RM Nowak, JE Hollander, AB Storrow, WT Abraham, PA McCullough, and A Maisel. Impact of atrial fibrillation on the diagnostic performance of b-type natriuretic peptide concentration in dyspneic patients: an analysis from the breathing not properly multinational study. *J Am Coll Cardiol*, 46:838–44, 2005.

- Koh, Nussenzveig, and Okolicany. Dynamics of atrial natriuretic factor- guanylate cyclase receptor and receptor-ligand complexes in cultured glomerular mesangial and renomedullary interstitial cells. *J Biol Chem*, 267:11987–11994, 1992a.
- GY Koh, DR Nussenzveig, J Okolicany, DA Price, and T Maack. Dynamics of atrial natriuretic factor- guanylate cyclase receptors and receptor-ligand complexes in cultured glomerular mesangial and renomedullary interstitial cells. *J Biol Chem*, 267:11987–11994, 1992b.
- Kohno, Horio, and Yokokawa. *title = C-type natriuretic peptide inhibits thrombin- and angiotensin II-stimulated endothelin release via cyclic guanosine, 3',5'-monophosphate.*, 19:320–325, 1992.
- Kohno, Horio, and Yasunari. Stimulation of brain natriuretic peptide release from heart by thyroid hormone. *Metabolism*, 42:1059–1064, 1993.
- UM Kokkonen, AR Pösö, S Hyypä, P Huttunen, and J Leppäluoto. Exercise-induced changes in atrial peptides in relation to neuroendocrine responses and fluid balance in the horse. *Veterinary Medicine. A, Physiology, Pathology, Clinical Medicine*, 49(3):144–50, 4 2002.
- KJ Koller, DG Lowe, GL Bennett, N Minamino, K Kangawa, H Matsuo, and DV Goeddel. Selective activation of the b natriuretic peptide receptor by c-type natriuretic peptide (cnp). *Science*, 252:120–123, 1991.
- Komeichi, Moreau R, and Cailmail S et al. Blunted natriuresis and abnormal systemic hemodynamic responses to c-type and brain natriuretic peptides in rats with cirrhosis. *J Hepatol*, 22:319–325, 1995.
- Konishi, Nishio S, and Tsutamoto T et al. Serum carvedilol concentration and its relation to change in plasma brain natriuretic peptide

- level in the treatment of heart failure: a preliminary study. *Int J Clin Pharmacol Ther*, 41:578–586, 2003.
- Kragelund, Gronning B, and Kober L et al. N-terminal pro-b-type natriuretic peptide and long-term mortality in stable coronary heart disease. *N Engl J Med*, 352:666–675, 2005.
- DG Krauser, DM Lloyd-Jones, CU Chae, R Cameron, S Anwaruddin, AL Baggish, A Chen, R Tung, and Jr Januzzi JL. Effect of body mass index on natriuretic peptide levels in patients with acute congestive heart failure: a probnp investigation of dyspnea in the emergency department (pride) substudy. *Am Heart J*, 149:744–50, 2005.
- Kruger, Graf, and Merx MW. Brain natriuretic peptide predicts right heart failure in patients with acute pulmonary embolism. *Am Heart J*, 147:60–65, 2004a.
- Kruger, Graf J, and Merx MW et al. Brain natriuretic peptide predicts right heart failure in patients with acute pulmonary embolism. *Am Heart J*, 147:60–65, 2004b.
- S Kruger, J Graf, D Kunz, T Stickel, P Hanrath, and U Janssens. brain natriuretic peptide levels predict functional capacity in patients with chronic heart failure. *J Am Coll Cardiol*, 40:718–722, 2002.
- Kucher, Printzen G, and Goldhaber SZ. Prognostic role of brain natriuretic peptide in acute pulmonary embolism. *Circulation*, 107: 2545–2547, 2003a.
- Kucher, Printzen, and Goldhaber SZ. Prognostic role of brain natriuretic peptide in acute pulmonary embolism. *Circulation*, 107: 2545–2547, 2003b.
- Kuhn. Molecular physiology of natriuretic peptide signalling. *Basic Res Cardiol*, 99:76–82, 2004.

- Kuhn, Voss M, and Mitko D et al. Left ventricular assist device support reverses altered cardiac expression and function of natriuretic peptides and receptors in end-stage heart failure. *Cardiovasc Res*, 64:308–314, 2004.
- M Kuhn, R Holtwick, HA Baba, JC Perriard, W Schmitz, and E Ehler. Progressive cardiac hypertrophy and dysfunction in atrial natriuretic peptide receptor (gc-a) deficient mice. *Heart*, 87:368–374, 2002.
- BR Kurnik, RL Allgren, FC Genter, RJ Solomon, ER Bates, and LS Weisberg. Prospective study of atrial natriuretic peptide for the prevention of radiocontrast-induced nephropathy. *Am J Kidney Dis*, 31:674–680, 1998.
- Kurokawa, Uede, and Ishiguro. Pathogenesis of hyponatremia following sub- arachnoid hemorrhage due to ruptured cerebral aneurysm. *Surg Neurol*, 46:500–507, 1996.
- Labrecque, McNiccol, Marquis, and De Lean. A disulfide-bridged mutant of natri- uretic peptide receptor-a displays constitutive activity. role of receptor dimerization in sig- nal transduction. *J Biol Chem*, 274:9752–9759, 1999.
- Labrecque, Deschenes, McNiccol, and De Lean. Agonistic induction of a covalent dimer in a mutant of natriuretic peptide receptor-a documents a juxtamembrane interaction that accompanies receptor activation. *J Biol Chem*, 276:8064–8072, 2001.
- RE Lang, H Tholken, D Ganten, FC Luft, H Ruskoaho, and T Unger. Atrial natriuretic factor-a circulating hormone stimulated by volume loading. *Nature*, 314:264–266, 1985.
- TH Langenickel, J Buttgereit, I Pagel-Langenickel, M Lindner, J Monti, K Beuerlein, N Al-Saadi, R Plehm, E Popova, J Tank, R Dietz, R Willenbrock, and M Bader. Cardiac hypertrophy in

- transgenic rats expressing a dominant-negative mutant of the natriuretic peptide receptor b. *Proc Natl Acad Sci U S A*, 103:4735–4740, 2006.
- Langub, Watson RE, and Herman JP. Distribution of natriuretic peptide precursor mRNAs in the rat brain. *J Comp Neurol*, 356:183–199, 1995.
- Lapinski, Stepniakowski, and Januszewicz. Plasma atrial natriuretic peptide concentration in patients with primary aldosteronism. *J Hypertens*, 9 (Suppl. 6):S260–261, 1991.
- LaPointe. Molecular regulation of the brain natriuretic gene. *Peptides*, 26:944–956, 2005.
- Latini, Masson S, De Angelis N, and Anand I. Role of brain natriuretic peptide in the diagnosis and management of heart failure: current concepts. (Review) *J Card Fail*, 8:288–299, 2002a.
- Latini, Masson S, and Anand I et al. The comparative prognostic value of plasma neurohormones at baseline in patients with heart failure enrolled in val-heft. *Eur Heart J*, 25:292–299, 2004.
- R Latini, S Masson, I Anand, D Judd, AP Maggioni, YT Chiang, M Bevilacqua, M Salio, P Cardano, PH Dunselman, NJ Holwerda, G Tognoni, and Cohn JN; Valsartan Heart Failure Trial Investigators. Effects of valsartan on circulating brain natriuretic peptide and norepinephrine in symptomatic chronic heart failure - the valsartan heart failure trial (val-heft). *Circulation*, 106:2454–8, 2002b.
- DC Leitman, JW Andresen, T Kuno, Y Kamisaki, JK Chang, and F Murad. Identification of multiple binding sites for atrial natriuretic factor by affinity cross-linking in cultured endothelial cells. *J Biol Chem*, 261:11650–11655, 1986.
- Leuchte, Holzapfel, and Baumgartner RA. Clinical significance of brain natriuretic peptide in primary pulmonary hypertension. *J Am Coll Cardiol*, 43:764–770, 2004a.

- Leuchte, Neurohr, and Baumgartner. Brain natriuretic peptide and exercise capacity in lung fibrosis and pulmonary hypertension. *Am J Respir Crit Care Med*, 170:360–365, 2004b.
- Lewin, Ledwidge M, and O’loughlin C et al. Clinical deterioration in established heart failure: What is the value of bnp and weight gain in aiding diagnosis? *Eur J Heart Fail*, 7:953–957, 2005.
- J Lewis, MM Salem, GM Chertow, LS Weisberg, F McGrew, TC Marbury, and RL Allgren. Atrial natriuretic factor in oliguric acute renal failure; anaritide acute renal failure study group. *Am J Kidney Dis*, 36:767–774, 2000.
- F Liang, A Atakilit, and DG Gardner. Integrin dependence of brain natriuretic peptide gene promoter activation by mechanical strain. *J Biol Chem*, 275:20355–60, 2000.
- Logeart, Saudubray C, and Beyne P et al. Comparative values of doppler echocardiography and b-type natriuretic peptide assay in the etiologic diagnosis of acute dyspnea. *J Am Col Cardiol*, 40: 1794–1800, 2002.
- Lohmann, Vaandrager, and Smolenski. Distinct and specific functions of cgmp-dependent protein kinases. *Trend Biochem Sci*, 22:307–312, 1997.
- MJ Lopez, SK Wong, I Kishimoto, S Dubois, V Mach, J Friesen, DL Garbers, and A Beuve. Salt-resistant hypertension in mice lacking the guanylyl cyclase-a receptor for atrial natriuretic peptide. *Nature*, 378:65–68, 1995.
- MJ Lopez, DL Garbers, and M Kuhn. The guanylyl cyclase-deficient mouse defines differential pathways of natriuretic peptide signaling. *J Biol Chem*, 272:23064–23068, 1997.
- De los Angeles Costa, Elesgaray, and Loria. Atrial natriuretic peptide influence on nitric oxide system in kidney and heart. *Regul Pept*, 118:151–157, 2004.

- B Lu, NP Gerard, LF Kolakowski, M Bozza, D Zurakowski, O Finco, MC Carroll, and C Gerard. Neutral endopeptidase modulation of septic shock. *J Exp Med*, 181:2271–2275, 1995.
- Lubien, DeMaria A, and Krishnaswamy P et al. Utility of b-natriuretic peptide in detecting diastolic dysfunction: comparison with doppler velocity recordings. *Circulation*, 105:595–601, 2002.
- Lucarelli, Iacoviello, and Dessi-Fulgheri. Associazione di una nuova variante genica del recettore a dei peptidi natriuretici con la familiarità per ipertensione arteriosa. *Ital Heart J 2 (Suppl 6)*, 10, 2001.
- Luchner, Burnett JC Jr, and Jougasaki M et al. Augmentation of the cardiac natriuretic peptides by beta-receptor antagonism: evidence from a population-based study. *J Am Coll Cardiol*, 32:1839–1844, 1998.
- A Luchner, C Hengstenberg, H Lowel, GA Riegger, H Schunkert, and S Holmer. Effect of compensated renal dysfunction on approved heart failure markers: direct comparison of brain natriuretic peptide (bnp) and n-terminal pro-bnp. *Hypertension*, 46:118–23, 2005.
- Ma, Ogawa, and de Bold AJ. Selective upregulation of cardiac brain natriuretic peptide at the transcriptional and translational levels by pro-inflammatory cytokines and by conditioned medium derived from mixed lymphocyte reactions via p38 map kinase. *J Mol Cell Cardiol*, 36:505–513, 2004.
- Maeder, Ammann, Rickli, and Diethelm. Elevation of b-type natriuretic peptide levels in acute respiratory distress syndrome. *Swiss Med Wkly*, 133:515–518, 2003.
- Maffei, Del Ry S, Prontera C, and Clerico A. Increase in circulating levels of cardiac natriuretic peptides after hormone replacement therapy in postmenopausal women. *Clin Sci*, 101:447–453, 2001.

- Magga, Puhakka, and Hietakorpi. Atrial natriuretic peptide, b-type natriuretic peptide, and serum collagen markers after acute myocardial infarction. *J Appl Physiol*, 96:1306–1311, 2004.
- M Magnusson, O Melander, B Israelsson, A Grubb, L Groop, and S Jovinge. Elevated plasma levels of nt-probnp in patients with type 2 diabetes without overt cardiovascular disease. *Diabetes Care*, 27:1929–35, 2004.
- Maisel. B-type natriuretic peptide (bnp) levels: diagnostic and therapeutic potential. (Review) *Rev Cardiovasc Med* 2(suppl 2):S13-18, 2001.
- Maisel, Koon J, and Krishnaswamy P et al. Utility of b-natriuretic peptide as a rapid, point-of-care test for screening patients undergoing echocardiography to determine left ventricular dysfunction. *Am Heart J*. 141:367–374, 2001.
- Maisel, Krishnaswamy P, and Nowak RM et al. Rapid measurement of b-type natriuretic peptide in the emergency diagnosis of heart failure. *N Engl J Med*, 347:161–167, 2002a.
- Maisel, McCord J, and Nowak RM et al. Bedside b-type natriuretic peptide in the emergency diagnosis of heart failure with reduced or preserved ejection fraction. results from the breathing not properly multinational study. *J Am Coll Cardiol*, 41:2010–2017, 2003.
- Maisel, Hollander JE, and Guss D et al. Rapid emergency department heart failure outpatient trial investigators. primary results of the rapid emergency department heart failure outpatient trial (redhot). a multicenter study of b-type natriuretic peptide levels, emergency department decision making, and outcomes in patients presenting with shortness of breath. *J Am Coll Cardiol*, 44:1328–1333, 2004.
- AS Maisel, P Krishnaswamy, RM Nowak, J McCord, JE Hollander, P Duc, T Omland, AB Storrow, WT Abraham, AH Wu, P Clifton, PG Steg, A Westheim, CW Knudsen, A Perez, R Kazanegra,

- HC Herrmann, and Investigators McCullough PA; Breathing Not Properly Multinational Study. Rapid measurement of b-type natriuretic peptide in the emergency diagnosis of heart failure. *N Engl J Med*, 347:161–7, 2002b.
- Mak, DeMaria A, Clopton P, and Maisel AS. Utility of b-natriuretic peptide in the evaluation of left ventricular diastolic function: comparison with tissue doppler imaging recordings. *Am Heart J*, 148: 895–902, 2004.
- Marchioli and Graziano SL. Paraneoplastic syndromes associated with small cell lung cancer. *Chest Surg Clin N Am*, 7:65–80, 1997.
- Marumoto, Hamada M, and Hiwada K. Increased secretion of atrial and brain natriuretic peptides during acute myocardial ischaemia induced by dynamic exercise in patients with angina pectoris. *Clin Sci*, 88:551–556, 1995.
- N Matsukawa, WJ Grzesik, N Takahashi, KN Pandey, S Pang, M Yamauchi, and O Smithies. The natriuretic peptide clearance receptor locally modulates the physiological effects of the natriuretic peptide system. *Proc Natl Acad Sci U S A*, 96:7403–7408, 1999.
- Anand-Srivastava MB. Natriuretic peptide receptor-c signaling and regulation. *Pep- tides*, 26:1044–1059, 2005.
- McCullough, Kuncheria J, and Mathur VS. Diagnostic and therapeutic utility of b-type natriuretic peptide in patients with renal insufficiency and decompensated heart failure. (Review) *Rev Cardiovasc Med*, 5:16–25, 2004a.
- McCullough, Kuncheria, and Mathur VS. Diagnostic and therapeutic utility of b-type natriuretic peptide in patients with renal insufficiency and decompensated heart failure. *Rev Cardiovasc Med*, 5: 16–25, 2004b.

- PA McCullough, RM Nowak, J McCord, JE Hollander, HC Herrmann, PG Steg, P Duc, A Westheim, T Omland, CW Knudsen, AB Storrow, WT Abraham, S Lamba, AH Wu, A Perez, P Clopton, P Krishnaswamy, R Kazanegra, and AS Maisel. B-type natriuretic peptide and clinical judgment in emergency diagnosis of heart failure: analysis from breathing not properly (bnp) multinational study. *Circulation*, 106:416–22, 2002.
- McDonagh, Cunningham AD, and Morrison CE et al. Left ventricular dysfunction, natriuretic peptides, and mortality in an urban population. *Heart*, 86:21–26, 2001.
- McGeoch, Lainchbury J, and Town GI et al. Plasma brain natriuretic peptide after longterm treatment for heart failure in general practice. *Eur J Heart Fail*, 4:479–483, 2002.
- McGirt, Blessing, and Nimjee SM. Correlation of serum brain natriuretic peptide with hyponatremia and delayed ischemic neurological deficits after subarachnoid hemorrhage. *Neurosurgery*, 54:1369–1373, 2004.
- McGrath and de Bold AJ. Determinants of natriuretic peptide gene expression. *Peptides*, 26:933–943, 2005.
- A McGregor, M Richards, E Espiner, T Yandle, and H Ikram. Brain natriuretic peptide administered to man: actions and metabolism. *J Clin Endocrinol Metab*, 70:1103–1107, 1990.
- McLean, Tang B, and Nalos M et al. Increased b-type natriuretic peptide (bnp) level is a strong predictor for cardiac dysfunction in intensive care unit patients. *Anaesth Intensive Care*, 31:21–27, 2003.
- M McNairy, N Gardetto, P Clopton, A Garcia, P Krishnaswamy, R Kazanegra, M Ziegler, and AS Maisel. Stability of b-type natriuretic peptide levels during exercise in patients with congestive heart failure: implications for outpatient monitoring with b-type natriuretic peptide. *Am Heart J*, 143:406–11, 2002.

- Mehra, Uber PA, and Park MH et al. Obesity and suppressed b-type natriuretic peptide levels in heart failure. *J Am Coll Cardiol*, 43: 1590–1595, 2004a.
- MR Mehra, PA Uber, MH Park, RL Scott, HO Ventura, BC Harris, and ED Frohlich. Obesity and suppressed b-type natriuretic peptide levels in heart failure. *J Am Coll Cardiol*, 43:1590–5, 2004b.
- AD Michaels, K Chatterjee, and T De-Marco. Effects of intravenous nesiritide on pulmonary vascular hemodynamics in pulmonary hypertension. *J Card Fail*, 11:425–431, 2005.
- RM Mills, TH LeJemtel, DP Horton, C Liang, R Lang, MA Silver, C Lui, and K Chatterjee. Sustained hemodynamic effects of an infusion of nesiritide (human b-type natriuretic peptide) in heart failure: a randomized, double-blind, placebo-controlled clinical trial; natreacor study group. *J Am Coll Cardiol*, 34:155–162, 1999.
- Mimura, Kanauchi, Ogawa, and Oohara. Resistance to natriuresis in patients with peritonitis carcinomatosa. *Horm Metab Res*, 28:183–186, 1996.
- KS Misono, RT Grammer, H Fukumi, and T Inagami. Rat atrial natriuretic factor: isolation, structure and biological activities of four major peptides. *Biochem Biophys Res Commun*, 123:444–451, 1984.
- Missouris, Grouzmann E, and Buckley MG et al. How does treatment influence endocrine mechanisms in acute severe heart failure? effects on cardiac natriuretic peptides, the renin system, neuropeptide y and catecholamines. *Clin Sci*, 94:591–599, 1998.
- Mizuno, Yoshimura, and Yasue. Aldosterone production is activated in failing ventricle in humans. *Circulation*, 103:72–77, 2001.
- P Moffatt, G Thomas, K Sellin, MC Bessette, F Lafreniere, O Akhouayri, R St-Arnaud, and C Lanctot. Osteocrin is a specific

- ligand of the natriuretic peptide clearance receptor that modulates bone growth. *J Biol Chem*, 282:36454–36462, 2007.
- Morimoto, Hayashino Y, and Shimbo T et al. Is b-type natriuretic peptide-guided heart failure management cost-effective? *Int J Cardiol*, 96:177–181, 2004.
- Morishige, Shimokawa, and Yamawaki. Local adenovirus-mediated transfer of c-type natriuretic peptide suppresses vascular remodeling in porcine coronary arteries in vivo. *J Am Coll Cardiol*, 35: 1040–1047, 2000.
- Morita, Yasue, and Yoshimura. Increased plasma levels of brain natriuretic peptide in patients with acute myocardial infarction. *Circulation*, 88:82–91, 1993.
- Mueller and Buser P. B-type natriuretic peptide (bnp): can it improve our management of patients with congestive heart failure? (*Review*) *Swiss Med Wkly*, 132:618–622, 2002.
- Mueller, Scholer A, and Laule-Kilian K et al. Use of b-type natriuretic peptide in the evaluation and management of acute dyspnea. *N Engl J Med*, 350:647–654, 2004.
- Mukkaddam-Daher, Tremblay J, and Fujio N et al. Alteration of lung atrial natriuretic peptide receptors in genetic cardiomyopathy. *Am J Physiol*, 271:138–145, 1996.
- Mukoyama, Nakao, and Hosoda. Brain natriuretic peptide as a novel cardiac hormone in humans. evidence for an exquisite dual natriuretic peptide system, atrial natriuretic peptide and brain natriuretic peptide. *J Clin Invest*, 87:1402–1412, 1991a.
- Mukoyama, Nakao, and Obata. Augmented secretion of brain natriuretic peptide in acute myocardial infarction. *Biochem Biophys Res Commun volume = 15*, 1991b.

- M Mukoyama, K Nakao, Y Saito, Y Ogawa, K Hosoda, S Suga, G Shirakami, M Jougasaki, and H Imura. Increased human brain natriuretic peptide in congestive heart failure. *N Engl J Med*, 323: 757–758, 1990.
- M Mukoyama, K Nakao, K Hosoda, S Suga, Y Saito, Y Ogawa, G Shirakami, M Jougasaki, K Obata, and H Yasue. Brain natriuretic peptide as a novel cardiac hormone in humans; evidence for an exquisite dual natriuretic peptide system, atrial natriuretic peptide and brain natriuretic peptide. *J Clin Invest*, 87:1402–1412, 1991c.
- Murdoch, McDonagh TA, and Byrne J et al. Titration of vasodilator therapy in chronic heart failure according to plasma brain natriuretic peptide concentration: randomized comparison of the hemodynamic and neuroendocrine effects of tailored versus empirical therapy. *Am Heart J*, 138:1126–1132, 1999.
- M Nagase, T Katafuchi, S Hirose, and T Fujita. Tissue distribution and localization of natriuretic peptide receptor subtypes in stroke-prone spontaneously hypertensive rats. *J Hypertens*, 15:1235–1243, 1997.
- Nagaya, Nishikimi, and Goto. Plasma brain natriuretic peptide is a biochemical marker for the prediction of progressive ventricular remodeling after acute myocardial infarction. *Am Heart J*, 135: 21–28, 1998.
- Nagaya, Nishikimi, and Uematsu. Plasma brain natriuretic peptide as a prognostic indicator in patients with primary pulmonary hypertension. *Circulation*, 102:865–870, 2000a.
- Nagaya, Nishikimi T, and Uematsu M et al. Plasma brain natriuretic peptide as a prognostic indicator in patients with primary pulmonary hypertension. *Circulation*, 102:865–870, 2000b.
- Nakagawa, Ogawa Y, and Itoh H et al. Rapid transcriptional activation and early mrna turnover of brain natriuretic peptide in car-

- diocyte hypertrophy. evidence for brain natriuretic peptide as an “emergency” cardiac hormone against ventricular overload. *J Clin Invest*, 96:1280–1287, 1995.
- Nakamura, Endo H, and Nasu M et al. Value of plasma b type natriuretic peptide measurement for heart disease screening in a japanese population. *Heart*, 87:131–135, 2002.
- Nakamura, Sakai T, and Osawa M et al. Comparison of positive cases for b-type natriuretic peptide and ecg testing for identification of precursor forms of heart failure in an elderly population. *Int Heart J*, 46:477–487, 2005.
- Nakanishi, Saito, and Kishimoto. Role of natriuretic peptide receptor guanylyl cyclase-a in myocardial infarction evaluated using genetically engineered mice. *Hypertension*, 46:1–7, 2005.
- K Nakao, A Sugawara, N Morii, M Sakamoto, T Yamada, H Itoh, S Shiono, Y Saito, K Nishimura, and T Ban. The pharmacokinetics of alpha-human atrial natriuretic polypeptide in healthy subjects. *Eur J Clin Pharmacol*, 31:101–103, 1986.
- Nakayama. The genetic contribution of the natriuretic peptide system to cardiovascular diseases. *Endocr J*, 52:11–21, 2005.
- Nakayama, Soma, and Takahashi. *title = Functional deletion mutation of the, 5'-flanking region of type A natriuretic peptide receptor gene and its association with essential hypertension and left ventricular hypertrophy in the Japanese.*, 86:841–845, 2000a.
- T Nakayama, M Soma, Y Takahashi, D Rehemudula, K Kanmatsuse, and K Furuya. Functional deletion mutation of the 5-flanking region of type a human natriuretic peptide receptor gene and its association with essential hypertension and left ventricular hypertrophy in the japanese. *Circ Res*, 86:841–845, 2000b.

- Nazario, Hu, and Pedram. Atrial and brain natriuretic peptides stimulate the production and secretion of c-type natriuretic peptide from bovine aortic endothelial cells. *J Clin Invest*, 95:1151–1157, 1995.
- Ng, Loke I, and Davies JE et al. Identification of previously undiagnosed left ventricular systolic dysfunction: community screening using natriuretic peptides and electrocardiography. *Eur J Heart Fail*, 5:775–782, 2003a.
- Ng, Loke I, and Davies JE et al. Identification of previously undiagnosed left ventricular systolic dysfunction: community screening using natriuretic peptides and electrocardiography. *Eur J Heart Fail*, 5:775–782, 2003b.
- Ng, Geeranavor, and Jennings SC. Diagnosis of heart failure using urinary natri- uretic peptides. *Clin Sci*, 106:129–133, 2004.
- Nicholls, Lainchbury JG, and Richards AM et al. Brain natriuretic peptide-guided therapy for heart failure. *Ann Med*, 33:422–427, 2001.
- Nielsen, McDonagh TA, Robb SD, and Dargie HJ. Retrospective analysis of the costeffectiveness of using plasma brain natriuretic peptide in screening for left ventricular systolic dysfunction in the general population. *J Am Coll Cardiol*, 41:113–120, 2003.
- Nishikimi, Matsuoka H, and Ishikawa K et al. Antihypertensive therapy reduces increased plasma levels of adrenomedullin and brain natriuretic peptide concomitant with regression of left ventricular hypertrophy in a patient with malignant hypertension. *Hypertens Res*, 19:97–101, 1996.
- Nousiainen, Jantunen E, and Vanninen E et al. Acute neurohumoral and cardiovascular effects of idarubicin in leukemia patients. *Eur J Haematol*, 61:347–353, 1998a.
- Nousiainen, Jantunen, and Vanninen. Acute neurohumoral and cardiovascular effects of idarubicin in leukemia patients. *Eur J Haematol*, 61:347–353, 1998b.

- Nousiainen, Jantunen E, and Vanninen E et al. Natriuretic peptides as markers of cardiotoxicity during doxorubicin treatment for non-hodgkin's lymphoma. *Eur J Haematol*, 62:135–141, 1999a.
- Nousiainen, Jantunen, and Vanninen. Natriuretic peptides as markers of cardiotoxicity during doxorubicin treatment for non-hodgkin's lymphoma. *Eur J Haematol*, 62:135–141, 1999b.
- Nugent, Onuoha, and McEneaney DJ. Variable patterns of atrial natriuretic peptide secretion in man. *Eur J Clin Invest*, 24:267–274, 1994.
- DR Nussenzveig, JA Lewicki, and T Maack. Cellular mechanisms of the clearance function of type c receptors of atrial natriuretic factor. *J Biol Chem*, 265:20952–20958, 1990.
- Ogawa, Itoh, and Tamura. Molecular cloning of the complementary dna and gene that encode mouse brain natriuretic peptide and generation of transgenic mice that overexpress the brain natriuretic peptide gene. *J Clin Invest*, 93:1911–1921, 1994a.
- Ogawa, Oida A, and Sugimura H et al. Clinical significance of blood brain natriuretic peptide level measurement in the detection of heart disease in untreated outpatients: comparison of electrocardiography, chest radiography and echocardiography. *Circ J*, 66:122–126, 2002.
- Ogawa, Qiu, Ogata, and Risono KS. Crystal structure of hormone-bound atrial natriuretic peptide receptor extracellular domain volume = , rotation mechanism for transmembrane signal transduction. *J Biol Chem*, 279:28625–28631, 2004.
- Y Ogawa, H Itoh, N Tamura, S Suga, T Yoshimasa, M Uehira, S Matsuda, S Shiono, H Nishimoto, and K Nakao. Molecular cloning of the complementary dna and gene that encode mouse brain natriuretic peptide and generation of transgenic mice that overexpress the brain natriuretic peptide gene. *J Clin Invest*, 93:1911–1921, 1994b.

- Y Ogawa, H Itoh, Y Yoshitake, M Inoue, T Yoshimasa, T Serikawa, and K Nakao. Molecular cloning and chromosomal assignment of the mouse c-type natriuretic peptide (cnp) gene (nppc): comparison with the human cnp gene (nppc). *Genomics*, 24:383–387, 1994c.
- Ohno, Itoh, and Ikeda. Accelerated reendothelialization with suppressed throm- bogenic property and neointimal hyperplasia of rabbit jugular vein grafts by adenovirus- mediated gene transfer of c-type natriuretic peptide. *Circulation*, 105:1623–1626, 2002.
- Ohta, Watanabe K, and Nakazawa M et al. Carvedilol enhances atrial and brain natriuretic peptide mrna expression and release in rat heart. *J Cardiovasc Pharmacol* 36(suppl 2):S19-23, 2000.
- S Oikawa, M Imai, A Ueno, S Tanaka, T Noguchi, H Nakazato, K Kangawa, A Fukuda, and H Matsuo. Cloning and sequence analysis of cdna encoding a precursor for human atrial natriuretic polypeptide. *Nature*, 309:724–726, 1984.
- Okumura and Iuchi K and Yoshida T et al. Brain natriuretic peptide is a predictor of anthracycline-induced cardiotoxicity. *Acta Haematol*, 104:158–163, 2000.
- Okumura, Iuchi, and Yoshida. Brain natriuretic peptide is a predictor of anthra- cycline-induced cardiotoxicity. *Acta Haematol*, 104:158–163, 2000.
- PM Oliver, JE Fox, R Kim, HA Rockman, HS Kim, RL Reddick, KN Pandey, SL Milgram, O Smithies, and N Maeda. Hypertension, cardiac hypertrophy, and sudden death in mice lacking natriuretic peptide receptor a. *Proc Natl Acad Sci U S A*, 94:14730–14735, 1997.
- PM Oliver, SW John, KE Purdy, R Kim, N Maeda, MF Goy, and O Smithies. Natriuretic peptide receptor 1 expression influences blood pressures of mice in a dose-dependent manner. *Proc Natl Acad Sci U S A*, 95:2547–2551, 1998.

- RC Olney, H Bukulmez, CF Bartels, TC Prickett, EA Espiner, LR Potter, and ML Warman. Heterozygous mutations in natriuretic peptide receptor-b (npr2) are associated with short stature. *J Clin Endocrinol Metab*, 91:1229–1232, 2006.
- Omland, Aakvaag A, and Bonarjee VV et al. Plasma brain natriuretic peptide as an indicator of left ventricular systolic function and long-term survival after acute myocardial infarction. comparison with plasma atrial natriuretic peptide and n-terminal proatrial natriuretic peptide. *Circulation*, 93:1963–1969, 1996a.
- Omland, Aakvaag A, and Vik-Mo H. Plasma cardiac natriuretic peptide determination as a screening test for the detection of patients with mild left ventricular impairment. *Heart*, 76:232–237, 1996b.
- Omland, Persson A, and Ng L et al. N-terminal pro-b-type natriuretic peptide and longterm mortality in acute coronary syndromes. *Circulation*, 106:2913–2918, 2002.
- Omland, Richards AM, and Wergeland R and Vik Mo H. B-type natriuretic peptide and longterm survival in patients with stable coronary artery disease. *Am J Cardiol*, 95:24–28, 2005.
- Ono, Tanabe K, and Asanuma T et al. Doppler echocardiography-derived index of myocardial performance (tei index): comparison with brain natriuretic peptide levels in various heart diseases. *Jpn Circ J*, 65:637–642, 2001.
- Opocher, Rocco, and Carpenne. Atrial natriuretic peptide in cushing’s disease. *J Endocrinol Invest*, 13:133–137, 1990.
- O’Neill, Bott-Silverman CE, and McRae AT 3rd et al. B-type natriuretic peptide levels are not a surrogate marker for invasive hemodynamics during management of patients with severe heart failure. *Am Heart J*, 149:363–369, 2005.

- Packer. The neurohormonal hypothesis: a theory to explain the mechanisms of disease progression in heart failure. (*Review*) *J Am Coll Cardiol*, 20:248–254, 1992.
- Pagani, Stefini F, and Panteghini M. Biological variation in serum concentrations of nterminal pro-brain natriuretic peptide (nt-probnp). (*Abstract*) *Clin Chem*, 49:A34, 2003.
- Palladini, Campana C, and Klersy C et al. Serum n-terminal pro-brain natriuretic peptide is a sensitive marker of myocardial dysfunction in al amyloidosis. *Circulation*, 107:2440–2445, 2003a.
- Palladini, Campana, and Klersy. Serum n-terminal pro-brain natriuretic peptide is a sensitive marker of myocardial dysfunction in al amyloidosis. *Circulation*, 107:2440–2445, 2003b.
- Pandey. Stoichiometric analysis of internalization, recycling and redistribution of photoaffinity-labeled guanylate cyclase/atrial natriuretic factor receptors in cultured murine leydig tumor cells. *J Biol Chem*, 268:4382–4390, 1993.
- Pandey. Dynamics of internalization and sequestration of guanylyl cyclase/atrial natriuretic peptide receptor-a. *Can J Physiol Pharmacol*, 79:631–639, 2001.
- Pandey. Biology of natriuretic peptides and their receptors. *Peptides*, 26:901–932, 2005.
- Pandey, Inagami, and Misono KS. Atrial natriuretic factor receptor on cultured ley- dig tumor cells volume = , ligand binding and photoaffinity labeling. *Biochemistry*, 25:8467–8472, 1986.
- Pandey, Kumar, Li, and Nguyen. Functional domains and expression of truncated atrial natriuretic peptide receptor-a volume = , the carboxyl-terminal regions direct the receptor internalization and sequestration in cos- / cells. *Mol Pharmacol*, 57:259–267, 2000.

- KN Pandey. Intracellular trafficking and metabolic turnover of ligand-bound guanylyl cyclase/atrial natriuretic peptide receptor-a into subcellular compartments. *Mol Cell Biochem*, 230:61–72, 2002.
- K Pankow, Y Wang, F Gembardt, E Krause, X Sun, G Krause, HP Schultheiss, WE Siems, and T Walther. Successive action of meprin a and neprilysin catabolizes b-type natriuretic peptide. *Circ Res*, 101:875–882, 2007.
- Panteghini and Clerico A. Understanding the clinical biochemistry of n-terminal pro-b-type natriuretic peptide: the prerequisite for its optimal clinical use. (Review) *Clin Lab*, 2004.
- Panteghini, Cuccia C, and Bonetti G et al. Rapid determination of brain natriuretic peptide in patients with acute myocardial infarction. *Clin Chem Lab Med*, 41:164–168, 2003.
- Parlapiano, Campana, and Alessandri. Plasma atrial natriuretic hormone in hyperthyroidism. *Endocr Res*, 24:105–112, 1998.
- Parsonage, Galbraith AJ, Koerbin GL, and Potter JM. Value of b-type natriuretic peptide for identifying significantly elevated pulmonary artery wedge pressure in patients treated for established chronic heart failure secondary to ischemic or idiopathic dilated cardiomyopathy. *Am J Cardiol*, 95:883–885, 2005.
- JB Patel, ML Valencik, AM Pritchett, JC Burnett, JA McDonald, and MM Redfield. Cardiac-specific attenuation of natriuretic peptide a receptor activity accentuates adverse cardiac remodeling and mortality in response to pressure overload. *Am J Physiol Heart Circ Physiol*, 289:H777–H784, 2005.
- Pedersen, Pedersen, and Jensen KT. Pulsatile secretion of atrial natriuretic peptide and brain natriuretic peptide in healthy humans. *Clin Sci*, 97:201–206, 1999.
- Perras, Schultes, and Behn. Intranasal atrial natriuretic peptide acts as central nervous inhibitor of the hypothalamo-pituitary-adrenal

- stress system in humans. *J Clin Endocrinol Metab*, 89:4642–4648, 2004.
- Pfeifer, Ruth, and Dostmann. Structure and function of cgmp-dependent protein kinases. *Rev Physiol Biochem Pharmacol*, 135: 105–149, 1999.
- A Pfeifer, A Aszodi, U Seidler, P Ruth, F Hofmann, and R Fassler. Intestinal secretory defects and dwarfism in mice lacking cgmp-dependent protein kinase ii. *Science*, 274:2082–2086, 1996.
- R Pfister and CA Schneider. Natriuretic peptides bnp and nt-pro-bnp: established laboratory markers in clinical practice or just perspectives? *Clin Chim Acta*, 349:25–38, 2004.
- Pichon, Cvitkovic F, and Hacene K et al. Drug-induced cardiotoxicity studied by longitudinal b-type natriuretic peptide assays and radionuclide ventriculography. *In vivo*, 19:567–576, 2005.
- Pilo, Iervasi, and Clerico. The circulatory model in metabolic studies of rapidly renewed hormones volume = , application to anp kinetics. *Am J Physiol*, 274, 1998.
- JG Porter, A Arfsten, F Fuller, JA Miller, LC Gregory, and JA Lewicki. Isolation and functional expression of the human atrial natriuretic peptide clearance receptor cDNA. *Biochem Biophys Res Commun*, 171:796–803, 1990.
- Potter. Phosphorylation-dependent regulation of the guanylyl cyclase-linked natriuretic peptide receptor volume = , dephosphorylation is a mechanism of desensitization. *Biochemistry*, 37:2422–2429, 1998a.
- Potter and Garber DL. Dephosphorylation of the guanyl cyclase-a receptor causes desensitization. *J Biol Chem*, 267:14531–14534, 1992.

- Potter and Garber DL. Protein kinase c-dependent desensitization of the atrial natriuretic peptide receptor is mediated by dephosphorylation. *J Biol Chem*, 269:14636–14642, 1994.
- Potter and Hunter. Identification and characterization of the major phosphorylation sites of the b-type natriuretic peptide receptor. *J Biol Chem*, 273:15533–15539, 1998a.
- Potter and Hunter. Phosphorylation of the kinase homology domain is essential for activation of the a-type natriuretic peptide receptor. *Mol Cell Biol*, 18:2164–2172, 1998b.
- Potter and Hunter T. Phosphorylation of the kinase homology domain is essential for activation of the a-type natriuretic peptide receptor. *Mol Cell Biol*, 18:2164–2172, 1998.
- LR Potter. Phosphorylation-dependent regulation of the guanylyl cyclase-linked natriuretic peptide receptor b: dephosphorylation is a mechanism of desensitization. *Biochemistry*, 37:2422–2429, 1998b.
- LR Potter and DL Garbers. Dephosphorylation of the guanylyl cyclase-a receptor causes desensitization. *J Biol Chem*, 267:14531–14534, 1992.
- LR Potter and DL Garbers. Protein kinase c-dependent desensitization of the atrial natriuretic peptide receptor is mediated by dephosphorylation. *J Biol Chem*, 269:14636–14642, 1994.
- LR Potter and T Hunter. Activation of pkc stimulates the dephosphorylation of natriuretic peptide receptor-b at a single serine residue: a possible mechanism of heterologous desensitization. *J Biol Chem*, 275:31099–31106, 2000.
- R Potthast, SE Abbey-Hosch, LK Antos, JS Marchant, M Kuhn, and LR Potter. Calcium-dependent dephosphorylation mediates the hyperosmotic and lysophosphatidic acid-dependent inhibition of natriuretic peptide receptor-b/guanylyl cyclase-b. *J Biol Chem*, 279:48513–48519, 2004.

- Prontera, Emdin M, and Zucchelli GC et al. Fully-automated ntprobnp and irma methods for bnp and anp in heart failure and healthy subjects. (*Letter*) *Clin Chem*, 49:1552–1554, 2003.
- Pruszczyk, Kostrubiec, and Bochowicz. N-terminal pro-brain natriuretic peptide in patients with acute pulmonary embolism. *Eur Respir J*, 22:649–653, 2003.
- WG Pyle and RJ Solaro. At the crossroads of myocardial signaling: the role of z-discs in intracellular signaling and cardiac function. *Circ Res*, 94:296–305, 2004.
- Qian, Haruno, and Asada. Local expression of c-type natriuretic peptide suppresses inflammation, eliminates shear stress-induced thrombosis, and prevents neointima formation through enhanced nitric oxide production in rabbit injured carotid arteries. *Circ Res*, 91: 1063–1069, 2002.
- SN Rahman, GE Kim, AS Mathew, CA Goldberg, R Allgren, RW Schrier, and JD Conger. Effects of atrial natriuretic peptide in clinical acute renal failure. *Kidney Int*, 45:1731–1738, 1994.
- Rahmutula, Nakayama, and Soma. Association study between the variants of the human anp gene and essential hypertension. *Hypertens Res*, 24:291–294, 2001.
- Ramasawmy, Kotea, and Lu. A new polymorphic restriction site at the human atrial natriuretic peptide (hanp) gene locus. *Hum Genet*, 91:509–510, 1993.
- Ray, Arthaud M, and Lefort Y et al. Usefulness of b-type natriuretic peptide in elderly patients with acute dyspnea. *Intensive Care Med*, 30:2230–2236, 2004.
- I Raymond, BA Groenning, PR Hildebrandt, JC Nilsson, M Baumann, J Trawinski, and F Pedersen. The influence of age, sex and other variables on the plasma level of n-terminal pro brain natriuretic

- peptide in a large sample of the general population. *Heart*, 89:745–751, 2003.
- Redfield, Rodeheffer RJ, and Jacobsen SJ et al. Plasma brain natriuretic peptide concentration: impact of age and gender. *J Am Coll Cardiol*, 40:976–982, 2002.
- Remes, Miettinen H, Reunanen A, and Pyorala K. Validity of clinical diagnosis of heart failure in primary health care. *Eur Heart J*, 12: 315–321, 1991.
- Remme and Swedberg K. Task force report. guidelines for the diagnosis and treatment of chronic heart failure. *Eur Heart J*, 22:1527–1560, 2001.
- Richards. Outpatient management of heart failure. (*Review*) *Heart Fail Rev*, 8:345–348, 2003.
- Richards and Troughton RW. Nt-probnp in heart failure: therapy decisions and monitoring. (*Review*) *Eur J Heart Fail*, 6:351–354, 2004.
- Richards, Nicholls MG, and Yandle TG et al. Plasma n-terminal pro-brain natriuretic peptide and adrenomedullin: new neurohormonal predictors of left ventricular function and prognosis after myocardial infarction. *Circulation*, 97:1921–1929, 1998.
- Richards, Doughty R, and Nicholls MG et al. Plasma n-terminal pro-brain natriuretic peptide and adrenomedullin: prognostic utility and prediction of benefit from carvedilol in chronic ischemic left ventricular dysfunction. australia-new zealand heart failure group. *J Am Coll Cardiol*, 37:1781–1787, 2001.
- Richards, Lainchbury JG, and Nicholls MG et al. Bnp in hormone-guided treatment of heart failure. (*Review*) *Trends Endocrinol Metab*, 13:151–155, 2002.

- AM Richards, G Tonolo, P Montorsi, J Finlayson, R Fraser, G Inglis, A Towrie, and JJ Morton. Low dose infusions of 26- and 28-amino acid human atrial natriuretic peptides in normal man. *J Clin Endocrinol Metab*, 66:465–472, 1988.
- AM Richards, R Doughty, MG Nicholls, S Macmahon, H Ikram, N Sharpe, EA Espiner, C Frampton, and TG Yandle. Neurohumoral prediction of benefit from carvedilol in ischemic left ventricular dysfunction, australia-new zealand heart failure group. *Circulation*, 99: 786–92, 1999.
- Brunner-La Rocca, Kaye DM, and Woods RL. Effects of intravenous brain natriuretic peptide on regional sympathetic activity in patients with chronic heart failure as compared with healthy control subjects. *J Am Coll Cardiol*, 37:1221–1227, 2001.
- Rodeheffer. Measuring plasma b-type natriuretic peptide in heart failure: good to go in 2004? (Review) *J Am Coll Cardiol*, 44:740–749, 2004.
- Rollin, Mediero, and Roldan-Pallares. Natriuretic peptide system in the human retina. *Mol Vis*, 10:15–22, 2004.
- RA Rose and WR Giles. Natriuretic peptide c receptor signalling in the heart and vasculature. *J Physiol*, 586:353–366, 2008.
- Ruskoaho. Cardiac hormones as diagnostic tools in heart failure. *Endocr Rev*, 24:341–356, 2003a.
- Ruskoaho. Cardiac hormones as diagnostic tools in heart failure. (Review) *Endocr Rev*, 24:341–356, 2003b.
- Ruskoaho, Leskinen, and Magga. Mechanisms of mechanical load-induced atrial natriuretic peptide secretion volume = , role of endothelin, nitric oxide, and angiotensin ii. *J Mol Med*, 75:876–885, 1997.

- H Ruskoaho. Cardiac hormones as diagnostic tools in heart failure. *Endocr Rev*, 24:341–356, 2003c.
- Magga J Ruskoaho H, Leskinen H. Mechanisms of mechanical load-induced atrial natriuretic peptide secretion volume = , role of endothelin, nitric oxide, and angiotensin ii. *J Mol Med*, 75:876–885, 1997.
- Rutledge, Sun, and Ross EA. Polymorphisms within the atrial natriuretic peptide gene in essential hypertension. *J Hypertens*, 13: 953–955, 1995.
- Sabatine, Morrow DA, and de Lemos JA et al. Multimarker approach to risk stratification in non-ST elevation acute coronary syndromes: simultaneous assessment of troponin I, C-reactive protein, and B-type natriuretic peptide. *Circulation*, 105:1760–1763, 2002.
- Sabatine, Morrow DA, and de Lemos JA et al. TIMI study group. Acute changes in circulating natriuretic peptide levels in relation to myocardial ischemia. *J Am Coll Cardiol*, 44:1988–1995, 2004.
- K Sabrane, MN Kruse, L Fabritz, B Zetsche, D Mitko, BV Skryabin, M Zwiener, HA Baba, M Yanagisawa, and M Kuhn. Vascular endothelium is critically involved in the hypotensive and hypovolemic actions of atrial natriuretic peptide. *J Clin Invest*, 115:1666–1674, 2005.
- Sackett and Haynes RB. The architecture of diagnostic research. *Br Med J*, 324:539–541, 2002.
- JD Sackner-Bernstein, M Kowalski, M Fox, and K Aaronson. Short-term risk of death after treatment with nesiritide for decompensated heart failure: a pooled analysis of randomized controlled trials. *Jama*, 293:1900–1905, 2005a.
- JD Sackner-Bernstein, HA Skopicki, and KD Aaronson. Risk of worsening renal function with nesiritide in patients with acutely decompensated heart failure. *Circulation*, 111:1487–1491, 2005b.

- Sagnella. Vasopeptidase inhibitors. (Review) *J Renin Angiotensin Aldosterone Syst*, 3:90–95, 2002.
- Saito, Nakao K, and Nishimura K et al. Clinical application of atrial natriuretic polypeptide in patients with congestive heart failure: beneficial effects on left ventricular function. *Circulation*, 76:115–124, 1987a.
- Y Saito, K Nakao, K Nishimura, A Sugawara, K Okumura, K Obata, R Sonoda, T Ban, H Yasue, and H Imura. Clinical application of atrial natriuretic polypeptide in patients with congestive heart failure: beneficial effects on left ventricular function. *Circulation*, 76:115–124, 1987b.
- Salo, Jimenez, and Kuhn. Urinary excretion of urodilatin in patients with cir- rhosis. *Hepatology*, 24:1428–1432, 1996.
- Sanderson, Chan WW, and Hung YT et al. Effect of low dose beta blockers on atrial and ventricular (b type) natriuretic factor in heart failure: a double blind, randomised comparison of metoprolol and a third generation vasodilating beta blocker. *Br Heart J*, 74:502–507, 1995.
- Sandri, Salvatici M, and Cardinale D et al. N-terminal pro-b-type natriuretic peptide after high-dose chemotherapy: a marker predictive of cardiac dysfunction? *Clin Chem*, 51:1405–1410, 2005.
- Sarzani, Dessi-Fulgheri, and Salvi. A novel promoter variant of the natriuretic peptide clearance receptor gene is associated with lower atrial natriuretic peptide and high- er blood pressure in obese hypertensives. *J Hypertens*, 17:1301–1305, 1999.
- Sarzani, Strazzullo P, and Salvi F et al. Natriuretic peptide clearance receptor alleles and susceptibility to abdominal adiposity. *Obes Res*, 12:351–356, 2004.

- Sayama, Nakamura Y, Saito N, and Kinoshita M. Why is the concentration of plasma brain natriuretic peptide in elderly inpatients greater than normal? *Coron Artery Dis*, 10:537–540, 1999.
- Schorr, Beige, and Ringel. Hpaai polymorphism of the atrial natriuretic peptide gene and blood pressure response to salt intake in normotensive men. *J Hypertens*, 15:715–718, 1997.
- RW Schrier and WT Abraham. Hormones and hemodynamics in heart failure. *New England Journal of Medicine*, 341:577–85, 1999.
- Schultz, Faber, and Kistorp. 2004.
- Scotland, Ahluwalia, and Hobbs AJ. C-type natriuretic peptide in vascular physiology and disease. *Pharmacol Ther*, 105:85–89, 2005.
- Segawa, Nakamura M, and Itai K et al. Plasma b-type natriuretic peptide levels and risk factors for congestive heart failure in a japanese general population. *Int Heart J*, 46:465–475, 2005.
- Shimizu, Masuta K, and Aono K et al. Molecular forms of human brain natriuretic peptide in plasma. *Clin Chim Acta*, 316:129–135, 2002.
- Shimizu, Masuta K, and Asada H et al. Characterization of molecular forms of probrain natriuretic peptide in human plasma. *Clin Chim Acta*, 334:233–239, 2003a.
- Shimizu, Masuta, and Asada. Characterization of molecular forms of probrain natriuretic peptide in human plasma. *Clin Chim Acta*, 334:233–239, 2003b.
- Sim, Hampton D, and Phillips C et al. The use of brain natriuretic peptide as a screening test for left ventricular systolic dysfunction - cost-effectiveness in relation to open access echocardiography. *Fam Pract*, 20:570–574, 2003.

- Squire, Orn S, and Ng LL et al. Plasma natriuretic peptides up to 2 years after acute myocardial infarction and relation to prognosis: an optimal substudy. *J Card Fail*, 11:492–497, 2005.
- Stanek, Frey B, and Hulsman M et al. Prognostic evaluation of neurohumoral plasma levels before and during beta-blocker therapy in advanced left ventricular dysfunction. *J Am Coll Cardiol*, 38:436–442, 2001.
- Stanton and Hansen M and Wijesundera HC et al. A direct comparison of the natriuretic peptides and their relationship to survival in chronic heart failure of a presumed non-ischaemic origin. *Eur J Heart Fail*, 7:557–565, 2005.
- Steg, Joubin L, and McCord J et al. B-type natriuretic peptide and echocardiographic determination of ejection fraction in the diagnosis of congestive heart failure in patients with acute dyspnea. *Chest*, 128: 21–29, 2005.
- Steinhilber, Cochrane, and Field LJ. Hypotension in transgenic mice expressing atrial natriuretic factor fusion genes. *Hypertension*, 16: 301–307, 1990a.
- ME Steinhilber, KL Cochrane, and LJ Field. Hypotension in transgenic mice expressing atrial natriuretic factor fusion genes. *Hypertension*, 16:301–307, 1990b.
- SL Stephenson and AJ Kenny. The hydrolysis of alpha-human atrial natriuretic peptide by pig kidney microvillar membranes is initiated by endopeptidase-24,11. *Biochem J*, 243:183–187, 1987.
- Straub, Hall, and Kramer BK. Atrial natriuretic factor and digoxin-like immunoreactive factor in diabetic patients volume = , their interrelation and the influence of the autonomic nervous system. *J Clin Endocrinol Metab*, 81:3385–3389, 1996.

- Struthers. Plasma concentrations of brain natriuretic peptide: will this new test reduce the need for cardiac investigations? (*Editorial*) *Br Heart J*, 70:397–398, 1993.
- T Sudoh, K Kangawa, N Minamino, and H Matsuo. A new natriuretic peptide in porcine brain. *Nature*, 332:78–81, 1988.
- T Sudoh, N Minamino, K Kangawa, and H Matsuo. C-type natriuretic peptide (cnp): a new member of natriuretic peptide family identified in porcine brain. *Biochem Biophys Res Commun*, 168:863–870, 1990.
- Suga, Itoh, and Komatsu. Cytokine-induced c-type natriuretic peptide (cnp) secretion from vascular endothelial cells - evidence for cnp as a novel autocrine/paracrine regulator from endothelial cells. *Endocrinology*, 133:3038–3041, 1993a.
- S Suga, K Nakao, K Hosoda, M Mukoyama, Y Ogawa, G Shirakami, H Arai, Y Saito, Y Kambayashi, K Inouye, and H Imura. Receptor selectivity of natriuretic peptide family, atrial natriuretic peptide, brain natriuretic peptide, and c-type natriuretic peptide. *Endocrinology*, 130:229–239, 1992a.
- S Suga, K Nakao, H Itoh, Y Komatsu, Y Ogawa, N Hama, and H Imura. Endothelial production of c-type natriuretic peptide and its marked augmentation by transforming growth factor-beta; possible existence of vascular natriuretic peptide system. *J Clin Invest*, 90:1145–1149, 1992b.
- S Suga, K Nakao, I Kishimoto, K Hosoda, M Mukoyama, H Arai, G Shirakami, Y Ogawa, Y Komatsu, and O Nakagawa. Phenotype-related alteration in expression of natriuretic peptide receptors in aortic smooth muscle cells. *Circ Res*, 71:34–39, 1992c.
- S Suga, H Itoh, Y Komatsu, Y Ogawa, N Hama, T Yoshimasa, and K Nakao. Cytokine-induced c-type natriuretic peptide (cnp) secretion from vascular endothelial cells-evidence for cnp as a novel au-

- tocrine/paracrine regulator from endothelial cells. *Endocrinology*, 133:3038–3041, 1993b.
- Sugawara, Nakao, and Itoh. Cosecretion of peptides derived from gamma- human atrial natriuretic polypeptide in normal volunteers and patients with essential hyper- tension and adrenal disorders. *J Hypertens*, 6(Suppl. 4):S327–329, 1988.
- Suzuki and Hayashi D and Yamazaki T et al. Elevated b-type natriuretic peptide levels after anthracycline administration. *Am Heart J*, 136:362–363, 1998.
- Suzuki, Hayashi, and Yamazaki. Elevated b-type natriuretic peptide levels after anthracycline administration. *Am Heart J*, 136:362–363, 1998.
- Suzuki, Yamaoki K, and Nakajima O et al. Screening for cardiac dysfunction in asymptomatic patients by measuring b-type natriuretic peptide levels. *Jpn Heart J*, 41:205–214, 2000.
- K Sward, F Valsson, J Sellgren, and SE Ricksten. Differential effects of human atrial natriuretic peptide and furosemide on glomerular filtration rate and renal oxygen consumption in humans. *Intensive Care Med*, 31:79–85, 2005.
- K Swedberg, J Cleland, H Dargie, H Drexler, F Follath, M Komajda, L Tavazzi, OA Smiseth, A Gavazzi, A Haverich, A Hoes, T Jaarsma, J Korewicki, S Levy, C Linde, JL Lopez-Sendon, MS Nieminen, L Pierard, and WJ Remme. Guidelines for the diagnosis and treatment of chronic heart failure: executive summary (update 2005). *Eur Heart J*, 26:1115–40, 2005.
- Takeda, Fukutomi T, and Suzuki S et al. Effects of carvedilol on plasma b-type natriuretic peptide concentration and symptoms in patients with heart failure and preserved ejection fraction. *Am J Cardiol*, 94: 448–453, 2004.

- Takemura, Takatsu, and Doyama. Expression of atrial and brain natriuretic peptides and their genes in hearts of patients with cardiac amyloidosis. *J Am Coll Cardiol*, 31:254–265, 1998.
- Takeuchi, Ohmori, and Kondo. Potentiation of c-type natriuretic peptide with ultrasound and microbubbles to prevent neointimal formation after vascular injury in rats. *Cardiovasc Res*, 58:231–238, 2003.
- Talwar, Squire IB, and Davies JE et al. Plasma n-terminal pro-brain natriuretic peptide and the ecg in the assessment of left-ventricular systolic dysfunction in a high risk population. *Eur Heart J*, 20: 1736–1744, 1999.
- S Talwar, IB Squire, PF Downie, AM Mccullough, MC Campton, JE Davies, DB Barnett, and LL Ng. Profile of plasma n-terminal probnp following acute myocardial infarction; correlation with left ventricular systolic dysfunction. *Eur Heart J*, 21:1514–21, 2000.
- N Tamura, Y Ogawa, H Chusho, K Nakamura, K Nakao, M Suda, M Kasahara, R Hashimoto, G Katsuura, M Mukoyama, H Itoh, Y Saito, I Tanaka, H Otani, and M Katsuki. Cardiac fibrosis in mice lacking brain natriuretic peptide. *Proc Natl Acad Sci U S A*, 97:4239–4244, 2000.
- N Tamura, TD Chrisman, and DL Garbers. The regulation and physiological roles of the guanylyl cyclase receptors. *Endocr J*, 48:611–634, 2001.
- N Tamura, LK Doolittle, RE Hammer, JM Shelton, JA Richardson, and DL Garbers. Critical roles of the guanylyl cyclase b receptor in endochondral ossification and development of female reproductive organs. *Proc Natl Acad Sci U S A*, 101:17300–17305, 2004.
- Tanaka, Nagai, and Date. Effects of bradykinin on cardiovascular remodeling in renovascular hypertensive rats. *Hypertens Res*, 27:865–875, 2005.

- ten Wolde, Tulevski II, and Mulder JW et al. Brain natriuretic peptide as a predictor of adverse outcome in patients with pulmonary embolism. *Circulation*, 107:2082–2084, 2003a.
- ten Wolde, Tulevski, and Mulder JW. Brain natriuretic peptide as a predictor of adverse outcome in patients with pulmonary embolism. *Circulation*, 107:2082–2084, 2003b.
- Tenhunen, Szokodi, and Ruskoaho. Posttranscriptional activation of bnp gene expression in response to increased left ventricular wall stressvolume = , role of calcineurin and pkc. *Regul Pept*, 128:187–196, 2005.
- Thomas, Fox KF, Coats AJ, and Sutton GC. The epidemiological enigma of heart failure with preserved systolic function. *Eur J Heart Fail*, 6:125–136, 2004.
- Thompson, Skrabal CA, and Loebe M et al. Plasma neurohormone levels correlate with left ventricular functional and morphological improvement in lvad patients. *J Surg Res*, 123:25–32, 2005.
- DJ Thuerauf, DS Hanford, and CC Glembofski. Regulation of rat brain natriuretic peptide transcription; a potential role for gata-related transcription factors in myocardial cell gene expression. *J Biol Chem*, 269:17772–17775, 1994.
- Toth, Vuorinen KH, and Vuolteenaho O et al. Hypoxia stimulates release of anp and bnp from perfused rat ventricular myocardium. *Am J Physiol* 266(4 Pt 2):H1572-1580, 1994.
- Totsune, Takahashi, and Satoh. Urinary immunoreactive brain natriuretic peptide in patients with renal disease. *Regul Pept*, 63: 141–146, 1996.
- Trindade and Rouleau JL. Vasopeptidase inhibitors: potential role in the treatment of heart failure. (Review) *Heart Fail Monit*, 2:2–7, 2001.

- Troughton and Frampton CM and Yandle TG et al. Treatment of heart failure guided by plasma aminoterminal brain natriuretic peptide (n-bnp) concentrations. *Lancet*, 355:1126–1130, 2000.
- Troughton, Richards AM, and Nicholls MG. Individualized treatment of heart failure. *Intern Med J*, 31:138–141, 2001.
- RW Troughton, CM Frampton, TG Yandle, EA Espiner, MG Nicholls, and AM Richards. Treatment of heart failure guided by plasma aminoterminal brain natriuretic peptide (n-bnp) concentrations. *Lancet*, 355:1126–30, 2000.
- H Tsuji, N Nishino, Y Kimura, K Yamada, M Nukui, S Yamamoto, T Iwasaka, and H Takahashi. Haemoglobin level influences plasma brain natriuretic peptide concentration. *Acta Cardiol*, 59:527–31, 2004.
- T Tsuji and T Kunieda. A loss-of-function mutation in natriuretic peptide receptor 2 (npr2) gene is responsible for disproportionate dwarfism in cn/cn mouse. *J Biol Chem*, 280:14288–14292, 2005.
- Tsukagoshi, Shimizu, and Kawata. Atrial natriuretic peptide inhibits tumor necrosis factor- α production by interferon- γ -activated macrophages via suppression of p38 mitogen-activated protein kinase and nuclear factor- κ B activation. *Reg Pept*, 99: 21–29, 2001.
- Tsunoda, Mendelsohn FAO, and Sexton PM et al. Decreased atrial natriuretic peptide binding in renal medulla in rats with chronic heart failure. *Circ Res*, 62:155–161, 1988.
- Tsutamoto, Kanamori T, and Morigami N et al. Possibility of down-regulation of atrial natriuretic peptide receptor coupled to guanylate cyclase in peripheral vascular beds of patients with chronic severe heart failure. *Circulation*, 87:70–75, 1993.

- Tsutamoto, Wada A, and Maeda K et al. Plasma brain natriuretic peptide level as a biochemical marker of morbidity and mortality in patients with asymptomatic or minimally symptomatic left ventricular dysfunction. comparison with plasma angiotensin ii and endothelin-1. *Eur Heart J*, 20:1799–1807, 1999.
- Tsutamoto, Wada A, and Maeda K et al. Effect of spironolactone on plasma brain natriuretic peptide and left ventricular remodeling in patients with congestive heart failure. *J Am Coll Cardiol*, 37: 1228–1233, 2001.
- T Tsutamoto, A Wada, K Maeda, T Hisanaga, Y Maeda, D Fukai, M Ohnishi, Y Sugimoto, and M Kinoshita. Attenuation of compensation of endogenous cardiac natriuretic peptide system in chronic heart failure: prognostic role of plasma brain natriuretic peptide concentration in patients with chronic symptomatic left ventricular dysfunction. *Circulation*, 96:509–16, 1997.
- T Tsutamoto, H Sakai, Wada, C Ishikawa, K Ohno, M Fujii, T Yamamoto, T Takayama, T Dohke, and M Horie. Torasemide inhibits transcardiac extraction of aldosterone in patients with congestive heart failure. *J Am Coll Cardiol*, 44:2252–3, 2004.
- II Tulevski, BJ Mulder, and DJ van Veldhuisen. Utility of a bnp as a marker for rv dysfunction in acute pulmonary embolism. *J Am Coll Cardiol*, 39:2080, 2002.
- Ueda, Yokouchi M, and Suzuki T et al. Prognostic value of high plasma brain natriuretic peptide concentrations in very elderly persons. *Am J Med*, 114:266–270, 2003.
- Uusimaa, Ruskoaho H, and Vuolteenaho O et al. Plasma vasoactive peptides after acute myocardial infarction in relation to left ventricular dysfunction. *Int J Cardiol*, 69:5–14, 1999.
- Valle, Aspromonte N, and Barro S et al. The nt-probnp assay identifies

- very elderly nursing home residents suffering from pre-clinical heart failure. *Eur J Heart Fail*, 7:542–551, 2005.
- van den Meiracker, Lameris TW, van de Ven LL, and Boomsma F. Increased plasma concentration of natriuretic peptides by selective beta1-blocker bisoprolol. *J Cardiovasc Pharmacol*, 42:462–468, 2003.
- Vanderheyden and Goethals M andVerstreken S et al. Wall stress modulates brain natriuretic peptide production in pressure overload cardiomyopathy. *J Am Coll Cardiol*, 44:2349–2354, 2004.
- Vanderheyden, Bartunek J, and Goethals M. Brain and other natriuretic peptides: molecular aspects. *Eur J Heart Fail*, 6:261–268, 2004.
- Y Vanneste, A Michel, R Dimaline, T Najdovski, and M Deschodt-Lanckman. Hydrolysis of alpha-human atrial natriuretic peptide in vitro by human kidney membranes and purified endopeptidase-24,11 evidence for a novel cleavage site. *Biochem J*, 254:531–537, 1988.
- Vasan, Benjamin EJ, and Levy D. Prevalence, clinical features and prognosis of diastolic heart failure: an epidemiologic perspective. *J Am Coll Cardiol*, 26:1565–1574, 1995.
- Vasan, Benjamin EJ, and Larson MG et al. Plasma natriuretic peptides for community screening for left ventricular hypertrophy and systolic dysfunction. *JAMA*, 288:1252–1259, 2002.
- Vatta, Presas, and Bianciotti LG. B and c types natriuretic peptides modulate norepinephrine uptake and release in the rat hypothalamus. *Regul Pept*, 16:175–184, 1996.
- Vatta, Presas, and Bianciotti LG. B and c types natriuretic peptides modify nor- epinephrine uptake and release in the rat adrenal medulla. *Peptides*, 18:1483–1489, 1997.
- Ventura, Gomes, and Reis WL. Nitrergic modulation of vasopressin, oxytocin and atrial natriuretic peptide secretion in response to

- sodium intake and hypertonic blood volume expansion. *Braz J Med Biol Res*, 35:1101–1109, 2002.
- Vesely. Natriuretic peptides and acute renal failure. *Am J Physiol Renal Physiol*, 285:F167–177, 2003a.
- Vesely. Natriuretic peptides and acute renal failure. (Review) *Am J Physiol Renal Physiol*, 285:F167–177, 2003b.
- Vieira, Gao, Nikonova, and Maack. Molecular and cellular physiology of the dissociation of atrial natriuretic peptide from guanylyl cyclase a receptors. *J Biol Chem*, 276:36438–36445, 2001a.
- MA Vieira, M Gao, LN Nikonova, and T Maack. Molecular and cellular physiology of the dissociation of atrial natriuretic peptide from guanylyl cyclase a receptors. *J Biol Chem*, 276:36438– 36445, 2001b.
- La Villa, Riccardi, and Lazzeri. Blunted natriuretic response to low-dose brain natriuretic peptide infusion in non azotemic cirrhotic patients with ascites and avid sodium retention. *Hepatology*, 22:1745–1750, 1995.
- Vollmar. The role of atrial natriuretic peptide in the immune system. *Peptides*, 26:1086–1094, 2005.
- Vollmar and Kiemer AK. Immunomodulatory and cytoprotective function of atrial natriuretic peptide. *Crit Rev Immunol*, 21:473–485, 2001.
- Vollmar and Schultz. Expression and differential regulation of natriuretic peptides in mouse macrophages. *J Clin Invest*, 95:2442–2450, 1995.
- Vollmar, Wolf, and Schulz. Co-expression of the natriuretic peptides (anp, bnp, cnp) and their receptors in normal and acutely involuted rat thymus. *J Neuroimmunol*, 57:117–127, 1995.

- Vollmer, Mytzka, Arendt, and Schulz. Atrial natriuretic peptide in bovine corpus luteum. *Endocrinology*, 123:762–767, 1988.
- Wallen, Landahl S, and Hedner T et al. Brain natriuretic peptide predicts mortality in the elderly. *Heart*, 77:264–267, 1997.
- Walther, Klostermann, and Hering-Walther. Fibrosis rather than blood pressure determines cardiac bnp expression in mice. *Regul Pept*, 116:95–100, 2003.
- Wang, Larson MG, and Levy D et al. Impact of age and sex on plasma natriuretic peptide levels in healthy adults. *Am J Cardiol*, 90:254–258, 2002.
- Wang, Larson MG, and Levy D et al. Impact of obesity on plasma natriuretic peptide levels. *Circulation*, 109:594–600, 2004a.
- DJ Wang, TC Dowling, D Meadows, T Ayala, J Marshall, S Minshall, N Greenberg, E Thattassery, ML Fisher, K Rao, and SS Gottlieb. Nesiritide does not improve renal function in patients with chronic heart failure and worsening serum creatinine. *Circulation*, 110:1620–1625, 2004b.
- TJ Wang, MG Larson, D Levy, EJ Benjamin, EP Leip, PW Wilson, and RS Vasan. Impact of obesity on plasma natriuretic peptide levels. *Circulation*, 109:594–600, 2004c.
- Y Wang, MC deWaard, A Sterner-Kock, H Stepan, HP Schultheiss, DJ Duncker, and T Walther. Cardiomyocyte-restricted overexpression of c-type natriuretic peptide prevents cardiac hypertrophy induced by myocardial infarction in mice. *Eur J Heart Fail*, 9: 548–557, 2007.
- Waschek. Developmental actions of natriuretic peptides in the brain and skeleton. *Cell Mol Life Sci*, 61:2332–2342, 2004.

- Weber, Blumenthal, and Hartung. Anp inhibits tnf-alpha-induced endothelial mcp-1 expression - involvement of p38 mapk and mkp-1. *J Leukoc Biol*, 74:932–941, 2003.
- Widecka, Ciechanowicz, and Adler. Analysis of polymorphisms sma (hpa ii) and sca i gene precursors of atrial natriuretic peptide (anp) in patients with essential hypertension. *Pol Arch Med Wewn*, 100: 27–34, 1998.
- EP Widmaier, H Raff, and KT Strang. *Vander's Human Physiology*. 11 edition, 2008.
- JN Wilcox, A Augustine, DV Goeddel, and DG Lowe. Differential regional expression of three natriuretic peptide receptor genes within primate tissues. *Mol Cell Biol*, 11:3454–3462, 1991.
- Williams, Ng LL, and O'Brien RJ et al. Comparison of plasma n-brain natriuretic peptide, peak oxygen consumption, and left ventricular ejection fraction for severity of chronic heart failure. *Am J Cardiol*, 93:1560–1561, 2004.
- MS Willis, ES Lee, and DG Grenache. Effect of anemia on plasma concentrations of nt-probnp. *Clin Chim Acta*, 358:175–81, 2005.
- RM Witteles, D Kao, D Christopherson, K Matsuda, RH Vagelos, D Schreiber, and MB Fowler. Impact of nesiritide on renal function in patients with acute decompensated heart failure and pre-existing renal dysfunction a randomized, double-blind, placebo-controlled clinical trial. *J Am Coll Cardiol*, 50:1835–1840, 2007.
- Witthaut, Busch, and Fraunberger. Plasma atrial natriuretic peptide and brain natriuretic peptide are increased in septic shock volume = , impact of interleukin-6 and sepsis-associated left ventricular dysfunction. *Intensive Care Med*, 29:1696–1702, 2003.
- Wiviott, de Lemos JA, and Morrow DA. Pathophysiology, prognostic significance and clinical utility of b-type natriuretic peptide in acute coronary syndromes. (Review) *Clin Chim Acta*, 346:119–128, 2004.

- KC Wold, H Vik-Mo, and T Omland. Blood haemoglobin is an independent predictor of b-type natriuretic peptide (bnp). *Clin Sci (Lond)*, 109:69–74, 2005.
- Woodard, Rosado, and Brown. Expression and control of c-type natriuretic peptide in rat vascular smooth muscle cells. *Am J Physiol Reg Int Comp Physiol*, 282:R156–165, 2002.
- Wright, Doughty RN, and Pearl A et al. Plasma amino-terminal pro-brain natriuretic peptide and accuracy of heart-failure diagnosis in primary care: a randomized, controlled trial. *J Am Coll Cardiol*, 42: 1793–1800, 2003.
- Wu and Smith A. Biological variation of the natriuretic peptides and their role in monitoring patients with heart failure. *Eur J Heart Fail*, 6:355–358, 2004.
- Wu, Smith A, and Wieczorek S et al. Biological variation for n-terminal pro- and btype natriuretic peptides and implications for therapeutic monitoring of patients with congestive heart failure. *Am J Cardiol*, 92:628–631, 2003a.
- Wu, Harrison A, and Maisel AS. Reduced readmission rate for alternating diagnoses of heart failure and pulmonary disease after implementation of b-type natriuretic peptide testing. *Eur J Heart Fail*, 6: 309–312, 2004a.
- AH Wu, T Omland, P Duc, J McCord, RM Nowak, and JE Hollander. The effect of diabetes on b-type natriuretic peptide concentrations in patients with acute dyspnea: an analysis from the breathing not properly multinational study. *Diabetes Care*, 27:2398–404, 2004b.
- AH Wu, T Omland, C Wold Knudsen, J McCord, RM Nowak, and JE Hollander. Relationship of b-type natriuretic peptide and anemia in patients with and without heart failure: A substudy from the breathing not properly (bnp) multinational study. *Am J Hematol*, 80:174–80, 2005.

- AHB Wu, A Smith, S Wiecezorek, JF Mather, B Duncan, CM White, C McGill, D Katten, and G Heller. Biological variation for n-terminal pro- and b-type natriuretic peptides and implications for therapeutic monitoring of patients with congestive heart failure. *Am J Cardiol*, 92:628–31, 2003b.
- C Wu, F Wu, J Pan, J Morser, and Q Wu. Furin-mediated processing of pro-c-type natriuretic peptide. *J Biol Chem*, 278:25847–25852, 2003c.
- Wylie, Murphy SA, and Morrow DA et al. Validated risk score predicts the development of congestive heart failure after presentation with unstable angina or non-st-elevation myocardial infarction: results from opus-timi 16 and tactics-timi 18. *Am Heart J*, 148:173–180, 2004.
- W Yan, F Wu, J Morser, and Q Wu. Corin, a transmembrane cardiac serine protease, acts as a pro-atrial natriuretic peptide-converting enzyme. *Proc Natl Acad Sci U S A*, 97:8525–8529, 2000.
- TG Yandle, AM Richards, MG Nicholls, R Cuneo, EA Espiner, and JH Livesey. Metabolic clearance rate and plasma half life of alpha-human atrial natriuretic peptide in man. *Life Sci*, 38:1827– 1833, 1986.
- TG Yandle, SO Brennan, EA Espiner, MG Nicholls, and AM Richards. Endopeptidase-24,11 in human plasma degrades atrial natriuretic factor (anf) to anf(99-105/106-126). *Peptides*, 10:891–894, 1989.
- A Yasoda, Y Komatsu, H Chusho, T Miyazawa, A Ozasa, M Miura, T Kurihara, T Rogi, S Tanaka, M Suda, N Tamura, Y Ogawa, and K Nakao. Overexpression of *cnf* in chondrocytes rescues achondroplasia through a mapk-dependent pathway. *Nat Med*, 10:80–86, 2004.
- Yasuda, Kanna, and Sakuragi. Local delivery of single low-dose of c-type natri- uretic peptide, an endogenous vascular modulator, in-

- hibits neointimal hyperplasia in a balloon-injured rabbit iliac artery model. *J Cardiovasc Pharmacol*, 39:784–788, 2002.
- Yasutake, Seino Y, and Kashiwagi M et al. Detection of cardiac sarcoidosis using cardiac markers and myocardial integrated backscatter. *Int J Cardiol*, 102:259–268, 2005.
- VT Yeung, SK Ho, MG Nicholls, and CS Cockram. Binding of cnp-22 and cnp-53 to cultured mouse astrocytes and effects on cyclic gmp. *Peptides*, 17:101–106, 1996.
- Yoshimura, Yasue H, and Tanaka H et al. Responses of plasma concentrations of a type natriuretic peptide and b type natriuretic peptide to alacepril, an angiotensin-converting enzyme inhibitor, in patients with congestive heart failure. *Br Heart J*, 72:528–533, 1994.
- M Yoshimura, H Yasue, E Morita, N Sakaino, M Jougasaki, M Kurose, M Mukoyama, Y Saito, K Nakao, and H Imura. Hemodynamic, renal, and hormonal responses to brain natriuretic peptide infusion in patients with congestive heart failure. *Circulation*, 84:1581–1588, 1991.
- Zeidel. Physiological responses to natriuretic hormones. in: Fray jcs, goodman hm (eds) handbook of physiology, section 7, the endocrine system, volume iii: Endocrine regulation of water and electrolyte balance. 2000.
- Zhang, Moalem, and Tse. Effects of natriuretic peptides on ventricular myocyte contraction and role of cyclic gmp signaling. *Eur J Pharmacol*, 510:209–215, 2005.