



جوستاف روسي انترناشيونال - مصر (EPE) * طلب علاج أول مرة / متردد

إسم المريض رباعياً : عواطف مخيمر انور احمد

المحافظة : القاهرة الحي / المدينة : المرج

العنوان : 42 ش بركة الحاج - ش زاهر المرج القاهرة

رقم البطاقة : 26708052501882 رقم التليفون : 01099808140 تاريخ تسجيل الإستمارة : 03-10-2026

العمر (تاريخ الميلاد) : 05-08-1967 المهنة : ربه منزل

الإقرار

أقر انا الموقع أدناه انني لا اتمتع باي نظام علاج مثل:- التأمين الصحي-القوات المسلحة-الشرطة-الهيئات-المؤسسات او اي نظام يتعارض مع نظام العلاج علي نفقة الدولة- وسوف اتحمل كافة نفقات العلاج في حالة ثبوت غير ذلك

الإخصائي الاجتماعي (.....) التوقيع: (.....)

المقر بما فيه (.....) التوقيع: (.....)

التشخيص الإكلينيكي : سرطان بالثدي (مبادرة)

نتائج الأبحاث : (مرفق)

التشخيص النهائي : سرطان بالثدي (مبادرة)

خطة العلاج : علاج مبادرة سرطان الثدي علاج هرموني خط ثاني مثبتات الاروماتيز [14161]

التعليق على التشخيص :

تقرير اللجنة الثلاثية المتخصصة

الاسم : د/محمد كيلاني الوظيفة : التوقيع :
الاسم : د/ندى حسن الوظيفة : التوقيع :
الاسم : د/نرمين رمضان الوظيفة : التوقيع :

أطلعت اللجنة علي لائحة العلاج الخاصة بالمجالس وتلتزم جهة العلاج بلائحة المجالس الطبية. كما تقر اللجنة بأن البيانات المدونة فى هذا النموذج صحيحة.

تعتمد صحة البيانات المرفقة رئيس اللجنة ومدير المستشفى.

تتعهد المستشفى أن لا يتم تنفيذ القرار إلا بعد مراجعة الملف الطبي للمريض وإعادة مناظرة المريض بالمستشفى قبل الصرف.

أعضاء اللجنة

..... /د. /د. /د. /د. /د. /د.

قرار اللجنة

التشخيص النهائي :
خطة العلاج :
المستشفى :
قسم :
جهة العلاج :
نفقات :
ملاحظات :

ختم
شعار الجمهورية

مدير عام
مديرية الشؤون الصحية

رئيس
المجلس الطبي المتخصص

Name: عواطف مخيمر انور احمد
National ID: 26708052501882



تقرير طبي

السيد

تحية طبية وبعد,,

بشأن تقرير طبي للمريض

إيماء للطلب المقدم من

رقم طبي:

بقسم:

نتشرف بأن نوضح فيما يلي تقريراً عن حالته

تاريخ وساعة الخروج :
PM 01:17 05-07-2026

تاريخ وساعة الدخول :
-- --:-- yyyy-mm-dd

وبعد الكشف والفحوصات والمتابعة وجد أنه

المریضة تعاني من ورم غير حميد بالثدي و تحتاج الي Second Line Hormonal Treatment Aromatase Inhibitor

الوقت :

-- --:--

التاريخ :

05-07-2026

الطبيب المعالج :
د/شيماء لاشين

جمهورية مصر العربية
بطاقة تحقيق الشخصية



عواطف
مخيمر انور احمد
٤٢ ش بركة الحاج - ش زاهر
المرج - القاهرة

٢٠١٧/١٧/٥٥ ٢٠١٧.٠٨.٥٢٥.١٨٨٢
J27004766

٢٠٢٣/٠٥/ ٢٠١٧.٠٨.٥٢٥.١٨٨٢



ربة منزل
انثى
مسلمة
متزوجة
سعد محمد عبدالراضي محمد
البطاقة مارية حتى ٢٠٣٠/٠٥/١٤





مبادرة رئيس الجمهورية
لدعم صحة المرأة المصرية

الاستشارة
معدة
مصر

MDT FORM

15/5/2022

Name	علاء كهنه انور	MRN	2155 / 0.7
Age	54 yr	ID	271.0.000.1A1e
Menopausal status		Phone	0109980114
OCP		marital status	متزوج
Co morbidity	Htnsion. , (1) Amenorrhea 1yr.	Family history	

Clinical Data			
Bx:	Rt- IDC grade II		
ER:	PR:	HER 2:	Ki67:
6/8	0/8	-ve	15%
Sonomammography:			
Post simple mastectomy x SLNB upfront surgery			
vets Workup:			
R L			
TNM			

Recommendations: Kt mastectomy + SLNBx 12/4/2022
IDC, GII
free vis. T3N0

MDT OUTCOME & PLAN

- Adjuvant AC X3 → Taxol x9 weeks
- PORT
- Adjuvant HT AI X5 yrs. after E2

SIGNATURE			
MEDICAL ONCOLOGY	SURGERY	RADIOLOGY	PATHOLOGY
د. محمد عبد الله		د. محمد عبد الله	✓

د. محمد عبد الله

Name: عواطف مخيمر انور احمد		
MR. : 3843		
Detailed Medical Report		
		السيد
تحتية طبية وبعد,,		
	بشأن تقرير طبي للمريض	إيماء للطلب المقدم من
	رقم طبي:	بقسم:
ننشرف بأن نوضح فيما يلي تقريراً عن حالته		
	تاريخ وساعة الخروج :	تاريخ وساعة الدخول :
وبعد الكشف والفحوصات والمتابعة وجد انه		
The patient is a 58-year-old female with a history of right breast cancer diagnosed in April 2022. Pathology was estrogen-receptor positive (ER 6/8), progesterone-receptor negative (PR 0/8), HER2-negative, staged T3 N0. She underwent a right mastectomy followed by adjuvant chemotherapy and completed a three-year course of tamoxifen. The treatment plan now calls for a transition to anastrozole (Arimidex) for a three-month duration, pending pharmacy processing. The patient reports severe bony pain and has been prescribed anastrozole 1 mg daily (1 × 1 × 90) with supportive measures. A dental assessment has been requested, and the patient is scheduled for follow-up in three months. A bone scan performed in August 2026 demonstrated extensive osteoblastic metastatic lesions. The patient denies axial skeletal pain on examination. Imaging with contrast for CTCAP was obtained, and the patient continues on anastrozole with supportive care for an additional three months.		
تحريراً في ,		
	الوقت :	التاريخ : 20-09-2026
	الطبيب المعالج : علياء احمد أبو النصر شهاب الدين	
يعتمد ,		

Request No : 13005027
Request Date : 03/04/2022
Verification Date : 12/04/2022
Financial Category : أروام دار السلام - ميانرو
Department : Out Patient
Registered by : marwa mohamed youss

Clinical Data:
Rt breast lesion

Rt simple mastectomy specimen & SLN biopsy

Rt simple mastectomy specimen measured 21 x 21 x 5 cm, with covering skin ellipse measured 18 x 12.5 cm.

- Microscopic examination:

- Diagnosis:

- Invasive duct carcinoma grade II
- Free nipple and areola

Printing Date : 20/04/2022 12:41:07

Printing By : marwa mohamed youss

Patient Name : السيدة / عواطف مكيور انور
Patient No. : 26708052501882
Path No. : 3045/21
Sex : Female
Age : 54 Y
Referring Doc. :

Request No. : 131537
Request Date : 09/12/2021
Verification Date : 23/12/2021
Financial Category : اورام دار السلام - مياطرة
Department : Out Patient
IHC No. :
Registered by : marwa mohamed yousse

IMMUNO HISTOCHEMISTRY REPORT

Clinical Data:

1) breast mass

Nature of the specimen:

tru cut biopsy

Gross Examination:

Multiple gray white tissue cores up to 1.5cm, totally submitted. Sections prepared from paraffin block or tissue staining using Bench Mark ULTRA system for Efr, PR,HER2, ER clone (1E2). PR clone SP9, HER-2 clone (4B5), & Ki67 clone (30-9)

Histoscopic examination:

Histologic type: IDC

Grade:

Angioinvasion: no

Ductal carcinoma in situ: no

Diagnosis:

1) breast mass, tru cut biopsy:

Invasive duct carcinoma GII

Immunomarkers:

R: Strong nuclear reaction in 35% of tumor cells (Allred score 6/8)

R: Negative nuclear reaction in tumor cells (Allred score 0/8)

Her-2: Negative membrane reaction in tumor cells (score 0)

Ki67: 15 % nuclear reaction

Prof.Gamal Abrie!Aty
Dr.Rana Fathy

Verified by:

Rana fathy

Printing Date : 09/02/2022 09:08:07

Page 1 of 2

Printed By : A/ mostafa

Patient Name : د. عروفت مجهر عروفت	Request No : 13005027
Patient No : 08779000000000000000	Request Date : 03/04/2022
Path No. : 920/22	Verification Date : 12/04/2022
Sex : Female	Financial Category : مبادرة - عيادة
Age : 54 Y	Department : Out Patient
Referring Doc :	Registered by : marwa mohamed youss

Pathology Report

- Free deep margin
- LN: Free SLNs (0/7)
- Stage : pT3 ,pN0

Dr.Menna Atef

Prof.Lubna Omar

Dr.

Verified by
dr.esraa magdy


Printing Date : 20/04/2022 12:55:07

Page 2 of 2

Printing By : marwa mohamed youss



Name : *Awatef Mekki* Date : *26-02-2022*

POST CONTRAST MULTI-SLICE CT SCAN OF THE CHEST, ABDOMEN & PELVIS REVEALED: -

1. Chest:

- Right breast irregular speculated mass lesions the largest measuring 4 X 3 cm.
- No sizable pulmonary nodules.
- No pathological hilar or mediastinal lymph nodes.
- No evidence of pleural effusion, pleural thickening or masses seen.
- Average cardiac size. No pericardial effusion detected.
- Normal CT appearance of the thoracic aorta and the great vessels.

2. Abdomen and pelvis:

- The liver is enlarged with homogenous CT texture. No definite focal lesions or contour irregularities.
- Patent homogeneously enhancing portal vein and its main branches as well as its main tributaries.
- No intra-hepatic or extra-hepatic biliary radicles dilatation.
- Normal size & parenchymal texture of the spleen with no focal lesions seen.
- No pathological abdominal or pelvic lymph nodes.
- Normal size, parenchymal thickness & density of both kidneys with no stones or back-pressure changes.
- Normal appearance of the pancreas, aorta, I.V.C and both supra-renals.
- Smooth filling of the urinary bladder with no filling defects or diverticular outpouching.
- No abdominal or pelvic masses or collections.

3. The scanned skeleton:

- Left sacral ala sclerotic focus mostly bone island.

OPINION:

- *Right breast malignant looking mass lesions.*
- *No definite pulmonary, hepatic or osseous metastatic deposits.*

Dr. Mohammed Mohsen, MSc

Dr. Amir Farouk MD

٣٨٤٣



مركز النيل للأشعة والتحاليل

SCANS LABS

Name : Awatif Mikhaymir Anwar Ahmed

Date : 31-07-2026 06:19

Branch : Dokki

Medical No. : 428975

Ref.Physician Prof : Goustav Roussey Egypt

Ins.Company : معمل مستشفيات

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

- Faint visualization of both kidneys.

Impression:

- Extensive widespread osteoblastic metastatic skeletal disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

Dr. Alaa Nagi,
MD Nuclear Medicine

R E C E N T E R C E N T R E D I A G N O S T I C N I L E

Call Center 19656

Date : 31-07-2026 06:19

Medical No.: 428975

تعاقد استشارات : Ins. Company :

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

Impression:

- Extensive widespread osteoblastic metastatic disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

**Dr. Alaa Nagi,
MD Nuclear Medicine**

٣٨٤٢



مركز النيل للأشعة والتحاليل

SCANS LABS

Name : Awatif Mikhaymir Anwar Ahmed

Date : 31-07-2026 06:19

Branch : Dokki

Medical No. : 428975

Ref.Physician Prof : Goustav Roussey Egypt

Ins.Company : معمل مستشفيات

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

- Faint visualization of both kidneys.

Impression:

- Extensive widespread osteoblastic metastatic skeletal disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

Dr. Alaa Nagi,
MD Nuclear Medicine

R E C E N T E R C E N T R E D I A G N O S T I C N I L E

٣٨٤٢



مركز النيل للأشعة والتحاليل

SCANS LABS

Name : Awatif Mikhaymir Anwar Ahmed

Date : 31-07-2026 06:19

Branch : Dokki

Medical No. : 428975

Ref.Physician Prof : Goustav Roussey Egypt

Ins.Company : معمل مستشفيات

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

- Faint visualization of both kidneys.

Impression:

- Extensive widespread osteoblastic metastatic skeletal disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

Dr. Alaa Nagi,
MD Nuclear Medicine

R E C E N T E R C E N T R E D I A G N O S T I C N I L E

٣٨٤٣



مركز النيل للأشعة والتحاليل

SCANS LABS

Name : Awatif Mikhaymir Anwar Ahmed

Date : 31-07-2026 06:19

Branch : Dokki

Medical No. : 428975

Ref.Physician Prof : Goustav Roussey Egypt

Ins.Company : معمل مستشفيات

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

- Faint visualization of both kidneys.

Impression:

- Extensive widespread osteoblastic metastatic skeletal disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

Dr. Alaa Nagi,
MD Nuclear Medicine

R E C E N T E R C E N T R E D I A G N O S T I C N I L E

٣٨٤٣



مركز النيل للأشعة و التحاليل

SCANS LABS

Name : Awatif Mikhaymir Anwar Ahmed

Date : 31-07-2026 06:19

Branch : Dokki

Medical No. : 428975

Ref.Physician Prof : Goustav Roussey Egypt

Ins.Company : معمل مستشفيات

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

- Faint visualization of both kidneys.

Impression:

- Extensive widespread osteoblastic metastatic skeletal disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

Dr. Alaa Nagi,
MD Nuclear Medicine

R E C E N T E R C E N T R E D I A G N O S T I C N I L E

٣٨٤٣



مركز النيل للأشعة و التحاليل

SCANS LABS

Name : Awatif Mikhaymir Anwar Ahmed

Date : 31-07-2026 06:19

Branch : Dokki

Medical No. : 428975

Ref.Physician Prof : Goustav Roussey Egypt

Ins.Company : معمل مستشفيات

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

- Faint visualization of both kidneys.

Impression:

- Extensive widespread osteoblastic metastatic skeletal disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

Dr. Alaa Nagi,
MD Nuclear Medicine

R E C E N T E R C E N T R E D I A G N O S T I C N I L E

٣٨٤٣



مركز النيل للأشعة و التحاليل

SCANS LABS

Name : Awatif Mikhaymir Anwar Ahmed

Date : 31-07-2026 06:19

Branch : Dokki

Medical No. : 428975

Ref.Physician Prof : Goustav Roussey Egypt

Ins.Company : معمل مستشفيات

Whole-Body Bone Scintigraphy Report

Clinical History: 58-year-old female with history of right breast cancer status post-surgical resection, chemotherapy, radiotherapy, and currently on hormonal therapy. Presenting with generalized bony aches.

Radiopharmaceutical: Tc-99m MDP

Procedure: Whole-body anterior and posterior planar scintigraphy

Findings:

- Diffuse extensive focal and confluent areas of markedly increased tracer uptake involve both the axial and appendicular skeleton. Prominent widespread involvement is demonstrated throughout the skull, cervical, thoracic and lumbar spine, ribs, sternum, pelvis, and bilateral long bones. Background soft-tissue clearance is reduced secondary to high osseous radiopharmaceutical avidity.

- Faint visualization of both kidneys.

Impression:

- Extensive widespread osteoblastic metastatic skeletal disease with heavy infiltration throughout the axial and appendicular skeleton.
- Follow up is recommended.

Dr. Alaa Nagi,
MD Nuclear Medicine

R E C E N T E R C E N T R E D I A G N O S T I C N I L E