جوستاف روسي انترناشيونال - مصر (EPE) * **طلب علاج أول مرة / متردد**

إسم المريض رباعياً : منى جابر محمود العاصى
المحافظة : القاهره **الحي / المدينة :** البساتين **العنوان :** ..
رقم البطاقة : 27501011708863 **رقم التليفون :** 01288415585 **تاريخ تسجيل الإستمارة :** 20-10-2026
العمر(تاريخ الميلاد) : 01-01-1975 **المهنة :** ربه منزل

الإقرار

أقر انا الموقع أدناه انني لا اتمتع باي نظام علاج مثل:-التأمين الصحي-القوات المسلحه-الشرطة-الهيئات-المؤسسات او اي نظام يتعارض مع نظام العلاج علي نفقة الدولة- وسوف اتحمل كافة نفقات العلاج في حالة ثبوت غير ذلك

الإخصائي الاجتماعي **المقر بما فيه**
التوقيع:(----------) **التوقيع:**(----------)

التشخيص الإكلينيكي : سرطان بالثدي (مبادرة)
نتائج الأبحاث : (مرفق)
التشخيص النهائي : سرطان بالثدي (مبادرة)
خطة العلاج : أبحاث أورام مسح ذري علي العظام [12966] **قسم العلاج :** الاورام
التعليق على التشخيص : ----------

تقرير اللجنة الثلاثية المتخصصة

الاسم: د/محمد كيلاني **الوظيفة:** ---------- **التوقيع:** ----------
الاسم: د/ندى حسن **الوظيفة:** ---------- **التوقيع:** ----------
الاسم: د/نرمين رمضان **الوظيفة:** ---------- **التوقيع:** ----------

أطلعت اللجنة علي لائحة العلاج الخاصة بالمجالس وتلتزم جهة العلاج بلائحة المجالس الطبية.
كما تقر اللجنة بأن البيانات المدونة فى هذا النموذج صحيحة.

تعتمد صحة البيانات المرفقة
رئيس اللجنة و مدير المستشفي.

تتعهد المستشفى أن لا يتم تنفيذ القرار إلا بعد مراجعة الملف الطبي للمريض وإعادة مناظرة المريض بالمستشفي قبل الصرف.

أعضاء اللجنة

ا.د/ ---------- **ا.د/** ---------- **د/** ---------- **د/** ----------

قرار اللجنة

التشخيص النهائي : ----------
خطة العلاج : ----------
المستشفي : ----------**قسم :** ----------
جهة العلاج : ----------**نفقات :** ----------
ملاحظات : ----------

رئيس	مدير عام	ختم
المجلس الطبي المتخصص	مديرية الشئون الصحية	شعار الجمهورية

Name: منى جابر محمود العاصي
National ID: 27501011708863

تقرير طبي

السيد

تحية طيبة وبعد,,

إيماء للطلب المقدم من	**بشأن تقرير طبي للمريض**
بقسم:	**رقم طبي:**

نتشرف بأن نوضح فيما يلي تقريرا عن حالته

تاريخ وساعة الدخول : -- --:-- yyyy-mm-dd	**تاريخ وساعة الخروج :** PM 01:17 05-07-2026

وبعد الكشف والفحوصات والمتابعة وجد انة

المريضة تعاني من ورم غير حميد بالثدى و تحتاج الي **Bone scintigraphy**

التاريخ : **05-07-2026**	**الوقت :** **-- --:--**
	الطبيب المعالج : د/شيماء لاشين



جمهورية مصر العربية
بطاقة تحقيق الشخصية

منى
جابر محمود العاصى
٦ ش جامع حسنين-ش مهران
البساتين - القاهره

١٩٧٥/٠١/٠١

٢٧٥٠١٠١١٧٠٨٨٦٣

LA6874169

٠١٢٨٨٤١٠٠٨٥



٢٧٥٠١٠١١٧٠٨٨٦٣ ٢٠٢٥/٠٣

ربة منزل

أنثى مسلمة متزوجة
اشرف سيد اسماعيل احمد
البطاقة سارية حتى ٢٠٣٢/٠٢/٢٨

١

National ID – Front & Back
[[DOCTYPE:ID_FRONT_BACK]]

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- 1**2 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY

Mammogram
[[DOCTYPE:MAMMOGRAM]]


جمهورية مصر العربية
بطاقة تحقيق الشخصية
منى
جابر محمود العاصى
٦ ش جامع حسنين-ش مهران
البساتين - القاهره
٢٧٥٠١٠١١٧٠٨٨٦٣
LA6874169


٢٧٥٠١٠١١٧٠٨٨٦٣ ٢٠٢٥/٠٣
ربة منزل
انثى مسلمة متزوجة
اشرف سيد اسماعيل احمد
١ البطاقة سارية حتى ٢٠٣٢/٠٢/٢٨
National ID - Back
[[DOCTYPE/ID_BACK]]

[[DOCTYPE:MAMMOGRAM]]
Mammogram

جوستاف روسي

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- **12 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International Egypt Pathology Lab
جوستاف روسي انترناشونال – مصر معمل الباثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	357584
Sex/Age	Female 51 Y	**Patient No.**	25778
Registered Date	12/8/2026 09:43:53 AM	**Referred By Dr.**	Interventional Radiology
Validation Date	20/9/2026 02:45:34 PM	**Branch**	Gustave Roussy
Reporting Date	24/9/2026 09:08:46 AM	**Pathology No.**	2548/26

Pathology Report

Clinical Data: Left breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 12/08/2026. Cold ischemia time is 20 seconds.

Gross Examination:
- Multiple fragmented grey, white tissue cores, ranging in size between 0.5 cm & 1.0 cm long, totally submitted.

Technique:
- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for;
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), SISH, & Ki67 clone (30-9).

Microscopic examination:
- **Histologic type:** Invasive lobular carcinoma.
- **Histologic grade:** 2 (differentiation 3, pleomorphism 2, mitosis 1).
- **DCIS:** Absent.
- **Microcalcification:** Absent.
- **Angioinvasion:** Absent.
- **Perineural invasion:** Absent.
- **Tumor infiltrating lymphocytes (TILs):** 1%.

Results:
* IPT (All controls showed appropriate reactivity):
- ER: Strong nuclear reaction in 95% of tumor cells (score 8/8).
- PR: Strong nuclear reaction in 99% of tumor cells (score 8/8).
- Her2: Moderate complete membrane reaction in more than 10% of tumor cells (score 2+).
- SISH: Amplified with clusters.
(Average Her2 copy No.: (6), Her2/CEP17 ratio: (4))
- Ki67%: 20%.

Conclusion:
Left breast mass, Tru-cut biopsy:
- Invasive lobular carcinoma, Grade 2.
IPT:

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International-Egypt- Pathology Lab
جوستاف روسي انترناشونال – مصر معمل الباثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	**357584**
Sex/Age	**Female 51 Y**	**Patient No.**	**25778**
Registered Date	**12/8/2026 09:43:53 AM**	**Referred By Dr.**	**Interventional Radiology**
Validation Date	**20/9/2026 02:45:34 PM**	**Branch**	**Gustave Roussy**
Reporting Date	**24/9/2026 09:08:46 AM**	**Pathology No.**	**2548/26**

- ER: Positive (score 8/8).
- PR: Positive (score 8/8).
- Her2: Equivocal (score 2+).
- SISH: Amplified with clusters.
- Ki67%: 20%.


***Scoring system: "ER, PR"*
ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66%
and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, week, 2, intermediate; and 3, strong). Total score from 0 to 8.
*** Scoring system: "Her2"*
Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.
IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within >10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.
ISH groups:
The ratio is calculated and the results is interpreted as:
• Group I (Positive): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is more than 4.
• Group II (Negative): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is less than 4. (Recounting give the same result).
• Group III (Positive): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 6 (Recounting give the same result).
• Group IV (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 4 but less than 6. (Recounting give the same result).
• Group V (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is less than 4.

Signature
Prof. Dr. Maha Guimei, MD, PhD.
Dr. Walaa Ibrahim Ghoneim. MD.


[[DOCTYPE:OTHER_PRESENTIAL_INITIATIVE_CARD]]
presential-initiative-card

Advanced Clinical Exam

الثدي الأيمن



الثدي الأيسر



- **Maximum mass dimension (mm):**
- **Side:**
  - ○Right
  - ○Left
  - ○Bilateral
- **Qiadrant:**
  - ○UOQ
  - ○UIQ
  - ○LOQ
  - ○LIQ
  - ○Central and retroareolar
  - ○Axilla
- **Imaging study requested:**
  - ○Sonography
  - ○Mammosonography
  - ○MRI breasts

[[DOCTYPE:MAMMOGRAM]]
Mammogram

جوستاف روسي
انترناشونال

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- **12 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY


جمهورية مصر العربية
بطاقة تحقيق الشخصية
منى
جابر محمود العاصى
٦ ش جامع حسنين-ش مهران
البساتين - القاهره
٢٧٥٠١٠١١٧٠٨٨٦٣
LA6874169


٢٧٥٠١٠١١٧٠٨٨٦٣
٢٠٢٥/٠٣
ربة منزل
انثى
مسلمة
متزوجة
اشرف سيد اسماعيل احمد
١
البطاقة سارية حتى ٢٠٣٢/٠٢/٢٨

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- **12 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International Egypt Pathology Lab
جوستاف روسي انترناشونال – مصر معمل البثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	357584
Sex/Age	Female 51 Y	**Patient No.**	25778
Registered Date	12/8/2026 09:43:53 AM	**Referred By Dr.**	Interventional Radiology
Validation Date	20/9/2026 02:45:34 PM	**Branch**	Gustave Roussy
Reporting Date	24/9/2026 09:08:46 AM	**Pathology No.**	2548/26

Pathology Report

Clinical Data: Left breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 12/08/2026. Cold ischemia time is 20 seconds.

Gross Examination:
- Multiple fragmented grey, white tissue cores, ranging in size between 0.5 cm & 1.0 cm long, totally submitted.

Technique:
- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for;
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), SISH, & Ki67 clone (30-9).

Microscopic examination:
- **Histologic type:** Invasive lobular carcinoma.
- **Histologic grade:** 2 (differentiation 3, pleomorphism 2, mitosis 1).
- **DCIS:** Absent.
- **Microcalcification:** Absent.
- **Angioinvasion:** Absent.
- **Perineural invasion:** Absent.
- **Tumor infiltrating lymphocytes (TILs):** 1%.

Results:
* IPT (All controls showed appropriate reactivity):
- ER: Strong nuclear reaction in 95% of tumor cells (score 8/8).
- PR: Strong nuclear reaction in 99% of tumor cells (score 8/8).
- Her2: Moderate complete membrane reaction in more than 10% of tumor cells (score 2+).
- SISH: Amplified with clusters.
(Average Her2 copy No.: (6), Her2/CEP17 ratio: (4))
- Ki67%: 20%.

Conclusion:
Left breast mass, Tru-cut biopsy:
- Invasive lobular carcinoma, Grade 2.
IPT:

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International-Egypt- Pathology Lab
جوستاف روسي انترناشونال – مصر معمل الباثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	**357584**
Sex/Age	**Female 51 Y**	**Patient No.**	**25778**
Registered Date	**12/8/2026 09:43:53 AM**	**Referred By Dr.**	**Interventional Radiology**
Validation Date	**20/9/2026 02:45:34 PM**	**Branch**	**Gustave Roussy**
Reporting Date	**24/9/2026 09:08:46 AM**	**Pathology No.**	**2548/26**


- ER: Positive (score 8/8).
- PR: Positive (score 8/8).
- Her2: Equivocal (score 2+).
- SISH: Amplified with clusters.
- Ki67%: 20%.

***Scoring system: "ER, PR"*
ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66%
and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, week, 2, intermediate; and 3, strong). Total score from 0 to 8.
*** Scoring system: "Her2"*
Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.
IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within >10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.
ISH groups:
The ratio is calculated and the results is interpreted as:
• Group I (Positive): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is more than 4.
• Group II (Negative): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is less than 4. (Recounting give the same result).
• Group III (Positive): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 6 (Recounting give the same result).
• Group IV (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 4 but less than 6. (Recounting give the same result).
• Group V (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is less than 4.


Signature
Prof. Dr. Maha Guimei, MD, PhD.
Dr. Walaa Ibrahim Ghoneim. MD.

Advanced Clinical Exam

الثدي الأيسر



الثدي الأيمن



- **Maximum mass dimension (mm):**
- **Side:**
  - ○Right
  - ○Left
  - ○Bilateral
- **Qiadrant:**
  - ○UOQ
  - ○UIQ
  - ○LOQ
  - ○LIQ
  - ○Central and retroareolar
  - ○Axilla
- **Imaging study requested:**
  - ○Sonography
  - ○Mammosonography
  - ○MRI breasts

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- **12 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY


جمهورية مصر العربية
بطاقة تحقيق الشخصية
منى
جابر محمود العاصى
٦ ش جامع حسنين-ش مهران
البساتين - القاهره
٢٧٥٠١٠١١٧٠٨٨٦٣
LA6874169


٢٧٥٠١٠١١٧٠٨٨٦٣
٢٠٢٥/٠٣
ربة منزل
انثى
مسلمة
متزوجة
اشرف سيد اسماعيل احمد
١
البطاقة سارية حتى ٢٠٣٢/٠٢/٢٨

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- **12 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International Egypt Pathology Lab
جوستاف روسي انترناشونال – مصر معمل البثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	357584
Sex/Age	Female 51 Y	**Patient No.**	25778
Registered Date	12/8/2026 09:43:53 AM	**Referred By Dr.**	Interventional Radiology
Validation Date	20/9/2026 02:45:34 PM	**Branch**	Gustave Roussy
Reporting Date	24/9/2026 09:08:46 AM	**Pathology No.**	2548/26

Pathology Report

Clinical Data: Left breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 12/08/2026. Cold ischemia time is 20 seconds.

Gross Examination:
- Multiple fragmented grey, white tissue cores, ranging in size between 0.5 cm & 1.0 cm long, totally submitted.

Technique:
- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for;
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), SISH, & Ki67 clone (30-9).

Microscopic examination:
- **Histologic type:** Invasive lobular carcinoma.
- **Histologic grade:** 2 (differentiation 3, pleomorphism 2, mitosis 1).
- **DCIS:** Absent.
- **Microcalcification:** Absent.
- **Angioinvasion:** Absent.
- **Perineural invasion:** Absent.
- **Tumor infiltrating lymphocytes (TILs):** 1%.

Results:
* IPT (All controls showed appropriate reactivity):
- ER: Strong nuclear reaction in 95% of tumor cells (score 8/8).
- PR: Strong nuclear reaction in 99% of tumor cells (score 8/8).
- Her2: Moderate complete membrane reaction in more than 10% of tumor cells (score 2+).
- SISH: Amplified with clusters.
(Average Her2 copy No.: (6), Her2/CEP17 ratio: (4))
- Ki67%: 20%.

Conclusion:
Left breast mass, Tru-cut biopsy:
- Invasive lobular carcinoma, Grade 2.
IPT:

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International-Egypt- Pathology Lab
جوستاف روسي انترناشونال – مصر معمل الباثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	**357584**
Sex/Age	**Female 51 Y**	**Patient No.**	**25778**
Registered Date	**12/8/2026 09:43:53 AM**	**Referred By Dr.**	**Interventional Radiology**
Validation Date	**20/9/2026 02:45:34 PM**	**Branch**	**Gustave Roussy**
Reporting Date	**24/9/2026 09:08:46 AM**	**Pathology No.**	**2548/26**


- ER: Positive (score 8/8).
- PR: Positive (score 8/8).
- Her2: Equivocal (score 2+).
- SISH: Amplified with clusters.
- Ki67%: 20%.

***Scoring system: "ER, PR"*
ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66%
and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, week, 2, intermediate; and 3, strong). Total score from 0 to 8.
*** Scoring system: "Her2"*
Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.
IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within >10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.
ISH groups:
The ratio is calculated and the results is interpreted as:
• Group I (Positive): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is more than 4.
• Group II (Negative): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is less than 4. (Recounting give the same result).
• Group III (Positive): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 6 (Recounting give the same result).
• Group IV (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 4 but less than 6. (Recounting give the same result).
• Group V (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is less than 4.


Signature
Prof. Dr. Maha Guimei, MD, PhD.
Dr. Walaa Ibrahim Ghoneim. MD.

Advanced Clinical Exam

الثدي الأيسر



الثدي الأيمن



- **Maximum mass dimension (mm):**
- **Side:**
  - ○Right
  - ○Left
  - ○Bilateral
- **Qiadrant:**
  - ○UOQ
  - ○UIQ
  - ○LOQ
  - ○LIQ
  - ○Central and retroareolar
  - ○Axilla
- **Imaging study requested:**
  - ○Sonography
  - ○Mammosonography
  - ○MRI breasts

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- **12 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY


جمهورية مصر العربية
بطاقة تحقيق الشخصية
منى
جابر محمود العاصى
٦ ش جامع حسنين-ش مهران
البساتين - القاهره
٢٧٥٠١٠١١٧٠٨٨٦٣
LA6874169


٢٧٥٠١٠١١٧٠٨٨٦٣ ٢٠٢٥/٠٣
ربة منزل
انثى مسلمة متزوجة
اشرف سيد اسماعيل احمد
١ البطاقة سارية حتى ٢٠٣٢/٠٢/٢٨
National ID - Back
[[DOCTYPE/ID_BACK]]

[[DOCTYPE:MAMMOGRAM]]
Mammogram

جوستاف روسي

Patient's Name:	**Mona Jaber Mahmoud Al-Asi**
Date:	**2026-07-29**
Referred by:	Prof.Dr. **no doctor**
Hospital:	**Goustave Roussey International Egypt**

Bilateral Digital soft tissue mammography with high resolution breast sonography revealed:

Clinical summary: Left retroareolar mass and nipple retraction..

Findings:

Right breast.

- Scattered densities of glandular tissue (ACR b)
- No speculations, masses asymmetry , distortion or suspicious microcalcifications.
- **Ultrasound** demonstrated homogeneous glandular appearance yet no solid or cystic lesions.
- **12 o'clock clusterd mircocyst measuring (12 x 4 mm).**
- Normal skin/nipple/areola complex.
- Non-specific axillary LNs.

Left breast.

- Scattered densities of glandular tissue (ACR b)
- **Retroareolar irregular speculated high density mass lesion associated with nipple retraction.**
- **Ultrasound demonstrated retroareolar irregular mass assocaited with intraductal extension measuring (1.5 x 1.6 x 2.4 cm) .**
- Normal skin.
- Non specfic axillary LNs .

Conclusion:-

- ***Right breast (BIRADS 2): clustered microcyst .***
- ***Left breast (BIRADS 5): Retroareolar suspicious mass.***

For biopsy .

Prof.MONA ZAKY

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International Egypt Pathology Lab
جوستاف روسي انترناشونال – مصر معمل الباثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	357584
Sex/Age	Female 51 Y	**Patient No.**	25778
Registered Date	12/8/2026 09:43:53 AM	**Referred By Dr.**	Interventional Radiology
Validation Date	20/9/2026 02:45:34 PM	**Branch**	Gustave Roussy
Reporting Date	24/9/2026 09:08:46 AM	**Pathology No.**	2548/26

Pathology Report

Clinical Data: Left breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 12/08/2026. Cold ischemia time is 20 seconds.

Gross Examination:
- Multiple fragmented grey, white tissue cores, ranging in size between 0.5 cm & 1.0 cm long, totally submitted.

Technique:
- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for;
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), SISH, & Ki67 clone (30-9).

Microscopic examination:
- **Histologic type:** Invasive lobular carcinoma.
- **Histologic grade:** 2 (differentiation 3, pleomorphism 2, mitosis 1).
- **DCIS:** Absent.
- **Microcalcification:** Absent.
- **Angioinvasion:** Absent.
- **Perineural invasion:** Absent.
- **Tumor infiltrating lymphocytes (TILs):** 1%.

Results:
* IPT (All controls showed appropriate reactivity):
- ER: Strong nuclear reaction in 95% of tumor cells (score 8/8).
- PR: Strong nuclear reaction in 99% of tumor cells (score 8/8).
- Her2: Moderate complete membrane reaction in more than 10% of tumor cells (score 2+).
- SISH: Amplified with clusters.
(Average Her2 copy No.: (6), Her2/CEP17 ratio: (4))
- Ki67%: 20%.

Conclusion:
Left breast mass, Tru-cut biopsy:
- Invasive lobular carcinoma, Grade 2.
IPT:

GUSTAVE ROUSSY INTERNATIONAL
Gustave Roussy International-Egypt- Pathology Lab
جوستاف روسي انترناشونال – مصر معمل الباثولوجي
جوستاف روسي

Patient Name	منى جابر محمود	**Case No.**	**357584**
Sex/Age	**Female 51 Y**	**Patient No.**	**25778**
Registered Date	**12/8/2026 09:43:53 AM**	**Referred By Dr.**	**Interventional Radiology**
Validation Date	**20/9/2026 02:45:34 PM**	**Branch**	**Gustave Roussy**
Reporting Date	**24/9/2026 09:08:46 AM**	**Pathology No.**	**2548/26**

- ER: Positive (score 8/8).
- PR: Positive (score 8/8).
- Her2: Equivocal (score 2+).
- SISH: Amplified with clusters.
- Ki67%: 20%.

***Scoring system: "ER, PR"*
ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66%
and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, week, 2, intermediate; and 3, strong). Total score from 0 to 8.
*** Scoring system: "Her2"*
Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.
IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within >10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.
ISH groups:
The ratio is calculated and the results is interpreted as:
• Group I (Positive): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is more than 4.
• Group II (Negative): HER2/CEP17 ratio is more than 2 & Average HER2 copy number/ cell is less than 4. (Recounting give the same result).
• Group III (Positive): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 6 (Recounting give the same result).
• Group IV (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is more than 4 but less than 6. (Recounting give the same result).
• Group V (Negative): HER2/CEP17 ratio is less than 2 & Average HER2 copy number/ cell is less than 4.

Signature
Prof. Dr. Maha Guimei, MD, PhD.
Dr. Walaa Ibrahim Ghoneim. MD.