



جوستاف روسي انترناشيونال - مصر (EPE) * طلب علاج أول مرة / متردد

إسم المريض رباعياً : سهير محمد حسين احمد
المحافظة : القاهرة
الحى / المدينة : الازهر
العنوان : ~
رقم البطاقة : 27312222101623 رقم التليفون : 01142132781
العمر (تاريخ الميلاد) : 22-12-1973 المهنة : بدون عمل
تاريخ تسجيل الإستمارة : 26-09-2026

الإقرار

أقر انا الموقع أدناه انني لا اتمتع باي نظام علاج مثل :-التأمين الصحي-القوات المسلحة-الشرطة-الهيئات-المؤسسات او اي نظام يتعارض مع نظام العلاج علي نفقة الدولة- وسوف اتحمل كافة نفقات العلاج في حالة ثبوت غير ذلك
الإخصائي الاجتماعي
التوقيع: (-----)
المقرر بما فيه
التوقيع: (-----)

التشخيص الإكلينيكي : سرطان بالثدي (مبادرة)

نتائج الأبحاث : (مرفق)

التشخيص النهائي : سرطان بالثدي (مبادرة)

خطة العلاج : أبحاث أورام اشعه مقطعيه علي البطن والصدر والحوض
بالصبغة [11127]

التعليق على التشخيص : -----

تقرير اللجنة الثلاثية المتخصصة

الاسم : د/محمد كيلاني
الاسم : د/ندى حسن
الاسم : د/نرمين رمضان
الوظيفة :
الوظيفة :
الوظيفة :
التوقيع :
التوقيع :
التوقيع :
أطلعت اللجنة علي لائحة العلاج الخاصة بالمجالس وتلتزم جهة العلاج بلائحة المجالس الطبية. كما تقرر اللجنة بأن البيانات المدونة فى هذا النموذج صحيحة.
تعتمد صحة البيانات المرفقة
رئيس اللجنة ومدير
المستشفى.

تتعهد المستشفى أن لا يتم تنفيذ القرار إلا بعد مراجعة الملف الطبي للمريض وإعادة مناظرة المريض بالمستشفى قبل الصرف.

أعضاء اللجنة

أ.د. / د. / د. / د. / د. / د. / د. / د. / د. / د.

قرار اللجنة

التشخيص النهائي :
خطة العلاج :
المستشفى :
قسم :
جهة العلاج :
نفقات :
ملاحظات :

ختم
شعار الجمهورية

مدير عام
مديرية الشؤون الصحية

رئيس
المجلس الطبي المتخصص

--

Name: سهير محمد حسين احمد National ID: 27312222101623	
--	---

تقرير طبي

السيد
تحية طبية وبعد,,

إيماء للطلب المقدم من	بشأن تقرير طبي للمريض
بقسم:	رقم طبي:

نتشرف بان نوضح فيما يلي تقريرنا عن حالته
--

تاريخ وساعة الدخول : -- --:-- yyyy-mm-dd	تاريخ وساعة الخروج : PM 01:17 05-07-2026
---	---

وبعد الكشف والفحوصات والمتابعة وجد انه
--

المریضة تعاني من ورم غير حميد بالثدى و تحتاج الي Contrast-enhanced CT Chest & abdomen and pelvis
--

التاريخ : 05-07-2026	الوقت : -- --:--
	الطبيب المعالج : د/شيماء لاشين



جمهورية مصر العربية
بطاقة تحقيق الشخصية

سهير
محمد حسين احمد
قرية البراجيل
مركز أوسيم - الجيزة

١٩٧٣/١٢

٢٧٣ ١٢ ٢٢ ٢١ ٠ ١٦ ٢٣

LT2414991

١١٤٩١٣٥٧٨١

١١٥٤٦٧٣٠١٢




٢٠٢٦/٠٥ ٢٧٣١٢٢٢٢١٠١٦٢٣

بدون عمل




أنثى
مسلمة
متزوجة
صابر اسماعيل السيد على
البطاقة سارية حتى ٢٠٣٣/٠٥/٠٥



Imaging report

Pat. Name: Sohair Mohamed	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026



كايرو سكان
للأشعة والتحاليل
CAIRO SCAN
Radiology & LABS

MR MAMMOGRAPHY OF BOTH BREASTS:

CLINICAL SUMMARY:

Age: 50yrs

Right breast pain and swelling

IMPRESSION:

LEFT BREAST: (BIRADS 1)

Normal study

RIGHT BREAST: (BIRADS 5)

Multi focal malignant mass with skin thickening and level II pathologically enlarged axillary LN.

RECOMENDATION:

For histopathological evaluation

TECHNIQUE:

Multiplanar imaging was performed on a 1.5T magnet with a dedicated breast coil, using the following frequencies:

- Axial T1, T2 IR
- Sagittal T2
- Subtraction of pre & post contrast 3D dynamic axial fat suppressed FLASH sequence 1.5 mm slice thickness after IV injection of Gd-DTPA (0.1 mmol/kg) with MIP image reformatting


FINDINGS:

The breasts are composed of heterogenous fibroglandular tissue and fat. There is mild background parenchymal enhancement

LEFT BREAST:

- T2WIs show no areas of abnormal signal intensity.
- After injection of contrast, no rapidly enhancing masses or other suspicious enhancement patterns were observed to suggest invasive malignancy.
- Hence, no Time -intensity enhancement curves were done.
- Normal skin / nipple / areolar complex.
- No pathological axillary lymph nodes.

Imaging report



كايرو سكان
للشعة والتصوير
CAIRO SCAN
Radiology & LABS

Pat. Name: Sohair Mohamed Hussein	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026

RIGHT BREAST:

- The middle aspect of the breast is the seat of two large speculated outline masses , 2 cm apart . The largest measures 5x4.6 cm reaching out to the nipple causing its retraction. The smaller measures 2.5x2.5 cm. They both elicit low T1 & T2 SI and high STIR SI
- After injection of contrast, they both showed dense contrast uptake.
- Diffuse skin thickening is noted.
- Level II pathological enlarged axillary lymph nodes are noted.

Best Regards
DR. DORRIA SALEM
Prof. of Radiology, Cairo University

For any inquiries regarding the findings in the report please contact us on
radreport@cairoscan.com.eg

Patient Name	سهر محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

Pathology Report

Clinical Data: Right breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 31/08/2026, Cold ischemia time is 20 seconds.

Gross Examination:

-Four greyish, white tissue cores, ranging in length from 0.4 cm & 1.5 cm, totally submitted.

Technique:

- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for:
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), & Ki67 clone (30-9).

Microscopic examination:

- Histologic type: Invasive duct carcinoma.
- Histologic grade: 3 (differentiation 3, pleomorphism 3, mitosis 2).
- DCIS: Absent.
- Microcalcification: Absent.
- Angioinvasion: Absent.
- Perineural invasion: Absent.
- Tumor infiltrating lymphocytes (TILs): Moderate.

Results:

* IPT (All controls showed appropriate reactivity):

- ER: Negative nuclear reaction in tumor cells (score 0/8).
- PR: Negative nuclear reaction in tumor cells (score 0/8).
- Her2: Strong complete membrane reaction in more than 10% of tumor cells (score 3+).
- Ki67%: 25%.

Conclusion:

Right breast mass, Tru-cut biopsy:

-Invasive duct carcinoma, Grade 3.

IPT:

- ER: Negative (score 0/8).
- PR: Negative (score 0/8).



Patient Name	سفيث محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

- Her2: Positive (score 3+).

- Ki67%: 25%.

****Scoring system: "ER, PR"**

ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66% and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, weak; 2, intermediate; and 3, strong). Total score from 0 to 8.

**** Scoring system: "Her2"**

Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.

IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within >10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.

Signature

Dr. Walaa Ibrahim Ghoneim, MD.



عقبة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر الجديدة - القاهرة 3 - مصر
Room No. (4) - First Floor-Building No. 987- Nile Corniche - Nisar El Qadima - Cairo, Egypt

naging report



Pat. Name: Sohair Mohamed Hussein	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026

RIGHT BREAST:

- The middle aspect of the breast is the seat of two large speculated outline masses , 2 cm apart . The largest measures 5x4.6 cm reaching out to the nipple causing its retraction. The smaller measures 2.5x2.5 cm. They both elicit low T1 & T2 SI and high STIR SI
- After injection of contrast, they both showed dense contrast uptake.
- Diffuse skin thickening is noted.
- Level II pathological enlarged axillary lymph nodes are noted.

Best Regards
DR. DORRIA SALEM
Prof. of Radiology, Cairo University

For any Inquiries regarding the findings in the report please contact us on
radreport@cairoscan.com.eg

Imaging report

Pat Name: Sohair Mohamed	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026



CAIRO SCAN
Radiology & MS

MR MAMMOGRAPHY OF BOTH BREASTS:

CLINICAL SUMMARY:

Age: 50yrs

Right breast pain and swelling

IMPRESSION:

LEFT BREAST: (BIRADS 1)

Normal study

RIGHT BREAST: (BIRADS 5)

Multi focal malignant mass with skin thickening and level II pathologically enlarged axillary LN.

RECOMMENDATION:

For histopathological evaluation

TECHNIQUE:

Multipolar imaging was performed on a 1.5T magnet with a dedicated breast coil, using the following frequencies:

- Axial T1, T2 IR
- Sagittal T2
- Subtraction of pre & post contrast 3D dynamic axial fat suppressed FLASH sequence 1.5 mm slice thickness after IV injection of Gd-DTPA (0.1 mmol/kg) with MIP image reformatting

FINDINGS:

The breasts are composed of heterogeneous fibroglandular tissue and fat. There is mild background parenchymal enhancement

LEFT BREAST:

- T2WIs show no areas of abnormal signal intensity.
- After injection of contrast, no rapidly enhancing masses or other suspicious enhancement patterns were observed to suggest invasive malignancy.
- Hence, no Time-intensity enhancement curves were done.
- Normal skin / nipple / areolar complex.
- No pathological axillary lymph nodes.

جمهورية مصر العربية
بطاقة تحقيق الشخصية

سهيير
محمد حسين احمد
قرية البراجيل
مركز اوسيم - الجيزة

٢٧٣ ١٢ ٢٢ ٢١ ٠١٦ ٢٣

LT2414991

٢٠٢٦/٠٥ ٢٧٣١٢٢٢٢١٠١٦٢٣
بدون عمل
أنثى مسلمة متزوجة
صابر اسماعيل السيد علي
البطاقة سارية حتى ٢٠٢٣/٠٥/٠٥ ١



Patient Name	سهر محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

Pathology Report

Clinical Data: Right breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 31/08/2026, Cold ischemia time is 20 seconds.

Gross Examination:

-Four greyish, white tissue cores, ranging in length from 0.4 cm & 1.5 cm, totally submitted.

Technique:

- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for:
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), & Ki67 clone (30-9).

Microscopic examination:

- Histologic type: Invasive duct carcinoma.
- Histologic grade: 3 (differentiation 3, pleomorphism 3, mitosis 2).
- DCIS: Absent.
- Microcalcification: Absent.
- Angioinvasion: Absent.
- Perineural invasion: Absent.
- Tumor infiltrating lymphocytes (TILs): Moderate.

Results:

* IPT (All controls showed appropriate reactivity):

- ER: Negative nuclear reaction in tumor cells (score 0/8).
- PR: Negative nuclear reaction in tumor cells (score 0/8).
- Her2: Strong complete membrane reaction in more than 10% of tumor cells (score 3+).
- Ki67%: 25%.

Conclusion:

Right breast mass, Tru-cut biopsy:

-Invasive duct carcinoma, Grade 3.

IPT:

- ER: Negative (score 0/8).
- PR: Negative (score 0/8).



Patient Name	سفيتر محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

- Her2: Positive (score 3+).

- Ki67%: 25%.

****Scoring system: "ER, PR"**

ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66% and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, weak; 2, intermediate; and 3, strong). Total score from 0 to 8.

**** Scoring system: "Her2"**

Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.

IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within > 10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.

Signature

Dr. Walaa Ibrahim Ghoneim, MD.



عقبة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر الجديدة - القاهرة 3 - مصر
 Room No. (4) - First Floor-Building No. 987- Nile Corniche - Nisar El Qadima - Cairo, Egypt

naging report



Pat. Name: Sohair Mohamed Hussein	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026

RIGHT BREAST:

- The middle aspect of the breast is the seat of two large speculated outline masses , 2 cm apart . The largest measures 5x4.6 cm reaching out to the nipple causing its retraction. The smaller measures 2.5x2.5 cm. They both elicit low T1 & T2 SI and high STIR SI
- After injection of contrast, they both showed dense contrast uptake.
- Diffuse skin thickening is noted.
- Level II pathological enlarged axillary lymph nodes are noted.

Best Regards
DR. DORRIA SALEM
Prof. of Radiology, Cairo University

For any inquiries regarding the findings in the report please contact us on
radreport@cairoscan.com.eg

Pat Name: Sohair Mohamed	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026



MR MAMMOGRAPHY OF BOTH BREASTS:

CLINICAL SUMMARY:

Age: 50yrs
Right breast pain and swelling

IMPRESSION:

LEFT BREAST: (BIRADS 1)
Normal study

RIGHT BREAST: (BIRADS 5)

Multi focal malignant mass with skin thickening and level II pathologically enlarged axillary LN.

RECOMMENDATION:

For histopathological evaluation

TECHNIQUE:

Multipolar imaging was performed on a 1.5T magnet with a dedicated breast coil, using the following frequencies:


- Axial T1, T2 IR
- Sagittal T2
- Subtraction of pre & post contrast 3D dynamic axial fat suppressed FLASH sequence 1.5 mm slice thickness after IV injection of Gd-DTPA (0.1 mmol/kg) with MIP image reformatting

FINDINGS:

The breasts are composed of heterogeneous fibroglandular tissue and fat. There is mild background parenchymal enhancement

LEFT BREAST:

- T2WIs show no areas of abnormal signal intensity.
- After injection of contrast, no rapidly enhancing masses or other suspicious enhancement patterns were observed to suggest invasive malignancy.
- Hence, no Time-intensity enhancement curves were done.
- Normal skin / nipple / areolar complex.
- No pathological axillary lymph nodes.



السيدة / سهيل محمد حسين

52 Years / Female

Herself

BIOCHEMISTRY REPORT

Kidney Functions:

Reference Range:

▲ Serum Creatinine

1.0

(Normal: 0.5 - 1.5 mg/dL)

signature
Dr. Hossain Hossain
LAB

Patient History In Our Lab.

Creat

26-Jun-2024

0.8



012277117105
01220066353

• شارع مصطفى كامل ، الشارع الكبير
البرجيل - أمام المسجد القبلي
مواعد العمل :
دائماً من الساعة ٩ صباحاً إلى ١١ مساءً

جمهورية مصر العربية
بطاقة تحقيق الشخصية

سهير
محمد حسين احمد
قرية البراجيل
مركز اوسيم - الجيزة

٢٧٣ ١٢ ٢٢ ٢١ ٠١٦ ٢٣

LT2414991

٢٠٢٦/٠٥ ٢٧٣١٢٢٢٢١٠١٦٢٣
بدون عمل
أنثى مسلمة متزوجة
صابر اسماعيل السيد علي
البطاقة سارية حتى ٢٠٢٣/٠٥/٠٥ ١



Patient Name	سهر محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

Pathology Report

Clinical Data: Right breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 31/08/2026, Cold ischemia time is 20 seconds.

Gross Examination:

-Four greyish, white tissue cores, ranging in length from 0.4 cm & 1.5 cm, totally submitted.

Technique:

- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for:
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), & Ki67 clone (30-9).

Microscopic examination:

- Histologic type: Invasive duct carcinoma.
- Histologic grade: 3 (differentiation 3, pleomorphism 3, mitosis 2).
- DCIS: Absent.
- Microcalcification: Absent.
- Angioinvasion: Absent.
- Perineural invasion: Absent.
- Tumor infiltrating lymphocytes (TILs): Moderate.

Results:

* IPT (All controls showed appropriate reactivity):

- ER: Negative nuclear reaction in tumor cells (score 0/8).
- PR: Negative nuclear reaction in tumor cells (score 0/8).
- Her2: Strong complete membrane reaction in more than 10% of tumor cells (score 3+).
- Ki67%: 25%.

Conclusion:

Right breast mass, Tru-cut biopsy:

-Invasive duct carcinoma, Grade 3.

IPT:

- ER: Negative (score 0/8).
- PR: Negative (score 0/8).



Patient Name	سفيث محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

- Her2: Positive (score 3+).

- Ki67%: 25%.

****Scoring system: "ER, PR"**

ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66% and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, weak; 2, intermediate; and 3, strong). Total score from 0 to 8.

**** Scoring system: "Her2"**

Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.

IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within > 10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.

Signature

Dr. Walaa Ibrahim Ghoneim, MD.



عقيدة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر القديمة - القاهرة 5 - مصر
 Room No. (4) - First Floor-Building No. 987- Nile Corniche - Mizr El Qadima - Cairo, Egypt

naging report



Pat. Name: Sohair Mohamed Hussein	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026

RIGHT BREAST:

- The middle aspect of the breast is the seat of two large speculated outline masses , 2 cm apart . The largest measures 5x4.6 cm reaching out to the nipple causing its retraction. The smaller measures 2.5x2.5 cm. They both elicit low T1 & T2 SI and high STIR SI
- After injection of contrast, they both showed dense contrast uptake.
- Diffuse skin thickening is noted.
- Level II pathological enlarged axillary lymph nodes are noted.

Best Regards
DR. DORRIA SALEM
Prof. of Radiology, Cairo University

For any inquiries regarding the findings in the report please contact us on
radreport@cairoscan.com.eg

Pat Name: Sohair Mohamed	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026



MR MAMMOGRAPHY OF BOTH BREASTS:

CLINICAL SUMMARY:

Age: 50yrs
Right breast pain and swelling

IMPRESSION:

LEFT BREAST: (BIRADS 1)
Normal study

RIGHT BREAST: (BIRADS 5)

Multi focal malignant mass with skin thickening and level II pathologically enlarged axillary LN.

RECOMMENDATION:

For histopathological evaluation

TECHNIQUE:

Multipolar imaging was performed on a 1.5T magnet with a dedicated breast coil, using the following frequencies:


- Axial T1, T2 IR
- Sagittal T2
- Subtraction of pre & post contrast 3D dynamic axial fat suppressed FLASH sequence 1.5 mm slice thickness after IV injection of Gd-DTPA (0.1 mmol/kg) with MIP image reformatting

FINDINGS:

The breasts are composed of heterogeneous fibroglandular tissue and fat. There is mild background parenchymal enhancement

LEFT BREAST:

- T2WIs show no areas of abnormal signal intensity.
- After injection of contrast, no rapidly enhancing masses or other suspicious enhancement patterns were observed to suggest invasive malignancy.
- Hence, no Time-intensity enhancement curves were done.
- Normal skin / nipple / areolar complex.
- No pathological axillary lymph nodes.



السيدة / سهيل محمد حسين

1 026080 510047

05-Aug-2026 11:48 AM

52 Years / Female

Herself

BIOCHEMISTRY REPORT

Kidney Functions:

Reference Range:

▲ Serum Creatinine

1.0

(Normal: 0.5 - 1.5 mg/dL)

signature
Dr. Hossam Hassan
LAB

Patient History In Our Lab.

Creat

26-Jun-2024

0.8



SEGMA LAB

☎ 01277117105

 012220066353

• شارع مصطفى كامل ، الشارع الكبير
البرجيل - أمام المسجد القبلي
مواعد العمل :
دائماً من الساعة ٩ صباحاً إلى ١١ مساءً

جمهورية مصر العربية
بطاقة تحقيق الشخصية

سهيير
محمد حسين احمد
قرية البراجيل
مركز اوسيم - الجيزة

٢٧٣ ١٢ ٢٢ ٢١ ٠ ١٦ ٢٣

LT2414991

٢٠٢٦/٠٥ ٢٧٣١٢٢٢٢١٠١٦٢٣
بدون عمل
أنثى مسلمة متزوجة
صابر اسماعيل السيد علي
البطاقة سارية حتى ٢٠٢٣/٠٥/٠٥ ١



Patient Name	سهر محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

Pathology Report

Clinical Data: Right breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 31/08/2026, Cold ischemia time is 20 seconds.

Gross Examination:

-Four greyish, white tissue cores, ranging in length from 0.4 cm & 1.5 cm, totally submitted.

Technique:

- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for:
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), & Ki67 clone (30-9).

Microscopic examination:

- Histologic type: Invasive duct carcinoma.
- Histologic grade: 3 (differentiation 3, pleomorphism 3, mitosis 2).
- DCIS: Absent.
- Microcalcification: Absent.
- Angioinvasion: Absent.
- Perineural invasion: Absent.
- Tumor infiltrating lymphocytes (TILs): Moderate.

Results:

* IPT (All controls showed appropriate reactivity):

- ER: Negative nuclear reaction in tumor cells (score 0/8).
- PR: Negative nuclear reaction in tumor cells (score 0/8).
- Her2: Strong complete membrane reaction in more than 10% of tumor cells (score 3+).
- Ki67%: 25%.

Conclusion:

Right breast mass, Tru-cut biopsy:

-Invasive duct carcinoma, Grade 3.

IPT:

- ER: Negative (score 0/8).
- PR: Negative (score 0/8).



Patient Name	سفيث محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

- Her2: Positive (score 3+).

- Ki67%: 25%.

****Scoring system: "ER, PR"**

ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66% and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, weak; 2, intermediate; and 3, strong). Total score from 0 to 8.

**** Scoring system: "Her2"**

Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.

IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within > 10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.

Signature

Dr. Walaa Ibrahim Ghoneim, MD.



عرة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر القديمة - القاهرة 5 - مصر
 Room No. (4) - First Floor-Building No. 987- Nile Corniche - Mkr El Qadima - Cairo, Egypt

naging report



Pat. Name: Sohair Mohamed Hussein	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026

RIGHT BREAST:

- The middle aspect of the breast is the seat of two large speculated outline masses , 2 cm apart . The largest measures 5x4.6 cm reaching out to the nipple causing its retraction. The smaller measures 2.5x2.5 cm. They both elicit low T1 & T2 SI and high STIR SI
- After injection of contrast, they both showed dense contrast uptake.
- Diffuse skin thickening is noted.
- Level II pathological enlarged axillary lymph nodes are noted.

Best Regards
DR. DORRIA SALEM
Prof. of Radiology, Cairo University

For any inquiries regarding the findings in the report please contact us on
radreport@cairoscan.com.eg

Imaging report

Pat Name: Sohair Mohamed	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026



CAIRO SCAN
Radiology & MS

MR MAMMOGRAPHY OF BOTH BREASTS:

CLINICAL SUMMARY:

Age: 50yrs
Right breast pain and swelling

IMPRESSION:

LEFT BREAST: (BIRADS 1)

Normal study

RIGHT BREAST: (BIRADS 5)

Multi focal malignant mass with skin thickening and level II pathologically enlarged axillary LN.

RECOMMENDATION:

For histopathological evaluation

TECHNIQUE:

Multipolar imaging was performed on a 1.5T magnet with a dedicated breast coil, using the following frequencies:


- Axial T1, T2 IR
- Sagittal T2
- Subtraction of pre & post contrast 3D dynamic axial fat suppressed FLASH sequence 1.5 mm slice thickness after IV injection of Gd-DTPA (0.1 mmol/kg) with MIP image reformatting

FINDINGS:

The breasts are composed of heterogeneous fibroglandular tissue and fat. There is mild background parenchymal enhancement

LEFT BREAST:

- T2WIs show no areas of abnormal signal intensity.
- After injection of contrast, no rapidly enhancing masses or other suspicious enhancement patterns were observed to suggest invasive malignancy.
- Hence, no Time-intensity enhancement curves were done.
- Normal skin / nipple / areolar complex.
- No pathological axillary lymph nodes.



Herself

Kidney Functions:

• شارع مصطفى كامل ، الشارع الكبير
البرجيل - أمام المسجد القبلي
مواعد العمل :
دائماً من الساعة ٩ صباحاً إلى ١١ مساءً

جمهورية مصر العربية
بطاقة تحقيق الشخصية

سهيير
محمد حسين احمد
قرية البراجيل
مركز اوسيم - الجيزة

٢٧٣ ١٢ ٢٢ ٢١ ٠١٦ ٢٣

LT2414991

٢٠٢٦/٠٥ ٢٧٣١٢٢٢٢١٠١٦٢٣
بدون عمل
أنثى مسلمة متزوجة
صابر اسماعيل السيد علي
البطاقة سارية حتى ٢٠٢٣/٠٥/٠٥ ١



Patient Name	سهر محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

Pathology Report

Clinical Data: Right breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 31/08/2026, Cold ischemia time is 20 seconds.

Gross Examination:

-Four greyish, white tissue cores, ranging in length from 0.4 cm & 1.5 cm, totally submitted.

Technique:

- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for:
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), & Ki67 clone (30-9).

Microscopic examination:

- Histologic type: Invasive duct carcinoma.
- Histologic grade: 3 (differentiation 3, pleomorphism 3, mitosis 2).
- DCIS: Absent.
- Microcalcification: Absent.
- Angioinvasion: Absent.
- Perineural invasion: Absent.
- Tumor infiltrating lymphocytes (TILs): Moderate.

Results:

* IPT (All controls showed appropriate reactivity):

- ER: Negative nuclear reaction in tumor cells (score 0/8).
- PR: Negative nuclear reaction in tumor cells (score 0/8).
- Her2: Strong complete membrane reaction in more than 10% of tumor cells (score 3+).
- Ki67%: 25%.

Conclusion:

Right breast mass, Tru-cut biopsy:

-Invasive duct carcinoma, Grade 3.

IPT:

-ER: Negative (score 0/8).

-PR: Negative (score 0/8).



Patient Name	سفيث محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

- Her2: Positive (score 3+).

- Ki67%: 25%.

****Scoring system: "ER, PR"**

ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66% and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, weak; 2, intermediate; and 3, strong). Total score from 0 to 8.

**** Scoring system: "Her2"**

Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.

IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within > 10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.

Signature

Dr. Walaa Ibrahim Ghoneim, MD.



عقبة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر الجديدة - القاهرة 3 - مصر
 Room No. (4) - First Floor-Building No. 987- Nile Corniche - Nisar El Qadima - Cairo, Egypt

naging report



Pat. Name: Sohair Mohamed Hussein	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026

RIGHT BREAST:

- The middle aspect of the breast is the seat of two large speculated outline masses , 2 cm apart . The largest measures 5x4.6 cm reaching out to the nipple causing its retraction. The smaller measures 2.5x2.5 cm. They both elicit low T1 & T2 SI and high STIR SI
- After injection of contrast, they both showed dense contrast uptake.
- Diffuse skin thickening is noted.
- Level II pathological enlarged axillary lymph nodes are noted.

Best Regards

DR. DORRIA SALEM

Prof. of Radiology, Cairo University

For any inquiries regarding the findings in the report please contact us on
radreport@cairoscan.com.eg

Pat Name: Sohair Mohamed	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026



MR MAMMOGRAPHY OF BOTH BREASTS:

CLINICAL SUMMARY:

Age: 50yrs
Right breast pain and swelling

IMPRESSION:

LEFT BREAST: (BIRADS 1)
Normal study

RIGHT BREAST: (BIRADS 5)

Multi focal malignant mass with skin thickening and level II pathologically enlarged axillary LN.

RECOMMENDATION:

For histopathological evaluation

TECHNIQUE:

Multipolar imaging was performed on a 1.5T magnet with a dedicated breast coil, using the following frequencies:


- Axial T1, T2 IR
- Sagittal T2
- Subtraction of pre & post contrast 3D dynamic axial fat suppressed FLASH sequence 1.5 mm slice thickness after IV injection of Gd-DTPA (0.1 mmol/kg) with MIP image reformatting

FINDINGS:

The breasts are composed of heterogeneous fibroglandular tissue and fat. There is mild background parenchymal enhancement

LEFT BREAST:

- T2WIs show no areas of abnormal signal intensity.
- After injection of contrast, no rapidly enhancing masses or other suspicious enhancement patterns were observed to suggest invasive malignancy.
- Hence, no Time-intensity enhancement curves were done.
- Normal skin / nipple / areolar complex.
- No pathological axillary lymph nodes.



السيدة / سهيل محمد حسين

52 Years / Female

Herself

BIOCHEMISTRY REPORT

Kidney Functions:

Reference Range:

▲ Serum Creatinine

1.0

(Normal: 0.5 - 1.5 mg/dL)

signature
Dr. Hossam Hassan
LAB

Patient History In Our Lab.

Creat

26-Jun-2024

0.8



SEGMA LAB

01277117105

 012220066353

• شارع مصطفى كامل ، الشارع الكبير
البرجيل - أمام المسجد القبلي
مواعد العمل :
دواماً من الساعة ٩ صباحاً إلى ١١ مساءً

جمهورية مصر العربية
بطاقة تحقيق الشخصية

سهيير
محمد حسين احمد
قرية البراجيل
مركز اوسيم - الجيزة

٢٧٣ ١٢ ٢٢ ٢١ ٠ ١٦ ٢٣

LT2414991

٢٠٢٦/٠٥ ٢٧٣١٢٢٢٢١٠١٦٢٣
بدون عمل

أنثى مسلمة متزوجة
صابر اسماعيل السيد علي

البطاقة سارية حتى ٢٠٢٣/٠٥/٠٥ ١



Patient Name	سهر محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

Pathology Report

Clinical Data: Right breast mass (BIRADS 5).

Nature of Specimen: Tru-cut biopsy, collected on 31/08/2026, Cold ischemia time is 20 seconds.

Gross Examination:

-Four greyish, white tissue cores, ranging in length from 0.4 cm & 1.5 cm, totally submitted.

Technique:

- Sections prepared from paraffin block for immune-staining using Bench Mark ULTRA system for:
- ER clone (1E2), PR clone (SP1), Her2 clone (4B5), & Ki67 clone (30-9).

Microscopic examination:

- Histologic type: Invasive duct carcinoma.
- Histologic grade: 3 (differentiation 3, pleomorphism 3, mitosis 2).
- DCIS: Absent.
- Microcalcification: Absent.
- Angioinvasion: Absent.
- Perineural invasion: Absent.
- Tumor infiltrating lymphocytes (TILs): Moderate.

Results:

* IPT (All controls showed appropriate reactivity):

- ER: Negative nuclear reaction in tumor cells (score 0/8).
- PR: Negative nuclear reaction in tumor cells (score 0/8).
- Her2: Strong complete membrane reaction in more than 10% of tumor cells (score 3+).
- Ki67%: 25%.

Conclusion:

Right breast mass, Tru-cut biopsy:

-Invasive duct carcinoma, Grade 3.

IPT:

-ER: Negative (score 0/8).

-PR: Negative (score 0/8).



Patient Name	سفيث محمد حسين	Case No.	361786
Sex/Age	Female 52 Y	Patient No.	26467
Registered Date	31/8/2026 06:30:17 PM	Referred By Dr.	Interventional Radiology
Validation Date	17/9/2026 11:01:21 AM	Branch	Gustave Roussy
Reporting Date	17/9/2026 01:07:15 PM	Pathology No.	2769/26

- Her2: Positive (score 3+).

- Ki67%: 25%.

****Scoring system: "ER, PR"**

ASCO-CAP ER/PR Test Guideline Recommendations: proportion of cells staining (0, no staining; 1, >1%; 2, between 1% and 10%; 3, between 11% and 33%; 4, between 34% and 66% and 5, between 67%-100% of the cells staining). Intensity of positive tumor cells (0, none; 1, weak; 2, intermediate; and 3, strong). Total score from 0 to 8.

**** Scoring system: "Her2"**

Reference: ASCO-CAP Her2 Test Guideline Recommendation. IHC 0 (Null): as defined by no staining observed. IHC 0+ (Ultra-low) as defined by membrane staining that is incomplete and is faint/barely perceptible and within < 10% of the invasive tumor cells. IHC 1+: as defined by incomplete membrane staining that is faint/barely perceptible and within > 10% of the invasive tumor cells.

IHC 2+: based on circumferential membrane staining that is incomplete and/or weak/moderate and within > 10% of the invasive tumor cells or complete and circumferential membrane staining that is intense and within < 10% of the invasive tumor cells. IHC 3+: based on circumferential membrane staining that is complete, intense > 10% of tumor cells.

Signature

Dr. Walaa Ibrahim Ghoneim, MD.



عقبة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر الجديدة - القاهرة 3 - مصر
 Room No. (4) - First Floor-Building No. 987- Nile Corniche - Nisar El Qadima - Cairo, Egypt

naging report



كايرو سكان
مؤسسة والتجارية
CAIRO SCAN
Radiology & LABS

Pat. Name: Sohair Mohamed Hussein	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026

RIGHT BREAST:

- The middle aspect of the breast is the seat of two large speculated outline masses , 2 cm apart . The largest measures 5x4.6 cm reaching out to the nipple causing its retraction. The smaller measures 2.5x2.5 cm. They both elicit low T1 & T2 SI and high STIR SI
- After injection of contrast, they both showed dense contrast uptake.
- Diffuse skin thickening is noted.
- Level II pathological enlarged axillary lymph nodes are noted.

Best Regards

DR. DORRIA SALEM

Prof. of Radiology, Cairo University

For any inquiries regarding the findings in the report please contact us on
radreport@cairoscan.com.eg

Imaging report

Pat Name: Sohair Mohamed	Pat. ID: 3618519
Procedure Name: MRI Breasts	Date: 06/08/2026



CAIRO SCAN
Radiology & MS

MR MAMMOGRAPHY OF BOTH BREASTS:

CLINICAL SUMMARY:

Age: 50yrs

Right breast pain and swelling

IMPRESSION:

LEFT BREAST: (BIRADS 1)

Normal study

RIGHT BREAST: (BIRADS 5)

Multi focal malignant mass with skin thickening and level II pathologically enlarged axillary LN.

RECOMMENDATION:

For histopathological evaluation

TECHNIQUE:

Multipolar imaging was performed on a 1.5T magnet with a dedicated breast coil, using the following frequencies:

- Axial T1, T2 IR
- Sagittal T2
- Subtraction of pre & post contrast 3D dynamic axial fat suppressed FLASH sequence 1.5 mm slice thickness after IV injection of Gd-DTPA (0.1 mmol/kg) with MIP image reformatting

FINDINGS:

The breasts are composed of heterogeneous fibroglandular tissue and fat. There is mild background parenchymal enhancement

LEFT BREAST:

- T2WIs show no areas of abnormal signal intensity.
- After injection of contrast, no rapidly enhancing masses or other suspicious enhancement patterns were observed to suggest invasive malignancy.
- Hence, no Time-intensity enhancement curves were done.
- Normal skin / nipple / areolar complex.
- No pathological axillary lymph nodes.



Patient ID.	 1 026080 510047	Patient Name	السيدة / سفيان محمد حسين
Request Date	05-Aug-2026 11:48 AM	Age / Sex	52 Years / Female
Referred By	Herself		

BIOCHEMISTRY REPORT

Kidney Functions :

▲ Serum Creatinine : 1.0 (Normal: 0.5 - 1.5 mg/dL)

Reference Range :

signature
Dr. Hossam Hassan
LAB

Patient History In Our Lab.

26-Jun-2024 Creat 0.8