

رقم البطاقة : 28510022101564 رقم التليفون : 01000428787
العمر(تاريخ الميلاد) : 02-10-1985 المهنة : حاصل على دبلوم
تاريخ تسجيل الإستمارة : 28-09-2026

أقر انا الموقع أدناه انني لا اتمتع باي نظام علاج مثل:- التأمين الصحي-القوات المسلحة-الشرطة-
الهيئات-المؤسسات او اي نظام يتعارض مع نظام العلاج علي نفقة الدولة- وسوف اتحمل كافة
نفقات العلاج في حالة ثبوت غير ذلك

الإخصائي الاجتماعي (-----) التوقيع: (-----)

المقرر بما فيه
التوقيع: (-----)

التعليق على التشخيص : -----

تتعهد المستشفى أن لا يتم تنفيذ القرار إلا بعد مراجعة الملف الطبي للمريض وإعادة مناظرة المريض بالمستشفى قبل الصرف.

----- / 2 ----- / 2 ----- / 2.1 ----- / 2.1

ملاحظات :

رئيس
المجلس الطبي المتخصص

--

Name: رشا ناجي عبد الهادي حجازي National ID: 28510022101564	
--	---

تقرير طبي

السيد
تحية طيبة وبعد,,

إيماء للطلب المقدم من	بشأن تقرير طبي للمريض
بقسم:	رقم طبي:

نتشرف بأن نوضح فيما يلي تقريراً عن حالته
--

تاريخ وساعة الدخول : -- --:-- yyyy-mm-dd	تاريخ وساعة الخروج : PM 01:17 05-07-2026
---	---

وبعد الكشف والفحوصات والمتابعة وجد أنه
--

A patient of uterine cancer for CT scan of the chest.

التاريخ : 05-07-2026	الوقت : -- --:--
	الطبيب المعالج : د/شيماء لاشين


جمهورية مصر العربية
بطاقة تحقيق الشخصية


رشا
ناجى عبدالهادى حجازى
قرية السبيل
منشأة القناطر - الجيزة

١٩٨٥/١ ٢٨٥ ١٠ ٠٢ ٢١ ٠١٥ ٦٤

KH4261189

٢٠٢٣/١١ ٢٨٥١٠٠٢٢١٠١٥٦٤
حاصله على دبلوم تجاره
أنثى مسلمة متزوجة
عاشور عبدالباري على عجور
البطاقة سارية حتى ٢٠٣٠/١١/٢٨ ١





Patient's ID & Sex: 01095186303RNA, F

Accession No: undefined

Study Date: 14-Sep-2026

Examination: MR PELVIS without contrast

Technique: Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.

Patient's Name: RASHA NAGY A.AL HADY 40, 040Y

Referred By: داعزة احمد

Reporting Date: 15-Sep-2026



Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity, and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD

كاشف
Scan
15/09/2026
د. غادة عوض



اسم المريض /

التاريخ: ٢٧

رقم التذكرة:

نموذج تقرير سونار (حوض)

Pelvic Ultrasound Study in Female

Uterus :

Av. ^{mean} gravid uterus, it shows ^{homogeneous}

echo pattern. It measures about 7 X 5 cm.

endometrial thicknes = 5 mm

Ovaries :

in size (— x —),

pathological lesions .

Rt. simple ovarian cyst
measured (2x2) —
for clinical correlation & treatment
lt. masked by gases

Free fluid :


Line of fluid in Douglas pouch

for clinical correlation

Signature :

Best Regards

صحتك أمانه بأيدينا



وادي النيل
للأشعة
wadi el Nile
scan



Patient's ID & Sex	01095186303RNA, F	Patient's Name	RASHA NACY A.AL HADY 40, 040Y
Accession No	undefined	Referved By	لاعزّة احمد
Study Date	14-Sep-2026	Reporting Date	15-Sep-2026
Examination	MR PELVIS without contrast		
Technique	Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.		

Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD

Ghada Awad
RASHA NACY A.AL HADY
15-Sep-2026
لاعزّة احمد

جمهورية مصر العربية
بطاقة تحقيق الشخصية



رشا
ناجى عبدالهادى حجازى
قرية السبيل
منشأة القناطر - الجيزة

١٩٨٥/١ ٢٨٥ ١٠ ٠٢ ٢١ ٠١٥ ٦٤

KH4261189





Patient's ID & Sex	01095186303RNA, F	Patient's Name	RASHA NACY A.AL HADY 40, 040Y
Accession No	undefined	Referved By	لاعزّة احمد
Study Date	14-Sep-2026	Reporting Date	15-Sep-2026
Examination	MRI PELVIS without contrast		
Technique	Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.		

Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD


RASHA NACY
15-Sep-2026
لاعزّة احمد



اسم المريض /
التاريخ: ٢٥/٥
رقم الملف: ٢٢٢

نموذج تقرير سونار (حوض)

Pelvic Ultrasound Study in Female

Trans-void
Trans-void

Av. 12w gravid uterus, it shows *homogenous*

Uterus :

echo pattern. It measures about *F 5* cm.

endometrial thickness = *5* mm

Ovaries :

in size (x x),

pathological lesions .

*Rt. simple cyst
measured (2x2) cm
for clinical condition
It. masked by gels*

Free fluid : *line of fluid in Douglas pouch*

for clinical condition
Signature : *Dr. H. S.*

Best Regards

جمهورية مصر العربية
بطاقة تحقيق الشخصية



رشا
ناجى عبدالهادى حجازى
قرية السبيل
منشأة القناطر - الجيزة

١٩٨٥/١ ٢٨٥ ١٠ ٠٢ ٢١ ٠١٥ ٦٤

KH4261189





Patient's ID & Sex	01095186303RNA, F	Patient's Name	RASHA NACY A.AL HADY 40, 040Y
Accession No	undefined	Referved By	لاعزّة احمد
Study Date	14-Sep-2026	Reporting Date	15-Sep-2026
Examination	MRI PELVIS without contrast		
Technique	Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.		

Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD


RASHA NACY
15-Sep-2026
لاعزّة احمد



اسم المريض /
التاريخ: ٢٥/٥
رقم الملف: ٢٢٢

نموذج تقرير سونار (حوض)

Pelvic Ultrasound Study in Female

Trans-void
Trans-void

Av. 12w gravid uterus, it shows *homoechoic*

Uterus :

echo pattern. It measures about *F 5* cm.

endometrial thickness = *5* mm

Ovaries :

in size (x x),

pathological lesions .

*Rt. simple ovarian cyst
measured (2x2) cm
for clinical correlation & follow up
It. masked by gels*

Free fluid : *Free of fluid in Douglas pouch*

for clinical correlation
Signature : *Dr. H. S.*

Best Regards

جمهورية مصر العربية
بطاقة تحقيق الشخصية



رشا
ناجى عبدالهادى حجازى
قرية السبيل
منشأة القناطر - الجيزة

١٩٨٥/١ ٢٨٥ ١٠ ٠٢ ٢١ ٠١٥ ٦٤

KH4261189





Patient's ID & Sex	01095186303RNA, F	Patient's Name	RASHA NACY A.AL HADY 40, 040Y
Accession No	undefined	Referved By	لاعزّة احمد
Study Date	14-Sep-2026	Reporting Date	15-Sep-2026
Examination	MRI PELVIS without contrast		
Technique	Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.		

Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD


RASHA NACY
15-Sep-2026
لاعزّة احمد



اسم المريض /
التاريخ: ٢٥/٥
رقم الملف: ٢٢٢

نموذج تقرير سونار (حوض)

Pelvic Ultrasound Study in Female

Trans-void
Trans-void

Av. 12w gravid uterus, it shows *homo-geneous*

Uterus :

echo pattern. It measures about *F 5* cm.

endometrial thickness = *5* mm

Ovaries :

in size (x x),

pathological lesions .

*Rt. simple ovarian cyst
measured (2x2) cm
for clinical correlation & follow up
It. masked by gels*

Free fluid : *Free of fluid in Douglas pouch*

for clinical correlation
Signature : *[Signature]*

Best Regards

جمهورية مصر العربية
بطاقة تحقيق الشخصية



رشا
ناجى عبدالهادى حجازى
قرية السبيل
منشأة القناطر - الجيزة

١٩٨٥/١ ٢٨٥ ١٠ ٠٢ ٢١ ٠١٥ ٦٤

KH4261189





Patient's ID & Sex	01095186303RNA, F	Patient's Name	RASHA NACY A.AL HADY 40, 040Y
Accession No	undefined	Referved By	لاعزّة احمد
Study Date	14-Sep-2026	Reporting Date	15-Sep-2026
Examination	MRI PELVIS without contrast		
Technique	Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.		

Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD


RASHA NACY
15-Sep-2026
لاعزّة احمد



اسم المريض /
التاريخ: ٢٥/٥
رقم الملف: ٢٢٢

نموذج تقرير سونار (حوض)

Pelvic Ultrasound Study in Female

Trans-void
Trans-void

Av. 12w gravid uterus, it shows *homo-geneous*

Uterus :

echo pattern. It measures about *F 5* cm.

endometrial thickness = *5* mm

Ovaries :

in size (x x),

pathological lesions .

*Rt. simple ovarian cyst
measured (2x2) cm
for clinical correlation & follow up
It. masked by gels*

Free fluid : *Free of fluid in Douglas pouch*

for clinical correlation
Signature : *Dr. H. S.*

Best Regards

جمهورية مصر العربية
بطاقة تحقيق الشخصية



رشا
ناجى عبدالهادى حجازى
قرية السبيل
منشأة القناطر - الجيزة

١٩٨٥/١ ٢٨٥ ١٠ ٠٢ ٢١ ٠١٥ ٦٤

KH4261189





Patient's ID & Sex	01095186303RNA, F	Patient's Name	RASHA NACY A.AL HADY 40, 040Y
Accession No	undefined	Referved By	لاعزّة احمد
Study Date	14-Sep-2026	Reporting Date	15-Sep-2026
Examination	MRI PELVIS without contrast		
Technique	Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.		

Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD


RASHA NACY A.AL HADY
15-Sep-2026
لاعزّة احمد



اسم المريض /
التاريخ: ٢٥/٥
رقم الملف: ٢٢٢

نموذج تقرير سونار (حوض)

Pelvic Ultrasound Study in Female

Trans-void
Trans-void

Av. 12w gravid uterus, it shows *homo*

Uterus :

echo pattern. It measures about **F X 5** cm.

endometrial thickness = **5** mm

Ovaries :

in size (**5** x **5**),

pathological lesions .

*Rt. simple cyst
measured (2x2) cm
for clinical condition
It. masked by gels*

Free fluid : *Free of fluid in Douglas pouch*

for clinical condition
Signature : *[Signature]*

Best Regards

جمهورية مصر العربية
بطاقة تحقيق الشخصية



رشا
ناجى عبدالهادى حجازى
قرية السبيل
منشأة القناطر - الجيزة

١٩٨٥/١ ٢٨٥ ١٠ ٠٢ ٢١ ٠١٥ ٦٤

KH4261189





Patient's ID & Sex	01095186303RNA, F	Patient's Name	RASHA NACY A.AL HADY 40, 040Y
Accession No	undefined	Referved By	لاعزّة احمد
Study Date	14-Sep-2026	Reporting Date	15-Sep-2026
Examination	MRI PELVIS without contrast		
Technique	Multiple weighted images in multiple planes in addition to axial diffusion-weighted images were obtained through the pelvis.		

Clinical Info

Pain

Comparison: No available studies for comparison.

Findings:

- The uterus is within mildly bulky in size measuring 9x5.5x5 cm , showing normal MRI features of the endometrial cavity and myometrium. endometrial thickness measuring about 10 mm.
- Mild diffuse thickening of the junctional zone measured 12.5 mm with few mainly fundal sub endometrial cysts are seen.
- No detectable myomas.
- Right adenexial loculated lesion is seen measuring 3x4 cm displaying low T1, high T2 .
- Unremarkable MRI appearance of the cervix.
- The left ovary showing normal follicular arrangement. No apparent pathological changes are seen.
- The pelvic bones show regular cortical outlines and normal homogenous medullary bone intensity, with no definite sizable focal erosive or destructive lesions seen.
- The surrounding muscles and soft tissue appear normal. No mass lesions or fluid collections.

Impression:

- The above described picture is impressive of diffuse adenomyosis.
- Right adenexial simple cyst.

Dr. Ghada Awad, MD


RASHA NACY
15-Sep-2026
لاعزّة احمد



اسم المريض /
التاريخ: ٢٥/٥
رقم الملف: ٢٢٢

نموذج تقرير سونار (حوض)

Pelvic Ultrasound Study in Female

Trans-void
Trans-void

Uterus :

Av. 12w gravid uterus, it shows homogeneous

echo pattern. It measures about 7 X 5 cm.

endometrial thickness = 5 mm

Ovaries :

in size (X X),

pathological lesions .

Rt. simple cyst
measured (2x2) cm
for clinical correlation
It. masked by gels

Free fluid : Rim of fluid in Douglas pouch

for clinical correlation
Signature :
.....

Best Regards