



جوستاف روسي انترناشيونال - مصر (EPE) * طلب علاج أول مرة / متكرر

إسم المريض رباعياً : شرين علي محمد جاد

المحافظة : بورسعيد

الحي / المدينة : الشرق

العنوان : الشراييه

رقم البطاقة : 27908030101701 رقم التليفون : 01128896060

تاريخ تسجيل الإستمارة : 26-09-2026

العمر (تاريخ الميلاد) : 03-08-1979 المهنة : ربه منزل

التشخيص الإكلينيكي : سرطان بالثدي (مبادرة)

نتائج الأبحاث : (مرفق)

التشخيص النهائي : سرطان بالثدي (مبادرة)

خطة العلاج : علاج مبادرة سرطان الثدي العلاج المساعد لوقاية العظام للمرضي
الذين يتلقوا علاج هرموني حمض الزيلودرونك [14157]

قسم العلاج :
الاورام

التعليق على التشخيص : -----

خطاب إداري

هذا الرقم القومي يتبع منظومة التأمين الصحي الشامل جهة الاختصاص



--

Name: شرين علي محمد جاد National ID: 27908030101701	
--	---

تقرير طبي

السيد
تحية طبية وبعد,,

إيماء للطلب المقدم من	بشأن تقرير طبي للمريض
بقسم:	رقم طبي:

نتشرف بأن نوضح فيما يلي تقريراً عن حالته
--

تاريخ وساعة الدخول : -- --:-- yyyy-mm-dd	تاريخ وساعة الخروج : PM 01:17 05-07-2026
---	---

وبعد الكشف والفحوصات والمتابعة وجد أنه
--

المریضة تعاني من ورم غير حميد بالثدي و تحتاج الي Zoledronic acid for bone protection during hormonal therapy
--

التاريخ : 05-07-2026	الوقت : -- --:--
	الطبيب المعالج : د/شيماء لاشين

١

جمهورية مصر العربية
بطاقة تحقيق الشخصية

لترين
علي محمد جواد
١٩ ش النور - هاشم اغا
الشرابية - القاهرة

٢٧٩٠٨٣٠١٠١٧٠١

٢٠١٩/٠٢/٠١

HX5516489

٢٠١٩/٠٢/٠١ ٢٧٩٠٨٣٠١٠١٧٠١

بدون عمل

أنشئ
مسجلة
اسمها حسين السيد بدر
البطاقة سارية حتى ٢٠٢٩/٠٢/٠١

١

Patient Name : الأستاذ / شربين على محمد جاد
Patient No. : 27908030101701
Path No. : 72/25
Sex : Female
Age : 45 Y
Referring Doc :

Request No. : 13001787
Request Date : 06/01/2025
Verification Date : 16/02/2025
Financial Category : اورام دار السلام - ميانره
Department : Out Patient
Registered by : Mrs.Mayada Hegazy

Pathology Report

Clinical Data: Right breast cancer, post NACT.

Nature Specimen: Right modified radical mastectomy specimen.

Gross Examination:

- * Procedure: Right MRM measured 24x17x8 cm with covering skin ellipse measured 16x22 cm.
- * Tumor extent: An ill defined greyish white rubbery tumor bed site.
- * Size: 1.5 x0.5 x0.5 cm.
- * Skin involvement: Nipple & areola were unremarkable grossly.
- * Distance from margins: 4 cm away from deep margin.
- * Lymph nodes number and size: Fibrofatty tissue piece measured 12x5x2 cm. Careful repeated dissection entangling only 9 L.Ns up to 2x1.5 cm.

Microscopic Examination:

- * Histological type: Residual viable invasive duct carcinoma.
- * Grade (differentiation+ nuclear pleomorphism+ mitosis): (3+2+1) Grade II.
- Maximal tumor diameter 1.5 x 0.5 cm
- Actual tumor cellularity: 25%, tumor necrosis index: 75%
- * Ductal carcinoma in situ: Positive DCIS (5%).
- * Skin involvement: Free nipple and areola.
- * Lymph node involvement: Positive in (4/9) with extranodal extension. Maximal tumour diameter 13 mm. Two lymph nodes showed effect of therapy.
- * Resection Margins: Free deep margin.
- * Histologic response to therapy: Partial pathological response.
- * Microcalcifications: No.
- * Additional pathologic findings: Fibrosis, hyalinosis (Effect of therapy).

Verified by :
DR.Menna Nabil

Printing Date : 19/02/2025 12:13:25

Page 1 of 2

Printing By : Mrs.Mayada Hegazy



غرفة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر القديمة - القاهرة - مصر
Room No. (4) - First Floor-Building No. 987- Nile Corniche - Misr El Qadema - Cairo, Egypt



Patient Name : الأستاذ / شرين على محمد جاد

Patient No. : 27908030101701

Path No. : 72/25

Sex : Female

Age : 45 Y

Referring Doc :

Request No. : 13001787

Request Date : 06/01/2025

Verification Date : 16/02/2025

Financial Category : أورام دار السلام - ميانره

Department : Out Patient

Registered by : Mrs.Mayada Hegazy

Pathology Report

Diagnosis:

Right breast cancer post NACT, MRM specimen:

- * Residual viable invasive duct carcinoma G II with focal DCIS (5%).
- * Free nipple and areola.
- * Lymph nodes were positive in (4 / 9) with extranodal extension.
- * Free deep margin.
- * Partial pathological response to therapy.
- Stage: ypT1c, ypN2a.
- RCB: 3.418, RCB class III.

Signature

Dr. Marwa Mahmoud.

Dr. Duaa Mohammed.

Verified by :

DR.Menna Nabil

Printing Date : 19/02/2025 12:13:25

Page 2 of 2

Printing By : Mrs.Mayada Hegazy



غرفة رقم (4) - الدور الأول - مبنى رقم (987) - كورنيش النيل - مصر القديمة - القاهرة - مصر
Room No. (4) - First Floor-Building No. 987- Nile Corniche - Misr El Qadema - Cairo, Egypt



Patient Name: Sherine Ali Mohamed Gad	Date : 08/10/2024
Exam : C.T. ABDOMEN & PELVIS WITH CONTRAST	Age : 45 Year(s)
Medical ID: 3881/2022	Pat. ID: 18987

POST-CONTRAST MULTI-SLICE CT SCAN OF THE CHEST, ABDOMEN & PELVIS REVEALED:

- As compared to previous study dated 27/03/2024.

1. Chest:

• **Stationary course regarding the previously seen:**

- Right LIQ area of parenchymal distortion despite being partially out of field.
- Right axillary prominent level I and II lymph nodes, the largest measures 1.8 x 1.4 cm.
- Few tiny scattered mostly subpleural pulmonary nodules ranging between 2-3 mm.
- No pathological hilar or mediastinal lymph nodes.
- No evidence of pleural effusion, pleural thickening or masses seen.
- Mild cardiomegaly. No pericardial effusion detected.

2. Abdomen and pelvis:

- The liver is still seen mildly enlarged with slightly inhomogeneous CT texture. No hypodense hepatic focal lesions could be appreciated or contour irregularities.
- Patent homogeneously enhancing portal vein and its major branches as well as its main tributaries.
- No intra-hepatic or extra-hepatic biliary radicles dilatation.
- Normal size & parenchymal texture of the spleen with no focal lesions on non-contrast basis.
- No pathological abdominal or pelvic lymph nodes.
- Normal size, parenchymal thickness & density of both kidneys with no stones or back-pressure changes.
- Normal appearance of the pancreas and both supra-renal.
- Smooth filling of the urinary bladder with no filling defects or diverticular outpouching.
- Normal appearance of the other pelvic organs.
- No abdominal or pelvic collections.

3. The scanned skeleton:

- Stable multi-level vertebral bodies sclerotic lesions.
- No vertebral structural collapse or extra-osseous soft tissue component.

OPINION:

- Follow up study showing stationary course since the last study dated 27/3/2024.

Dr. Mostafa Abdelwahab, MSc

Dr. Wafid Mohamed, M.D.


			
رئيسة الجمهورية د. س. ح.		مباركة رئيسة الجمهورية لدعم صحة المرأة المصرية	
MDT FORM			
Name:	Shirin Ali Mohamed Jad	MRN:	1111
Age:	47 Y	ID:	27908030101701
Menopausal Sta		Phone:*	01140423136
NRS SCORE:		Marital Status:	
BMI:		Family History:	
Gender		Reason Of MDT:	
CO Morbidity:		progressive disease	
Surgical History:* mrm		ECOG Performance Status	☐ 0 ☒ 1 ☐ 2 ☐ 3 ☐ 4

Clinical Data:

BX:	2022: IDC GII		
ER:	8/8	PR: 5/8	HER2: +ve KI67: 40%

History:

Patient diagnosed Rt oligo metastatic breast cancer 10/2022
CTCAP: small bony lesion D8 MDT 12/2022: For Femara- Herceptin-
Zoladex 3 months as patient was refusing chemotherapy CT 2/2023: Non
conclusive HFL 14x15 mm at segment VIII MRI AP 4/2023: 9mm FL at
segment V likely cyst for follow up MRI DL spine 11/2023: L4 & L5 few
osseous lesions likely metastatic BS 3/2024: Free
Patient ended up receiving 27 cycles IV Herceptin last 8/2024 (full course)
+ Femara + Zoladex 12-2022 till 6/2025
11/2024 CTCAP: SD, BS: Free She underwent RT MRM 1/2025
Residual viable IDC GII è DCIS ypT1cN2b (4/9) free margins
MDT 3/2025: 6 TCPh + PORTH > AI + Zoladex She received only 6 cycles
TXT-carboplatin and no anti her2, last cycle 10/2025

[illegible]

Clinical Staging:

Summary Statement:

Pathological Staging:

100

Sonomammography:

CTCAP 9/2026: diffuse heterogeneous density of lower thoracic lumber vertebrae as well as iliac bone and sacrum likely metastatic + ant wedge fracture of L1,2 &4 (pathological)

Biomarker/Molecular

Mets Workup:

MRI DL spine 9/2026: Widely scattered marrow infiltrative lesions within almost all dorsal, lumbar and sacral vertebrae with no definitive extra osseous soft tissue component
+ L1,2 &4 ant wedge compression fracture likely pathological

Recommendations:

Downloaded from <http://ajphaphysiol.org/> at University of California, San Diego on June 11, 2015

MDT Outcome & Plan

for xeloda/ tykerb
zometa every 3 ms after dental assessment

SIGNATURE

bout:blank

اسم المريض: شيرين علي محمد	PM تاريخ الحالة: 10:20:35 10/09/2026
الطبيب المعالج:	كود أمر الشغل: 105365
كود المريض: 59798	النوع: Female

MULTISLICE POST CONTRAST CT SCAN OF THE ABDOMEN AND PELVIS:

TECHNIQUE: Multislice axial CT images of the abdomen and pelvis were obtained followed by multiplanar reconstruction of the source images. IV contrast was used.

FINDINGS:

****Reported patient with breast cancer.**

- The liver is average in size showing homogenous CT density and enhancement with no focal lesions.
- Patent and homogeneously enhancing portal vein.
- No intra or extra hepatic biliary radicles dilatation.
- Normal CT appearance of the spleen, pancreas, suprarenal glands, both kidneys, aorta and IVC.
- Regular filling of the urinary bladder showing no stones, masses or diverticulae.
- Unremarkable other pelvic organs.
- No abdominal or pelvic pathologically enlarged lymph nodes.
- No ascites.
- Scanned lung bases show middle lobar small consolidative opacity with parenchymal bands.
- Diffuse heterogenous density of the lower thoracic and lumbar vertebrae as well as the iliac bones and sacrum showing mixed lytic and sclerotic lesions likely metastatic. Anterior wedging of L1, L2 and L4.

OPINION:

- **Diffuse heterogenous density of the lower thoracic and lumbar vertebrae as well as the iliac bones and sacrum showing mixed lytic and sclerotic lesions likely metastatic.**
- **Anterior wedge fractures of L1, L2 and L4, likely pathological.**

Dr. Asmaa Ali, MD

اسم المريض: شيرين علي محمد	PM تاريخ حاله: 10:20:35 10/09/2026
الطبيب المعالج:	كود أمر الشغل: 105365
كود المريض: 59798	النوع: Female

MRI OF THE DORSAL AND LUMBAR SPINE

TECHNIQUE:

- Sagittal STIR, T1 and T2 WIs
- Axial T1 and T2 WIs

FINDINGS:

****Reported patient with breast cancer.**

- Widely scattered marrow infiltrative lesions are seen within almost all the dorsal, lumbar and sacral vertebrae. They appear of low T1 and heterogenous high T2 and STIR signal. No definite extraosseous soft tissue component.
- L1, L2 and L4 anterior wedging.
- Similar lesions are seen within both iliac bones.
- D4-5 central disc protrusion is seen effacing the central CSF space.
- L4-5 posterior disc bulge is seen indenting the theca and partially encroaching upon the neural foramina.
- Normal dimensions of the spinal canal.
- Normal size and signal of the spinal cord.

OPINION:

- **Reported patient with breast cancer shows widely scattered marrow infiltrative lesions as described, likely metastatic.**
- **L1, L2 and L4 anterior wedge compression fractures, likely pathological.**

Dr. Asmaa Ali, MD