

From Autoencoders to Variational Autoencoders: The Encoder

Issues with vanilla autoencoders

- The latent space plot isn't symmetrical around the origin
- Some labels are represented over small areas, others over large ones
- There are discontinuities in the latent space

From Autoencoders to Variational Autoencoders

- Modify encoder component
- Modify loss function


From Autoencoders to Variational Autoencoders

- Modify encoder component
- Modify loss function

Encoder mapping: AE vs VAE

Encoder mapping: AE vs VAE


Map sample to a point in latent space




*Images taken from *Generative Deep Learning* by David Foster

Encoder mapping: AE vs VAE

Map sample to a point in latent space



Map sample to a multivariate normal distribution in latent space



*Images taken from *Generative Deep Learning* by David Foster

WHAT THE HECK IS A

**MULTIVARIATE
NORMAL DISTRIBUTION?**

imgflip.com

Defining the normal distribution in 1d

- AKA Gaussian distribution

Defining the normal distribution in 1d

- AKA Gaussian distribution
- Probability distribution with bell curve shape

Defining the normal distribution in 1d

- AKA Gaussian distribution
- Probability distribution with bell curve shape
- Describe real-valued random variables (e.g., population height, weight)


Defining the normal distribution in 1d

- AKA Gaussian distribution
- Probability distribution with bell curve shape
- Describe real-valued random variables (e.g., population height, weight)
- Defined by 2 variables:
 - μ - mean value
 - σ - standard deviation

Defining the normal distribution in 1d

- AKA Gaussian distribution
- Probability distribution with bell curve shape
- Describe real-valued random variables (e.g., population height, weight)
- Defined by 2 variables:
 - μ - mean value -> centre of the distribution
 - σ - standard deviation -> variability of the distribution

Visualising the (standard) normal distribution



Probability density function

$$f(x|\mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

Defining the normal distribution in 1d

- The greater σ the flatter the curve
- Changing μ shifts the curve right and left
- [Let's play around with the normal distro!](#)

Sampling a point from a normal distribution

$$z = \mu + \sigma \varepsilon$$

Sampling a point from a normal distribution

$$z = \mu + \sigma \boxed{\varepsilon}$$


Sampled point from
standard normal distribution



**LET'S GENERALISE A NORMAL
DISTRIBUTION TO MORE THAN 1 DIMENSION**

MULTIVARIATE NORMAL DISTRIBUTION

Visualising the multivariate normal distribution



Formalising the multivariate normal distribution

$$f(x_1, \dots, x_k) = \frac{e^{-\frac{1}{2}(\vec{x} - \vec{\mu})^T \Sigma^{-1} (\vec{x} - \vec{\mu})}}{\sqrt{(2\pi)^k |\Sigma|}}$$

Formalising the multivariate normal distribution

$$f(x_1, \dots, x_k) = \frac{e^{-\frac{1}{2}(\vec{x} - \vec{\mu})^T \Sigma^{-1} (\vec{x} - \vec{\mu})}}{\sqrt{(2\pi)^k |\Sigma|}}$$

∝ Mahalanobis distance:

Distance between \mathbf{x} and $\boldsymbol{\mu}$

Formalising the multivariate normal distribution

$$f(x_1, \dots, x_k) = \frac{e^{-\frac{1}{2}(\vec{x} - \vec{\mu})^T \Sigma^{-1} (\vec{x} - \vec{\mu})}}{\sqrt{(2\pi)^k |\Sigma|}}$$

$$\vec{\mu} = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix}$$

Formalising the multivariate normal distribution

$$f(x_1, \dots, x_k) = \frac{e^{-\frac{1}{2}(\vec{x} - \vec{\mu})^T \Sigma^{-1} (\vec{x} - \vec{\mu})}}{\sqrt{(2\pi)^k |\Sigma|}}$$

$$\vec{\mu} = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix}$$

$$\Sigma = \begin{pmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{pmatrix}$$

Covariance
matrix

Formalising the multivariate normal distribution

$$f(x_1, \dots, x_k) = \frac{e^{-\frac{1}{2}(\vec{x} - \vec{\mu})^T \Sigma^{-1} (\vec{x} - \vec{\mu})}}{\sqrt{(2\pi)^k |\Sigma|}}$$

$$\vec{\mu} = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix}$$

$$\Sigma = \begin{pmatrix} \sigma_x^2 & \rho \sigma_x \sigma_y \\ \rho \sigma_x \sigma_y & \sigma_y^2 \end{pmatrix}$$

Covariance
matrix

Correlation
between x, y

VAE and multivariate normal distribution

- VAE assumes all dimensions are independent ($\rho = 0$)

VAE and multivariate normal distribution

- VAE assumes all dimensions are independent ($\rho = 0$)

$$\Sigma = \begin{pmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{pmatrix} \Rightarrow \begin{pmatrix} \sigma_x^2 & 0 \\ 0 & \sigma_y^2 \end{pmatrix}$$

VAE and multivariate normal distribution

- VAE assumes all dimensions are independent ($\rho = 0$)
- Encoder maps each input to a mean vector and a variance vector in the latent space

VAE and multivariate normal distribution


- VAE assumes all dimensions are independent ($\rho = 0$)
- Encoder maps each input to a mean vector and a variance vector in the latent space

$$\vec{\mu} = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix} \quad \vec{\sigma}^2 = \begin{pmatrix} \sigma_x^2 \\ \sigma_y^2 \end{pmatrix}$$

VAE and multivariate normal distribution

- VAE assumes all dimensions are independent ($\rho = 0$)
- Encoder maps each input to a mean vector and a variance vector in the latent space

$$\vec{\mu} = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix} \quad \vec{\sigma}^2 = \begin{pmatrix} \sigma_x^2 \\ \sigma_y^2 \end{pmatrix}$$



Encoding data into a point in latent space

$$z = \vec{\mu} + \Sigma \epsilon$$

Encoding data into a point in latent space

$$z = \vec{\mu} + \Sigma \boxed{\varepsilon}$$

Sampled point from
standard normal
distribution

Encoding data into a point in latent space

$$z = \vec{\mu} + \Sigma \epsilon$$

Sampled point from
standard normal
distribution

Encoding data into a point in latent space

$$z = \vec{\mu} + \Sigma \epsilon$$

Sampled point from
standard normal
distribution

$$\Sigma = e^{\frac{\log(\vec{\sigma}^2)}{2}}$$

Encoding data into a point in latent space

$$z = \vec{\mu} + \Sigma \epsilon$$

Sampled point from
standard normal
distribution

$$\Sigma = e^{\frac{\log(\vec{\sigma}^2)}{2}}$$

Log can take any
value in - / + infinity

**WHY SHOULD WE USE
MULTIVARIATE DISTRO?**



A major autoencoder drawback

There are discontinuities
in the latent space



Some generated images
will be poor

The multivariate normal distro fix

- We're sampling a random point around the mean value

The multivariate normal distro fix

- We're sampling a random point around the mean value
- Points in the same area produce very similar images when decoded

The multivariate normal distro fix

- We're sampling a random point around the mean value
- Points in the same area produce very similar images when decoded
- Reconstruction loss is small

The multivariate normal distro fix

- We're sampling a random point around the mean value
- Points in the same area produce very similar images when decoded
- Reconstruction loss is small
- VAE ensures a quasi-continuous latent space

The multivariate normal distro fix

- We're sampling a random point around the mean value
- Points in the same area produce very similar images when decoded
- Reconstruction loss is small
- VAE ensures a quasi-continuous latent space
- We can sample any point in the latent space, expecting the decoder to create a well formed image

What next?

- Discuss techniques to fix ulterior AEs' issues
- Focus on loss function for VAEs