


CE 641 FATE AND TRANSPORT MODELING OF ECOSYSTEMS

Tidal Exchange Mass Transport

1. A harbor of a large bay receives a discharge at one end as shown. The harbor is divided into two segments for simplicity.



A conservative substance is released into segment 1 at a rate of 1 kg/day. There is no advective flow through the system. A survey gave the following results:

$$C_1 = 12 \text{ mg/L}, \quad C_2 = 5 \text{ mg/L}, \quad \text{concentration of bay} = 1 \text{ mg/L}$$

- (a) What are the dispersive exchange flows due to mixing of the bay and the harbor, (i.e., between segments 1 and 2 and between segment 2 and the bay, respectively) in m^3/day ?
 - (b) What is the input to the bay from this harbor in kg/day ? That is, what is the dispersive flux across the interface between segment 2 and the bay?
2. As a first rough approximation, the lower 50 miles of the Potomac Estuary may be assumed to be composed of two well-mixed segments of water – one above the other – with flow entering the lower layer from the Chesapeake Bay. Mixing between the two layers also occurs. The following conditions are appropriate:

Inflow of fresh water to upper layer = 11,000 cfs

Salinity of inflow = 0

Salinity of upper layer = 10 ppt

Salinity of lower layer = 12 ppt

Salinity of bottom bay water = 14 ppt

Assuming that horizontal dispersion is not significant, but including vertical mixing:

- (a) What is the flow entering the lower layer of the estuary from the bay in cfs?
- (b) What is the surface outflow from the upper layer to the bay in cfs?
- (c) If the surface area is $7.68 \times 10^7 \text{ m}^2$ and each segment depth = 3 m, what is the vertical dispersion coefficient in cm^2/sec ?