

SUSPECT ADVERSE REACTION REPORT

CIOMS FORM I: CASE 3

I. REACTION INFORMATION			
1. PATIENT INITIALS (first, last) Not Reported	1a. COUNTRY United States	2. DATE OF BIRTH Not Reported	2a. AGE 54 Years
3. SEX Female	4-6. REACTION ONSET Unspecified (Signs developed following 5 days of consecutive administration prior to admission)		
8-12. CHECK ALL APPROPRIATE TO ADVERSE REACTION ☐ PATIENT DIED ☒ INVOLVED OR PROLONGED INPATIENT HOSPITALISATION (Required 13 days of acute inpatient care) ☐ INVOLVED PERSISTENCE OR SIGNIFICANT DISABILITY OR INCAPACITY ☒ LIFE THREATENING (Severe febrile neutropenia with ANC of 0.01 K/uL) ☒ OTHER MEDICALLY IMPORTANT CONDITION (Severe multi-systemic overdose toxicity)			
7 + 13. DESCRIBE REACTION(S) (including relevant tests/lab data) Severe iatrogenic methotrexate toxicity presenting as profound myelosuppression/pancytopenia, severe oral hemorrhagic sores on the hard and soft palate, significant dysphagia, aphonia, nausea, vomiting, severe diarrhea, hematochezia, extreme fatigue, generalized weakness, and erythematous papules on the buttocks and posterior lower limbs. An abdominal CT scan confirmed acute inflammatory mucositis and colitis. Lab Values: • Admission (Day 1): WBC: 0.7 K/uL, Hb: 11 g/dL, Platelets: 133 K/uL, Neutrophils: 12%, ANC: 0.09 K/uL, K+: 3.1 mmol/L, Creatinine: 0.59 mg/dL. • Nadir (Day 4/5/6): WBC: 0.5 K/uL, Platelets: 15 K/uL, ANC: 0.01 K/uL. • Serum Methotrexate: <0.04 mg/ml (expected in delayed/chronic tissue toxicity). • Discharge (Day 13): WBC: 7.5 K/uL, Platelets: 299 K/uL, ANC: 4.4 K/uL (fully normalized).			

II. SUSPECT DRUG(S) INFORMATION	
14. SUSPECT DRUG(S) (include generic name)	Methotrexate (2.5 mg tablets)
15. DAILY DOSE(S)	15 mg daily (3 tablets of 2.5 mg twice a day due to medication administration error)
16. ROUTE(S) OF ADMINISTRATION	Oral
17. INDICATION(S) FOR USE	Polyarthritis secondary to mixed connective tissue disease

18. THERAPY DATES (from/to)	Start: Unspecified (Administered for 5 consecutive days prior to admission) Stop: Discontinued after 5 days upon identification of error
19. THERAPY DURATION	5 days
20. DID REACTION ABATE AFTER STOPPING DRUG?	☒ YES ☐ NO ☐ NA (Positive Dechallenge: Hematological indices and mucosal lesions gradually normalized following drug withdrawal and leucovorin rescue).
21. DID REACTION REAPPEAR AFTER REINTRODUCTION?	☐ YES ☐ NO ☒ NA (Rechallenge not performed).

III. CONCOMITANT DRUG(S) AND HISTORY

22. CONCOMITANT DRUG(S) AND DATES OF ADMINISTRATION (exclude those used to treat reaction)
• Mesalamine (Stopped during admission) • Budesonide (Stopped during admission)
23. OTHER RELEVANT HISTORY (e.g. diagnostics, allergies, pregnancy with last month of period, etc.)
• Medical History / Comorbidities: Mixed connective tissue disease, ulcerative colitis, hypertension, Lyme disease, and sick sinus syndrome.

IV. MANUFACTURER INFORMATION

24a. NAME AND ADDRESS OF MANUFACTURER Unknown (Literature report)	24b. MFR CONTROL NO. Unknown (Literature report)	24c. DATE RECEIVED BY MANUFACTURER Unknown (Literature report)
24d. REPORT SOURCE ☐ STUDY ☒ LITERATURE ☒ HEALTH PROFESSIONAL	DATE OF THIS REPORT 19-Jul-2026	25a. REPORT TYPE ☒ INITIAL ☐ FOLLOWUP
Literature Citation Details: • Authors: Kristina L. Quan Soon, Sundeep Shah, Ramy Ibrahim, Khalid Alzwahereh • Journal: Cureus (July 2024) • Article ID: Cureus 16(7): e64081 • DOI: https://doi.org/10.7759/cureus.64081		