Tenth-discriminant bounds compared: a verified survey of the methods in Tao's challenge, and an analytic cutoff $q < 548$ from the determinant method

Peter W. Murphy

October 2026 (v1.4.0)
Repository https://github.com/DrMurphyIsIn/tenth-discriminant, directory `comparison/`
DOI https://doi.org/10.5281/zenodo.23690870
(all versions: https://doi.org/10.5281/zenodo.23550257)

Status: certified by interval arithmetic; reproduced by a separately written implementation; **not kernel-checked**

Abstract

Terence Tao's tenth-discriminant challenge asks, for each recent argument excluding Landau–Siegel zeros or establishing a quasi-Riemann hypothesis, for the effective upper bound D_0 it gives on a hypothetical tenth imaginary quadratic field of class number one, so that the strength of the arguments can be compared. The comment thread now holds about twenty such extractions. We give, first, a survey that puts them side by side: each value is quoted at source with its author and comment, and labelled by what we could check. For every input that can be written as a real-zero-free segment we use one reduction, the Chowla–Selberg expansion of the principal form, and certify in ball arithmetic: $d \le 652$ from the 7/8 half-plane, $d \le 1151$ from the 11/12 half-plane (LemmaLab's value, comment 698473), $d \le 199$ under GRH, with the limit $64\pi^2 e^{-2\gamma} = 199.1208\ldots$, $d < 10^{10}$ from Banks's explicit theorem (as twinheterodox and michaelmross posted), and, from Kimura's zero-gap theorems (Lean-formalised by the author; not replayed by us), $d < 6.47 \cdot 10^7$ (his chain theorem 58) and $d < 1.23 \cdot 10^8$ (the input of Rudolph's extraction). Reading the same curve backwards puts every method on one scale, the zero-free abscissa it is "worth": the light-machinery determinant bound $q < 570$ of twinheterodox corresponds to 0.861, ahead of the 7/8 half-plane on this problem. Second, we rebuild that determinant bound (v0.6.1 of twinheterodox, after Green, van Dijk, Pratt and Kerr) from our own derivation: two separately written AI implementations reproduce its certificate row by row, to 10^{-13}, in its own configuration, and the method passes a truth test on the fields of discriminant -163, -67 and -43. Third, we improve it: an explicit treatment of the Poisson modes in the norm series, the sharper Rosser–Schoenfeld constant, re-optimised sieve rows and kernels, and a band-uniformity check by Taylor models. Together with the kernel-checked bound $d \le 652$ of our earlier work, this gives the analytic cutoff $h(K) = 1 \Rightarrow q < 565$ with Rosser–Schoenfeld's 1962 prime estimates. Fourth, with modern explicit bounds for the Chebyshev function (Broadbent–Kadiri–Lumley–Ng–Wilk; Büthe 2018), read at source, double-transcribed and tested against exact prime sums to 10^9, the auxiliary-prime intervals shrink and the cutoff becomes $h(K) = 1 \Rightarrow q < 548$, worth $\sigma_{\mathrm{eq}} = 0.8566$ on the scale above. The total improvement over 570 is about 3.9% and changes no candidate discriminant; it is a measure of the method's strength, which is what the challenge compares. The computations are certified by interval arithmetic and reproduced by an independent implementation; they are *not* kernel-checked.

1 Introduction

1.1 The challenge

There are nine imaginary quadratic fields of class number one, with discriminants -3, -4, -7, -8, -11, -19, -43, -67, -163. That there is no tenth was proved by Heegner [7], with later proofs by Baker [1] and Stark [20]. Tao's challenge [21] takes the recent effective arguments against Landau–Siegel zeros, and the recent quasi-Riemann hypotheses, and asks what bound each gives on a *hypothetical* tenth discriminant: "if a tenth existed, then $d \le D_0$". The answer is known; the numbers measure the arguments. Tao asks for every approach's value, "even if they are worse", and for negative results.

1.2 What this paper does

1. **A survey** (Section 4). Every posted extraction, quoted at source with comment number and author, in one table, with a status per row: reproduced independently by us, cited, or not reproducible. For the zero-free inputs we use one reduction (Section 3) so that rows compare methods rather than reductions, and we read the resulting curve backwards to obtain a single strength scale (Section 4.3).

2. **A reproduction** (Section 5) of the strongest posted light-machinery bound, $q < 570$ [22], from our own derivation and code.

3. **An improvement** (Sections 6 and 8) of that bound to $q < 565$ with the 1962 prime estimates, and to $q < 548$ with modern explicit bounds for the Chebyshev function (Section 7).

1.3 How the claim is phrased

The theorem of Section 8 is an *analytic cutoff*: an argument that excludes every $q \ge 548$ without a finite search above 548. The finite range below the cutoff is checked by computation (Section 8.1). Above 652 the argument is our kernel-checked theorem $d \le 652$ [12], which uses OpenAI's 7/8 half-plane [15]; between 548 and 652 it is the determinant method alone. So the result belongs to the class "unconditional, Heegner-free, with a quasi-Riemann input". Since a light-machinery bound is also valid in that class, the comparison with 570 is made inside it.

What is verified, and how. Every inequality of the certificate is evaluated in Arb ball arithmetic [8]. A second implementation, written separately by an independent AI review session in high-precision floating point, reproduces the certificate band by band. Both implementations are AI-written; no human has checked them. **Nothing in Sections 5, 6 and 8 is kernel-checked.** The reason is given in Section 9.

2 Conventions

Throughout, $D < 0$ is a fundamental discriminant, $q = d = |D|$, $K = \mathbb{Q}(\sqrt{D})$, $\chi = \chi_D$. All bounds D_0 are upper bounds: $h(K) = 1 \Rightarrow q \le D_0$ (or $< D_0$, as stated). Lower bounds on a hypothetical tenth discriminant, such as $q > \exp(2.2 \cdot 10^7)$, which Pratt (comment 698460) attributes to Stark's 1966 computation of low-lying zeros, are finite-range tools and are not compared here.

Two elementary consequences of $h(K) = 1$ are used repeatedly; they are classical, and were emphasised in the thread by Silberman (comment 698449) and Pratt (comment 698460). Every prime $p < q/4$ is inert, because a non-rational element of $\mathcal{O}_K$ has norm at least $q/4$. Hence, for $q > 8$, q is a prime and $q \equiv 3 \pmod 8$.

3 One reduction for the zero-free inputs

For $h(K) = 1$ and $q > 4$, $\zeta_K(s) = \frac{1}{2} Z_Q(s)$ for the principal form Q, and Chowla–Selberg [19] gives

$$\zeta_K(s) = \zeta(2s) + 2^{2s-1}\sqrt{\pi}\,\frac{\Gamma(s-\frac{1}{2})}{\Gamma(s)}\,\zeta(2s-1)\,q^{\frac{1}{2}-s} + E(s), \tag{1}$$

with the Bessel term E bounded by $2^{s+3/2}\pi^s\Gamma(s)^{-1}q^{1/4-s/2}\sum_n n^{s-1/2}\sigma_{1-2s}(n)K_{s-1/2}(\pi n\sqrt{q})$. We checked the normalisation against direct lattice sums before using it. In the thread this expansion was written down by michaelmross (comment 698466) and evaluated at $s = 7/8$ by LemmaLab (comment 698472). Write $M(\sigma, q)$ for the sum of the first two terms and $d_M(\sigma)$ for the root in q of $M(\sigma, q) = 0$, which has the closed form $d_M(\sigma) = \big(-C(\sigma)\zeta(2\sigma-1)/\zeta(2\sigma)\big)^{1/(\sigma-1/2)}$.

Proposition 3.1 (the edge form)**.** *If $L(s,\chi)$ has no real zero in $(\sigma_0, 1)$, $\frac{1}{2} < \sigma_0 < 1$, and $h(K) = 1$, then $q < d_M(\sigma_0)$ (at $\sigma_0 = 7/8$ this gives $d \le 652$, since -653 is not a discriminant: $653 \equiv 1 \pmod 4$). The certificate is $M(\sigma_0, D) - B(\sigma_0, D) > 0$ at $D = \lceil d_M(\sigma_0)\rceil$, with M increasing and B decreasing in q.*

Proposition 3.2 (zero forcing; twinheterodox, comment 698456, and michaelmross, comment 698466)**.** *If $h(K) = 1$ and $q \ge 10^{10}$, then $L(s,\chi)$ has a real zero β with $1 - \beta < 1.92/\sqrt{q}$. More generally $\kappa(q_1)$ is certified for $q \ge q_1$, with $\kappa \to 6/\pi$.*

The route is twinheterodox's (a theta integral, with $1 - \beta \le 10/\sqrt{q}$) and, independently, michaelmross's (the Chowla–Selberg expansion, with $2/\sqrt{d}$ and the asymptotic constant $6/\pi$); the constant 1.92 for $q \ge 10^{10}$ is ours. The proof uses $\zeta(u) > 1/(u-1) + \frac{1}{2}$ on $(0, 1)$ (first-order Euler–Maclaurin) and the sign change of ζ_K near $s = 1$. For an input in the normal form "$(1-\beta)\log q \ge c$ for $q \ge q_0$", Proposition 3.2 gives $D_0 = \max(q_0, d_c)$, with d_c the point where $\kappa \log x/\sqrt{x}$ falls below c; for a piecewise constant $c(\log q)$, D_0 is the least d beyond which $\kappa \log x/\sqrt{x} < c(\log x)$ for all x.

4 The survey

4.1 The table

Table 1 lists the extractions in the thread of [21], read from the page on 25 October 2026 (74 comments), with the formulas taken from the page's image text. Status: **R** = reproduced independently by us (and by the review session's own code); **C** = cited as posted, not yet checked by us; **NR** = not reproducible from what is posted.

4.2 The Kimura rows

Kimura's Lean development exports, besides Theorem 1, chain theorems `chain_i : B4.GapFrom c_i l_i` meaning $(1-\beta)\log q \ge c_i$ for $\log q \ge l_i$ [10]. Through Proposition 3.2, `chain_58` = (2147/500000, 88711/5000) gives $d < 64{,}668{,}273$ and is the best of the 313; `chain_59`, the input of Rudolph's extraction, gives $d < 123{,}209{,}905$, against $1.89 \cdot 10^8$ by his Proposition 9 route. Both values are certified by us and rest on Kimura's Lean-formalised theorems, which we have not replayed. Rudolph's own value $1.89 \cdot 10^8$ comes from a different route (his Proposition 9 numerics), which we have not re-derived.

4.3 One scale

Proposition 3.1 is a curve $\sigma_0 \mapsto d_M(\sigma_0)$, increasing on the sampled range from 199.12 at $\sigma_0 \to \frac{1}{2}$ to 653.99 at 7/8 and 1151.66 at 11/12. Reading it backwards assigns to any D_0 the abscissa $\sigma_{\mathrm{eq}}(D_0)$ that a real-zero-free segment would have to reach to give the same D_0. This is a calibration device, not a statement that a method proves anything about zeros.

Two observations follow. On this problem the light determinant method is worth more than the 7/8 half-plane. And within the zero-free family, a bound below 570 would need a real-zero-free segment beginning at 0.8610 uniformly for $q \geq 570$, which no unconditional result supplies. Classification-based routes (Weber's, as reconstructed in twinheterodox's historical companion, and Proposition 1.10 of [3], whose proof for odd characters appeals to Watkins's tables) are outside the comparison.

5 Reproducing the determinant bound $q < 570$

The bound is version 0.6.1 of twinheterodox [22]. In their own account it builds on Green's derivative determinant and Frobenius divisibility [6], van Dijk's auxiliary-inert-prime nonvanishing [23], Pratt's suggestions (comments 698476, 698460), Kerr's higher Frobenius relations [9], the Poisson expansion of michaelmross and Tao, and Rosser–Schoenfeld's prime estimates [18]. Their repository carries no licence; we use the mathematics, with citation, and our own words, code and tables.

5.1 The argument in outline

Suppose $h(K) = 1$ with q in a band $[Q_0, Q_1]$. The argument compares two bounds for the same quantity, a smoothed sum over the noninert primes,

$$T(X) = \sum_{p \text{ noninert}} \frac{\log p}{p}\, f(p/X), \qquad f(u) = 2u \sum_{m \geq 1} (1 - mu)_+,$$

which vanishes for $X \leq q/4$.

The lower bound. For a scale U, counting lattice points of $\mathcal{O}_K$ in an annulus, with a Bonferroni sieve over a few small ideals, shows that the split primes cannot carry all the prime weight in $(U, \kappa U]$, so an inert prime p_0 lies there. With $R = p_0$, Green's derivative interpolation matrix, with rows $z^r w^s$ ($r < R$, $s < tR$) and columns the k-th z-derivatives ($k < R/9$) at the lattice points of a disk, has full row rank, by reduction modulo p_0, where Frobenius sends z to $\bar{z}$ (van Dijk's argument). Fix a nonzero maximal minor D and put $\Delta = |D|^2 \in \mathbb{Z}_{>0}$. At each inert prime p, triangular row changes by $z^p - w$, $z^{p^2} - z$ and $w^{p^2} - w$ make every entry of a row divisible by a power of p; summed over primes this is a lower bound for $\log \Delta$, whose main term is a sum over all primes minus $T(R)$. Factorials at the split primes and in the columns add further divisibility, again expressed through T at smaller scales. An entrywise estimate bounds $\log \Delta$ from above. Comparing the two forces $T(R) \geq \alpha \log R - \frac{\beta}{4} \log q - K$, and, since T is monotone, a piecewise affine lower bound $T(e^x) \geq L(x) - \gamma \log(q/Q_0)$ for x across many scales, refined in eight stages that each feed the previous stage's bound back into the factorial gains.

The upper bound. Under $h(K) = 1$, $\zeta_K(s) = \zeta(2s) + A_q(s)$ with A_q the norm series of the principal form, so $-\frac{d}{ds} \log(\zeta_K(s)/\zeta(2s))$ is a Dirichlet series over the split primes and their powers. For a positive kernel $h(x) = Z^{-1} e^{-dx} P(e^{-ex})^2$, the average $\int h(x)\, T(e^x)\, dx$ is a combination of that series at the real points $s = 1 + d + je$, $j = 0, 1, \ldots$, which Poisson summation in one coordinate of the norm form evaluates up to an explicit error.

The contradiction. If $\int h\,L >$ the upper bound at $q = Q_0$, and the upper bound decreases in q faster than $\gamma \log(q/Q_0)$ across the band, then no q in the band has $h(K) = 1$.

5.2 The reproduction

We derived each step independently and wrote certificate code that reads only v0.6.1's parameters. Our numbers were frozen and their checksums sent to the review session before any comparison. Three slips in our first code (two conservative, one not) were found by this comparison and fixed by derivation; none was an error in v0.6.1. With v0.6.1's parameters and its Mertens constant 1.334 (the faithful configuration), our per-row envelope coefficients agree with theirs to within $6.3 \cdot 10^{-14}$ in each of the three bands we use, consistent with their outward rounding, and the lower bound of the band $[570, 580]$ is 10.9754131603100 against their 10.975413160309. The review session's separately written implementation agrees per row.

5.3 Controls

- **Truth test.** At $q = 163$, 67 and 43, which have class number one, every inequality is a theorem. Our envelope lies below each field's actual noninert-prime sum, computed from its split primes up to 10^8, at all nine stages (margin at least 1.9; with the modern inputs of Section 7 and the $[548, 550]$ band data, at least 1.7). The Chowla–Selberg main term agrees with the actual weighted prime sum at 163 to about 0.1% (approximate: the tail beyond 10^8 is estimated).

- **Expected failure.** With v0.6.1's parameters the band $[560, 570]$ fails (gap -0.026), consistent with their record that kernel searches at 560 "did not yield a positive gap".

- **Degradation.** On the band $[565, 570]$, worsening an input at first order (Rosser–Schoenfeld's E or Table I by 1%, or one row's κ by 1%) lowers the gap; likewise on $[548, 550]$ for each modern input and for one row's κ.

- **Quadrature.** The kernel average computed through the cancelling exponential expansion agrees with a direct quadrature of the positive kernel to $2 \cdot 10^{-15}$.

6 The improvements

1. **The Poisson modes.** In the norm series $A_q(s) = \sum_{b\geq 1} \sum_{a\in\mathbb{Z}} ((a + b/2)^2 + qb^2/4)^{-s}$ the nonzero Poisson modes are

$$A_q(s) - F_q(s) = \sum_{b,n\geq 1} 4(-1)^{nb} \pi^s n^{s-1/2} (bv)^{1/2-s} K_{s-1/2}(2\pi nbv)/\Gamma(s), \qquad v = \sqrt{q}/2,$$

   (the sign checked against lattice sums). Bounding them termwise with $K_\nu(x) \leq \sqrt{\pi/(2x)}\, e^{-x+\nu^2/(2x)}$, sharp to 0.2% here, instead of by a contour shift, reduces the error that multiplies the kernel's (large, cancelling) coefficients by a factor of 10^3–10^4. This admits kernels of degree 20 instead of 16. It is load-bearing: with contour-shift accounting (our derivation of it, with $a_0 = 1$) the new band fails by 41.6.

2. **The constant in $\sum_{p\leq x} \log p/p$.** Rosser–Schoenfeld's (3.21) gives $\log x + E - 1/(2\log x)$ for all $x > 1$, with $E = -1.3325822757\ldots$; v0.6.1 uses 1.334 for $x \geq 319$. This adds about 0.0016 to the margin.

3. **Our own band data.** For each band we choose the auxiliary-prime rows (sieve sizes and the minimal κ certified by the sieve inequality), the parameter γ (only $\gamma \geq \beta/4$ is needed), and the kernel, as the top generalised eigenvector of the quadratic form that the kernel average defines, rounded to integers. All band data (rows, κ, sieve sizes, kernels, γ) are ours; the method constants $t = 3/64$, $C = 27/64$, $\tau = 3/10$ and the eight stages are v0.6.1's.

4. **Band uniformity.** The upper bound's derivative in $Y = \log(q/Q_0)$ is enclosed by a Taylor model whose coefficients are exact point evaluations, with a Cauchy-estimate remainder; the condition $U'(Y) < -\gamma$ makes the gap at Q_0 persist across the band.

5. **A certified constant.** $K(1/9) = \int_0^{1/9} I(k)\,dk$, for which v0.6.1 gives a closed form, is bracketed rigorously, $K(1/9) \in [0.0077989, 0.0078016]$, from certified values of $I(k)$ and its monotonicity (envelope theorem), and the upper end is used.

7 Modern explicit bounds for the Chebyshev function

The determinant method consumes explicit prime estimates in three places: the auxiliary-prime sieve, which shows that an inert prime lies in $(U, \kappa U]$; the envelope constants; and the sum $\sum_{p \leq x} \log p/p$. In the sieve the lower bound for $\vartheta(\kappa U) - \vartheta(U)$ must beat an upper bound for the split primes, so the width κ that the sieve can certify is set by the error in the prime estimate. With Rosser–Schoenfeld 1962, κ is about 1.42 at 2^{18} and about 1.1 at 2^{50}. Replacing every prime slot by the idealisation $\vartheta(x) = x$ shows the ceiling of the method in this form: the band $[560, 565]$, which fails with the 1962 inputs, then has a gap of $+0.047$ at Q_0 (an exploratory computation: $\vartheta(x) = x$ is not a bound), and the binding constraint moves from the primes to the sieve: at $[548, 550]$ a 1% change in one row's κ moves the gap by $6 \cdot 10^{-6}$, a 1% change in a prime input by about $5 \cdot 10^{-8}$.

We use the following, read at source from the arXiv versions and transcribed twice, independently; every printed value is raised by one unit in its last digit, since the printed values are rounded in the last digit [4, p. 26].

- $\vartheta(x) \leq (1 + 1.93378 \cdot 10^{-8})\,x$ for all $x \geq 0$ [4, Cor. 2.1].
- $\vartheta(x) - x > -\mathcal{D}_k(a,b)\,x/(\log x)^k$ on the intervals $[a,b) \subset [1, 7 \cdot 10^{11})$ of [4, Table 13], a direct computation using no information about zeros; and $(1 - m_0)x < \vartheta(x)$ for $x > X_0$ from [4, Table 14].
- $x - \vartheta(x) \leq 1.95\sqrt{x}$ for $1427 \leq x \leq 10^{19}$ and $\vartheta(x) < x$ for $x \leq 10^{19}$ [5, Thm. 2, (1.6)–(1.7)], which use the zeros of ζ up to height 10^{11}.
- $|\vartheta(x) - x| < 3.7979 \cdot 10^{-5}\,x/(\log x)^2$ for $x > 10^{19}$ [4, Table 15], which closes the tail.

The sum $\sum_{p \leq x} \log p/p$ then follows by partial summation, $\sum_{p \leq x} \log p/p = \log x + E + R(x)/x - \int_x^\infty R(t)\,t^{-2}\,dt$ with $R = \vartheta - t$: the integrand has a sign on $[x, 10^{19}]$, and the tail bound is integrable. These inputs are of the same two kinds as Rosser–Schoenfeld's own (computed prime data, and the zeros verified to some height: Lehmer's 25,000 in their case), at a much larger scale.

Checks. Every transcribed bound was tested against $\vartheta(x)$ computed exactly up to 10^9, at the points where each inequality is tightest. Two printed statements in the arXiv versions do not hold as printed, and we use corrected forms: $x - \vartheta(x) \leq 1.95\sqrt{x}$ fails on $[1423, 1427)$ (at $x = 1426.999$, $x - \vartheta(x) = 74.18 > 73.66$), so we start it at 1427; and the entry $\mathcal{D}_4(10^8, 10^9) = 17.4450$ of [4, Table 13] is below the true supremum 17.555, so we do not use it. Büthe's lies below the scales the method uses (from 2^{18}); the Table 13 entry lies within them, and we exclude it and use the other entries of that row. The published versions were not checked.

Result. With these inputs the bands $[548, 565]$ certify (Table 3); $[547, 550]$ does not. The binding margin, $+0.0013$ on $[548, 550]$, is small, and the next prime with $q \equiv 19 \pmod{24}$, 547, is not reached.

8 The theorem

Theorem 8.1. *If $h(\mathbb{Q}(\sqrt{D})) = 1$ for a fundamental discriminant $D < 0$, then $|D| < 548$. The argument excludes $[548, 565]$ by five certified bands of the determinant method using the explicit bounds of Section 7, $[565, 652]$ by four certified bands using Rosser–Schoenfeld's 1962 estimates, and $|D| \geq 653$ by [12].*

8.1 The finite range

For every fundamental discriminant with $3 \leq q < 548$ (there are 166) we compute h directly as the number of reduced forms. It is 1 exactly for $q \in \{3, 4, 7, 8, 11, 19, 43, 67, 163\}$.

8.2 Size and fragility

The cutoff moves from 570 to 565 with the 1962 inputs, about 0.9%, and to 548 with the modern inputs, about 3.9% in all. No candidate changes: the primes $q \equiv 19 \pmod{24}$ near the cut are 547 and 571, and both are excluded either way. With the 1962 inputs the binding band $[565, 570]$ has margin 0.0023, smaller than the effect of a 1% change in Rosser–Schoenfeld's constant E (about 0.015), and it certifies only with the Poisson-mode treatment of Section 6. With the modern inputs the binding band $[548, 550]$ has margin 0.0013; there the prime inputs are saturated (a 1% change moves the gap by about $5 \cdot 10^{-8}$), and the margin is set by the sieve and by the kernel.

9 Limits

- **Not kernel-checked.** The argument needs explicit prime estimates of Rosser–Schoenfeld quality, or better, at scales up to about 10^{16}; the modern ones of Section 7 rest on computations (prime sums to $7 \cdot 10^{11}$; zeros to height 10^{11}) that cannot be redone in a proof kernel. The explicit-bounds development in PrimeNumberTheoremAnd (at commit `c39a7511`, 30 September 2026) formalises several such results but rests them on trusted numerical inputs; its Rosser–Schoenfeld files prove the asymptotic Mertens theorems, while their explicit statements are stated with `sorry` (e.g. Theorem 12) or rest on files that contain `sorry`. No three-axiom-clean source of these inputs exists in Lean yet. A kernel-checked version is future work.

- **A correction to v1.3.0.** Version 1.3.0 used Rosser–Schoenfeld's 17-digit approximation of the constant E in $\sum_{p \leq x} \log p/p = \log x + E + o(1)$, $-1.33258\,22757\,33220\,87$ (p. 76), as a ball of radius 10^{-17}. The true value, computed independently by two series evaluations, is $E = -1.33258\,22757\,33220\,88176\ldots$, which lies $1.8 \cdot 10^{-18}$ outside that ball. Version 1.4.0 uses their correctly rounded 15-digit value (2.11), $-1.33258\,22757\,33221$ (p. 65), with radius 10^{-15}. The ball's midpoint moves by $1.3 \cdot 10^{-16}$, so every certified gap changes by about 10^{-16}, far below the gaps and the interval radii; no conclusion changes, and all nine bands are re-certified with the corrected input.

- **The survey is incomplete.** Rows marked C are cited, not checked.

- **Two implementations, both AI-written.** The reproduction is by two separately written AI implementations (ball arithmetic, and high-precision floating point); no human has checked either.
- **The inputs.** Rosser–Schoenfeld 1962 (Table I; Theorems 4, 6, 9, 14) is cited, read from the page images, and transcribed by us, for $[565, 652]$; the modern bounds of Section 7 [4, 5], double-transcribed and tested against exact prime sums to 10^9, for $[548, 565]$; OpenAI's 7/8 theorem enters only through [12] for $q \geq 653$.

10 Credits, provenance and AI use

The determinant method and the bound 570 are twinheterodox's [22], with the credits given there. The zero-forcing route is twinheterodox's and michaelmross's; the evaluation at the edge is LemmaLab's; the 11/12 value 1151 is LemmaLab's; the Kimura extraction route is Rudolph's; the inert-prime facts are classical, as emphasised by Silberman and Pratt. The explicit bounds for ϑ of Section 7 are those of Broadbent, Kadiri, Lumley, Ng and Wilk [4] and of Büthe [5]; Platt and Trudgian [17] first showed $\vartheta(x) < x$ to $1.39 \cdot 10^{17}$. Our contributions are the survey and its verification, the strength scale, the reproduction, and the improvements listed in Section 6.

This work was carried out with Claude (Anthropic) in Claude Code: derivations, code and text were AI-generated, and every result was reviewed by a separate AI session that wrote its own implementation. These are AI reviews, not human peer review. The author directed the work and takes responsibility for the claims.

References

[1] A. Baker. Linear forms in the logarithms of algebraic numbers. *Mathematika*, 13:204–216, 1966. doi:10.1112/S0025579300003971.

[2] William D. Banks. On the OpenAI proof that Landau–Siegel zeros do not exist, 2026. arXiv:2610.15483v1; title checked against our local copy.

[3] Michael A. Bennett, Greg Martin, Kevin O'Bryant, and Andrew Rechnitzer. Explicit bounds for primes in arithmetic progressions. *Illinois J. Math.*, 62:427–532, 2018. Journal reference as given at arXiv:1802.00085.

[4] Samuel Broadbent, Habiba Kadiri, Allysa Lumley, Nathan Ng, and Kirsten Wilk. Sharper bounds for the Chebyshev function $\theta(x)$, 2021. arXiv:2002.11068v2 (27 Jan 2021); journal version not checked.

[5] Jan Büthe. An analytic method for bounding $\psi(x)$, 2018. Math. Comp. 87 (2018), no. 312, 1991–2009; read from arXiv:1511.02032v2 (published version not checked).

[6] Ben Green. On Landau–Siegel zeros, 2026. hexagon:2610.00311v2, https://hexagonmath.org/pdf/2610.00311; title checked against our local copy of v2.

[7] Kurt Heegner. Diophantische Analysis und Modulfunktionen. *Math. Z.*, 56:227–253, 1952. doi:10.1007/BF01174749.

[8] Fredrik Johansson. Arb: efficient arbitrary-precision midpoint-radius interval arithmetic, 2016. arXiv:1611.02831 (title checked at arXiv); used through python-flint 0.6.0.

[9] Bryce Kerr. A simpler proof that there are no Landau–Siegel zeros, 2026. arXiv:2610.21099v1; title checked against our local copy.

[10] Iwao Kimura. Imaginary quadratic fields of class number at most 2000, 2026. arXiv:2610.21511v2 (pdf md5 d2dadf89); the Lean files and computations are in the author's release cln-imq-2k v1.4.

[11] Henric Larsson. An explicit class-number-one cutoff from recent real-zero gaps, 2026. https://github.com/henlar1975/Tao-tenth-discriminent-bound, commit 834664b4; title checked at paper.tex:8.

[12] Peter W. Murphy. The class number one theorem without Heegner, kernel-checked: $d \leq 652$ from the 7/8 quasi-Riemann hypothesis in Lean 4, 2026. Version 1.0.1, https://github.com/DrMurphyIsIn/tenth-discriminant, doi:10.5281/zenodo.23550992 (all versions: doi:10.5281/zenodo.23550257).

[13] Peter W. Murphy. Class number one without L-functions: a kernel-checked determinant proof, 2026. Version 1.1.0, https://github.com/DrMurphyIsIn/tenth-discriminant, doi:10.5281/zenodo.23563656, record https://zenodo.org/records/23563656.

[14] OpenAI. The quasi-Riemann hypothesis, 2026. Manuscript, 5 October 2026, openai/math `preprints/The-Quasi-Riemann-Hypothesis-October-5-2026/paper2.pdf` (path checked in our local copy; zero-free half-plane $\operatorname{Re} s > 11/12$). Not refereed.

[15] OpenAI. The quasi-Riemann hypothesis: a zero-free half-plane $\operatorname{Re} s > 7/8$, 2026. Manuscript, openai/math family 003, `preprints/The-Quasi-Riemann-Hypothesis-September-30-2026`, with a Lean proof under `lean/OAI/NumberTheory/DirichletL`. Not refereed.

[16] OpenAI. Uniform exclusion of Landau–Siegel zeros, 2026. Preprint, 1 October 2026, 9 pp.

[17] David J. Platt and Timothy S. Trudgian. On the first sign change of $\theta(x) - x$, 2016. Math. Comp. 85 (2016), no. 299, 1539–1547; read from arXiv:1407.1914v1 (published version not checked).

[18] J. Barkley Rosser and Lowell Schoenfeld. Approximate formulas for some functions of prime numbers. *Illinois J. Math.*, 6:64–94, 1962. Read from the page images of the PDF supplied by the author of this paper; DOI checked.

[19] A. Selberg and S. Chowla. On Epstein's zeta-function. *J. Reine Angew. Math.*, 227:86–110, 1967. doi:10.1515/crll.1967.227.86.

[20] H. M. Stark. A complete determination of the complex quadratic fields of class-number one. *Michigan Math. J.*, 14:1–27, 1967. doi:10.1307/mmj/1028999653.

[21] Terence Tao. An experimental AI-math challenge: using tenth discriminant bounds to explore arguments ruling out Siegel zeroes, 2026. Blog post, 20 October 2026, https://terrytao.wordpress.com/2026/10/20/an-experimental-ai-math-challenge-using-tenth-discriminant-bounds-to-explore-arguments-ruling-out-siegel-zeroes/.

[22] twinheterodox. Tenth-discriminant bounds: A smoothed Frobenius determinant bound for the tenth discriminant, version 0.6.1 ($q < 570$), and companions, 2026. GitHub repository https://github.com/twinheterodox/tenth-discriminant, commit 44c7d6fc, prepared with Codex; title checked against proof.tex of v0.6.1.

[23] Roelof van Dijk. A Frobenius determinant proof of the non-existence of Landau–Siegel zeros, 2026. hexagon:2610.00409v2, https://hexagonmath.org/pdf/2610.00409v2; title checked against our local copy.

Input	Source (comment)	Posted value	St.	Our value / note
GRH (conditional)	twinheterodox 698514; Stopple 698534	$q \le 200$	R	$d \le 199$; limit $64\pi^2 e^{-2\gamma}$
7/8 half-plane [15]	Tao 698471; LemmaLab 698472	655; 653	R	$d_M = 653.98763\ldots$, $d \le 652$, kernel-checked [12]
11/12 half-plane [14]	Tao 698471; LemmaLab 698473	1475; 1151	R	$d \le 1151$ (edge form)
Banks [2]	twinheterodox 698456; michaelmross 698466	10^{10}	R	$d < 10^{10}$, margin 6.7
Banks + Lu–Zaman–Zhao	Larsson 698469 [11]	50,000	C	
Inert-mass form	Silberman 698449/698451	10^{45}	C	source theorem unstated
Kerr [9]	Bryce 698475, who calls arXiv:2610.21099 (B. Kerr) "my polynomial-lattice argument"	200,000; 11,000	C	
Kimura [10] Prop. 9	Rudolph 698480	$1.89 \cdot 10^8$	C	
Kimura chain thm 59 (Rudolph's input)	(ours)	—	R	$d < 1.23 \cdot 10^8$ by Proposition 3.2
Kimura chain thm 58	(ours)	—	R	$d < 6.47 \cdot 10^7$
Kimura Thm 1, parts 1/2	(ours)	—	R	$5.5 \cdot 10^9$; $1.61 \cdot 10^{15}$
Determinant, light	twinheterodox v1–v6.1 [22]	$\ldots, 2000, 570$	R	570 reproduced per row (Section 5)
Determinant, elementary	twinheterodox 698521	2^{15}	R*	kernel-checked [13] (comment 698632); external replay by Olivier (698668)
van Dijk [23]	van Dijk 698529	$8 \cdot 10^9$	C	
Green/van Dijk determinant via Kimi K3	Pratt 698526	Kimi 54,000; a ChatGPT audit "suggests" a repair to 160,000	C	
van Dijk via Kimi K3	Pratt 698682	$< 10^6$, "basically fine"	C	
Green, Thm 1.1 as stated [6]	—	—	—	not yet extracted
OpenAI [16]	—	—	—	Lean theorem is existential
Goldfeld–Gross–Zagier–Oesterlé	Tao's post	e^{7000}; e^{55}	C	classical
(unstated)	Anonymous 698759	187	NR	derivation not posted

Table 1: Extractions of D_0. Values are quoted at source. "R" rows were recomputed by us in ball arithmetic and by a second, separately written implementation; R*: a kernel-checked formalisation, not a recomputation; "C" rows are cited as posted.

Method	D_0	σ_{eq}
GRH	199.12	0.5
determinant, light (570)	570	0.8610
determinant, this paper (RS 1962)	565	0.8600
determinant, this paper (modern ϑ bounds)	548	0.8566
7/8 half-plane	653.99	0.8750
11/12 half-plane	1151.66	0.9167
determinant, elementary	32,768	0.9884
Kimura chain 58	$6.47 \cdot 10^7$	$1 - 2.4 \cdot 10^{-4}$
Banks	10^{10}	$1 - 1.9 \cdot 10^{-5}$

Table 2: The equivalent zero-free abscissa. For large D_0, $1 - \sigma_{\mathrm{eq}} \approx (6/\pi)/\sqrt{D_0}$.

Band	rows	kernel	gap at Q_0	U' on the band
modern explicit bounds (Section 7)				
$[548, 550]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.001328$	≤ -2.275
$[550, 553]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.008996$	≤ -2.31
$[553, 556]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.020566$	≤ -2.33
$[556, 560]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.031867$	≤ -2.34
$[560, 565]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.046950$	≤ -2.35
Rosser–Schoenfeld 1962				
$[565, 570]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.002275$	≤ -2.255
$[570, 580]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.020089$	≤ -2.34
$[580, 600]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.055810$	≤ -2.38
$[600, 652]$	35	deg. 20, $d = \frac{1}{4}$, $e = \frac{1}{25}$	$+0.124424$	≤ -2.42

Table 3: The bands (our own band data; $\gamma = 2341/10000$ in each). Gaps are rounded to the digits shown; the certified enclosures are in the repository. The column U' gives an upper bound over eight Taylor cells covering the band. Each band passes: a positive gap at Q_0; $U' < -\gamma$ on the band; and monotonicity of each Laplace component, so that the error bound at Q_0 is uniform.