

# Artificial General Intelligence for Radiation Oncology

Chenbin Liu, PhD<sup>1</sup>, Zhengliang Liu, MS<sup>2</sup>, Jason Holmes, PhD<sup>3</sup>, Lu Zhang, MS<sup>4</sup>, Lian Zhang, PhD<sup>3</sup>, Yuzhen Ding, PhD<sup>3</sup>, Peng Shu, MS<sup>2</sup>, Zihao Wu, MS<sup>2</sup>, Haixing Dai, PhD<sup>2</sup>, Yiwei Li, MS<sup>2</sup>, Dinggang Shen, PhD<sup>5,6,7</sup>, Ninghao Liu, PhD<sup>2</sup>, Quanzheng Li, PhD<sup>8</sup>, Xiang Li, PhD<sup>8</sup>, Dajiang Zhu, PhD<sup>4</sup>, Tianming Liu, PhD<sup>2</sup>, and Wei Liu, PhD<sup>3</sup>

<sup>1</sup>Department of Radiation Oncology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital & Shenzhen Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Shenzhen, Guangdong, China.

<sup>2</sup>School of Computing, University of Georgia

<sup>3</sup>Department of Radiation Oncology, Mayo Clinic

<sup>4</sup>Department of Computer Science and Engineering, The University of Texas at Arlington

<sup>5</sup>School of Biomedical Engineering, ShanghaiTech University

<sup>6</sup>Shanghai United Imaging Intelligence Co., Ltd.

<sup>7</sup>Shanghai Clinical Research and Trial Center

<sup>8</sup>Department of Radiology, Massachusetts General Hospital and Harvard Medical School

## Abstract

The emergence of artificial general intelligence (AGI) is transforming radiation oncology. As prominent vanguards of AGI, large language models (LLMs) such as GPT-4 and PaLM 2 can process extensive texts and large vision models (LVMs) such as the Segment Anything Model (SAM) can process extensive imaging data to enhance the efficiency and precision of radiation therapy. This paper explores full-spectrum applications of AGI across radiation oncology including initial consultation, simulation, treatment planning, treatment delivery, treatment verification, and patient follow-up. The fusion of vision data with LLMs also creates powerful multimodal models that elucidate nuanced clinical patterns. Together, AGI promises to catalyze a shift towards data-driven, personalized radiation therapy. However, these models should complement human expertise and care. This paper provides an overview of how AGI can transform radiation oncology to elevate the standard of patient care in radiation oncology, with the key insight being AGI's ability to exploit multimodal clinical data at scale.

## 1 Introduction

An estimated 600,000 people in the United States die from cancer every year. Beyond surgery and chemotherapy (now augmented by immunotherapy), radiotherapy has proven as a standard and effective treatment option for nearly 50-70 percent of cancer patients. There exist diverse modalities for the delivery in radiotherapy: (1) brachytherapy involves the surgical implantation of radioactive sources into the patient to kill tumors[1]; (2) early external beam radiotherapy utilizes strong radioactive sources like Cobalt60 positioned at a distance from patients with collimators used to shape and direct beams[2]; (3) contemporary external beam radiotherapy utilizes medical linear accelerator to generate high-energy electron beams or photon beams via bremsstrahlung interactions, optimized for tumor targeting through treatment planning systems[3]; (4) particle therapy with protons or heavier ions produces conformal dose distributions with reduced exit doses, demonstrating

---

efficacy in select cases [4, 5, 6, 7].

Radiotherapy treatment involves six basic stages: initial consultation, simulation, treatment planning, treatment delivery, treatment verification, and patient follow-up. The initial consultation includes a radiation oncologist reviewing the patient's medical history including demographics, operative notes, pathology reports, radiology reports, lab results, discharge, and consults notes to determine the appropriateness of radiotherapy [8]. Simulation precisely localizes the tumor using CT/MR imaging and customized immobilization devices to ensure reproducibility between fractions. In treatment planning, medical professionals delineate target and organ-at-risk volumes on simulation images[9]. Medical dosimetrists and physicists design individualized treatment plans balancing tumor control and normal tissue toxicity[10, 11, 12]. Treatment verification involves the evaluation of dosimetric and geometric accuracy for radiation safety[13, 14]. Qualified radiation therapists perform image-guided radiation delivery as prescribed. Periodic patient follow-up monitors the progress and addresses any side effects. Longitudinal surveillance further guides the management of potential recurrence or residual disease[15][16]. With the emergence of deep learning algorithms, radiotherapy is undergoing substantial transformation. Artificial intelligence (AI) achieves human-level accuracy in the auto-segmentation of organs-at-risk and tumor volumes from CT/MR images, which saves clinicians' times spent on delineation[17, 18]. Beyond auto-segmentation, AI-based algorithms were also implemented in tumor staging[19], image registration[20, 21], automatic treatment planning[22], quality assurance[23], outcomes prediction[24, 25, 26, 27], as well as other areas.

The introduction of the transformer architecture marked a significant milestone in the development of deep learning models, leading to remarkable advancements in terms of parameter size and model complexity [28]. There has been an exponential growth in the scale of these models which were trained on massive amounts of text data. Notable examples of large language models (LLM) that have emerged include OpenAI's generative pre-trained transformer (GPT), Google's pathways language model (PaLM), etc. Some domain-specific LLMs can serve as virtual assistants, facilitating healthcare advice, medical decision support, and administrative tasks[29] [30]. Building upon the success of pre-trained large models in natural language processing (NLP), researchers have embarked on exploring pre-trained large models in the domain of computer vision. Large visual models (LVMs), such as vision transformers (ViT) [31] and VideoMAE V2 [32], exhibit exceptional accuracy in recognizing objects and scenes within images and videos. Researchers are exploring the potential of combining language and visual models to develop advanced AI systems that possess a more human-like understanding of the world [33]. Since AGI models are trained on large and diverse datasets, AGI can achieve impressive zero-shot/few-shot generalization on unseen/limited datasets and perform various real-world tasks with proper prompts.

In this review, we will focus on the different aspects of radiation oncology and the emerging applications and potentials of AGI. The structure of this paper is as follows. In section 2, we introduced the application of LLMs in radiation oncology, including the automatic selection of radiotherapy modalities, patient follow-ups, knowledge extraction from multi-center data, and standardization of clinical data. Section 3 presents how the LVMs and multimodal models shape the domain of medical imaging, and the potentials in radiation oncology. Section 4 discusses the transformative potential of AGI in radiotherapy dose prediction and automatic treatment planning. Section 5 summarizes the deep learning methods in the generation of synthetic CT and talks about the potential of AGI models. Section 6 focuses on the application of deep neural networks in medical image registration. In Section 7, we explore the future directions of AGI in radiation oncology. The potential developments and bottlenecks in the field are discussed as well.




Figure 1: Large Language Models for Radiation Oncology

## 2 Large Language Models for Radiation Oncology

Since the advent of the transformer architecture, there has been a remarkable burgeoning in the domain of Natural Language Processing (NLP). A plethora of sophisticated models have emerged, ranging from early pre-trained BERT models [34, 35], the Generative Pre-trained Transformer (GPT) series [36, 37, 38, 39], PaLM 2 [40] and recent open-source LLMs such as BLOOMZ [41] and the Llama series [42, 43]. These architectures and domain-specific variants [44, 45, 46, 47, 48] exhibit diverse capabilities in effectively tackling an extensive array of NLP tasks [49, 50, 51, 52, 53] and downstream applications [54, 55, 56, 57, 58].

Radiation oncology is a dynamic field, marked by nuanced clinical decision-making and a relentless drive for efficacy and precision. Within this intricate landscape, the application of LLMs can usher in a new era of enhanced patient care and clinical efficiency. With a proven record of success in diverse science domains [59, 60, 49, 61, 62], these models can sift through vast textual datasets to offer informed recommendations tailored to radiation oncology. In a seminal paper by Holmes et al. [58], researchers explored the capabilities of LLMs in radiation oncology physics. The study benchmarked four LLMs against medical physicists and non-experts using an exam developed at Mayo Clinic. ChatGPT (GPT-4) outperformed other models and even medical physicists in some tests, highlighting the potential of LLMs in specialized fields like radiation oncology physics. The impressive accuracy of GPT-4, and to a lesser extent GPT-3.5, in answering questions on the topic of radiation oncology physics suggests that LLMs may be adequate for a wide range of applications in radiation oncology.

Figure 1 illustrates the central role of LLMs in connecting data to clinical applications. For example, selecting the most appropriate radiation modality for each patient is a complex and time-consuming process in radiation oncology [63]. Presently, oncologists often have to spend considerable time reviewing extensive clinical notes, which include a plethora of text documents like demographics, operative notes, pathology and radiology reports, lab results, and discharge and consults notes. These notes are frequently peppered with subtexts and templated information from the Electronic

---

Medical Record (EMR) systems, making them cumbersome to navigate. The manual nature of this task not only requires significant labor but is also prone to human errors and inconsistencies in judgment. Advanced LLMs, fine-tuned specifically on radiation oncology datasets, can ameliorate this issue. Such models can rapidly parse through intricate clinical narratives to recommend the most suitable radiation modality, thereby enhancing both efficiency and the likelihood of optimal patient outcomes.

LLMs also can be pivotal in post-treatment scenarios. By analyzing patient feedback in clinical notes or electronic communications, they can identify patterns suggestive of complications or side-effects. Furthermore, they can generate detailed, patient-specific educational materials, offering insights into the radiation procedures they underwent, potential side effects, and the logic behind the selected treatment modalities.

In addition, radiation oncology practices often differ across institutions. LLMs can process diverse clinical notes, research findings, and treatment narratives from multiple institutions [56, 49]. This vast reservoir of knowledge can then be used to inform best practices or highlight innovative treatment approaches that have found success in particular settings.

Another long-standing problem in radiation oncology is poorly labeled structure names [64, 65, 66, 67, 68, 69]. When analyzing data for a set of patients where the structure contours are required, we want to identify a particular set of structure contours. To do this, the structure name is usually used, however there is typically a lot of variation in the naming of a structure across patients and institutions, which makes the subsequent data analysis very cumbersome. For example, for one patient, prostate might be "prostate" and for another it may be "pstate". For the femur head, the label may be "Femur\_head\_left" or "fem\_head\_l". It is very time-consuming and tedious to standardize the structure names and it is also prone to human errors.

This problem led to the development of a standard for labeling structure names. In 2018, the American Association of Physicists in Medicine (AAPM) created a task group (TG) report known as TG-263[70], which defined a standard for naming structures. However, even with a standard defined, clinics still often are slow to adopt the standard or choose not to. For clinics that choose to adopt the TG-263 standard, data prior to 2018 remains poorly labeled. Tons of these historical patient data in radiation oncology still need to be standardized if we want to take advantage of them for data-mining. For this reason, researchers have investigated methods for re-labeling structure names.

Prospective methods typically employ search algorithms and look-up tables [65] or machine learning [64, 66, 67, 68, 69]. The inputs for machine learning approaches always include the structure names and structures converted to binary masks. Additionally, inputs may include CT information, dose information, or reduced information such as the structure volume. Typically, the inclusion of additional information leads to better results, however studies consistently show that the most significant factor in the accuracy of these models to correctly re-label structure names is the structure name itself.

We have tested GPT-4's ability to re-label structure names on a per-patient basis for 50 prostate patients (never previously seen by GPT-4). The prompt includes the structure names, the TG-263 guidelines, and a list of standard structure names (no patient information). Importantly, the prompt also includes institution-specific instructions or guidelines. Figure 2 shows the results. These results are comparable to the prior reported studies, however there are important distinctions. In the prior reported studies, structure names are always treated as stand-alone, not as part of a set of structure names for the patient. In doing this, they lose important contextual information. Additionally, since GPT-4 does not need to be trained, we may include rare structure names where very little




Figure 2: The accuracy of GPT-4 in re-labeling structure names according to the TG-263 report.

data exists. All of the prior reported studies only test the performance for a small set of structure names. The generality of GPT-4 is far beyond existing AI-based methods and may allow for easy implementation across institutions.

In the evolving landscape of radiation oncology, LLMs serve as valuable tools, amplifying the depth and breadth of human expertise. They promise a synthesis of vast knowledge, aiding clinicians in their quest to offer unparalleled patient care. However, it remains imperative that these models work in tandem with human experts, ensuring a blend of computational efficiency with compassionate care.

### 3 Large Multimodal Models for Radiation Oncology

The success [35, 71, 72, 73, 74, 75, 76, 77] of language models in the field of NLP also provides a revolutionizing paradigm for the advancement of the visual domain. The transformer architecture, in particular, has become building blocks for constructing LVMs [78, 79, 80]. Notable instances include the Vision Transformer (ViT) [81], Swin Transformer [82], VideoMAE V2 [32], and others [83, 84, 85, 86]. These LVMs undergo pre-training using extensive image datasets, equipping them with the capability to capture the complexities of image content and extract intricate semantic information. This empowers them to be highly effective across a wide range of downstream applications [87, 18, 49, 88, 89, 90, 91, 21, 21, 92, 93, 94].

Recently, the emergence of the Segment Anything Model (SAM) [78] has introduced a novel approach to tackling downstream tasks. Unlike traditional pre-trained visual models, SAM operates as a promptable image segmentation model[18]. It utilizes prompts provided by users to guide the model




Figure 3: Large Multimodal Foundation Models for Radiation Oncology

in accurately segmenting the desired areas. By skillfully engineering prompts, SAM is capable of addressing a broad range of downstream tasks effectively. Its remarkable zero-shot generalization capability underscores the pivotal role of prompt engineering in enhancing the efficacy of downstream tasks.

The progress made in the field of LLMs and LVMs has provided a robust groundwork for the evolution of large multimodal foundation models [95], marking a significant milestone in the integration of language and visual modalities. This harmonious combination of various modalities holds the promise of unraveling complex real-world situations and enhancing modeling capabilities. Capitalizing on this complementary synergy, the convergence of unimodal large language models and vision models has gained significant attention, ushering in the era of large multimodal models with boundless potential.

Transitioning from an unimodal framework to a multimodal one requires adjustments in both data and model aspects. In the realm of data, researchers commonly pivot towards creating multimodal-instruction text datasets. This can be achieved by either reconfiguring existing benchmark datasets [96, 97, 98, 99] or employing innovative self-instruction methodologies [100, 96, 101]. On the model horizon, a prevailing approach entails the fusion of information from disparate modalities into LLMs, endowing them with robust reasoning capabilities. Existing studies have taken one of two paths: the direct alignment of multimodal embeddings with LLMs [96, 96, 97, 98] or the utilization of expert models to translate data from various modalities into a form assimilable by LLMs [102, 103]. Through these methodologies, these works reshape LLMs into multimodal conversational agents [100, 96, 102] and versatile task solvers [96, 97, 99] by fine-tuning their performance based on multimodal instructions.

Across various domains, large multimodal foundation models hold profound promise and unparalleled

---

potential. Radiation Oncology [58] is such a quintessential example of a multimodal domain, where multimodality takes on a significant role. Multimodality in radiation oncology refers to the convergence and effective utilization of diverse medical imaging data and complementary information sources. This integration is aimed at elevating the precision and efficacy of radiation therapy. The significance of large multimodal foundation models is notably exemplified in this domain. These models possess the capacity to illuminate intricate patterns, forecast treatment outcomes, and facilitate tailored and precise therapeutic strategies. Their potential is not confined to augmenting existing processes; instead, they hold the power to catalyze a transformative shift within the field. By charting a course towards data-driven insights, these models lay the foundation for a new era of data-driven radiation oncology. Figure 3 highlights some applications of multimodal foundation models in radiation oncology.

One pivotal part of radiation therapy workflow is the accurate prediction of patient outcomes, which include metrics like tumor control, toxicity levels, and overall survival rates [104]. Patient outcome modeling plays a vital role in personalized cancer management across the spectrum of available treatment modalities and has specific applications in areas such as adaptive radiotherapy [105][106]. Conventionally, this process demands the integration of diverse data types, including clinical, imaging, treatment, dosimetric, and biological data. The unstructured nature of clinical notes adds to the complexity, as it obliges clinicians to dedicate extensive hours to data extraction and analysis. Domain-adapted AGI models can dramatically transform this workflow by swiftly navigating through the clinical notes to extract relevant information. In doing so, these models contribute to the development of more robust predictive algorithms. Automation in this domain not only streamlines operations but also holds the potential to increase the accuracy of outcome predictions, thus informing better treatment planning and elevating the standard of patient care.

In recent years, AI-based outcome models have made significant progress, such as acute skin toxicity for breast cancer[107], radiation pneumonitis for lung cancer[108], and overall survival for liver cancer,[109] etc. However, the successful integration of outcome models into clinical practice is not solely contingent upon the accuracy of the models themselves. It also hinges upon the ability of clinicians to interpret and comprehend the specific decisions made by the models. Consequently, the interpretability and explainability of the AI or AGI models hold equal importance to their accuracy when considering their clinical implementation.

## 4 AGI for Treatment Planning

Radiation therapy serves as a key treatment strategy in clinical practice. Over recent years, its effectiveness has been remarkably enhanced, largely attributed to state-of-the-art modalities like Intensity-Modulated Radiation Therapy (IMRT) and Intensity-Modulated Proton Therapy (IMPT) [110, 111, 112, 113, 114, 115]. This advancement has dual implications: It has improved treatment plan quality but has concurrently augmented the intricacy of these plans with longer planning duration and potential challenges in maintaining treatment accuracy. Clinicians often have to balance lots of planning-related parameters which necessitates intricate communications between dosimetrists and physicians to fine-tune the planning parameters. As a remedy to this, the research landscape is increasingly focusing on optimizing dose distribution and setting reasonable constraints[22, 116, 117, 118]. The overarching ambition is to efficiently and accurately generate an optimal dose distribution comparable to a manual plan based on previous treatment planning knowledge and ensure both time efficiency and uniform excellence, irrespective of the planner's level of expertise [119, 120].




Figure 4: AI based treatment planning workflow. a) KBP method b) Dose Prediction method. The dotted box means an improved enhancement for the workflow.

In clinical practice, one common strategy for enhancing the efficiency and quality of manual treatment planning involves reviewing prior cases deemed to be exemplary. In particular, parameters from these previous cases—such as beam configurations and dose-volume histogram (DVH) objectives used in inverse planning—can be directly integrated into the current planning process or serve as benchmarks for decision-making [121, 122]. Building on this concept, researchers have employed statistical models to distill specific attributes from these superior cases, utilizing the best of clinical acumen and knowledge. The initial class of methodologies focuses on predicting viable plan parameters, such as DVH objectives to guide the optimization process. Approaches informed by artificial intelligence (AI) display remarkable advantages over conventional rule-based algorithms. Commonly known as Knowledge-Based Planning (KBP), these techniques anticipate possible dose distribution patterns for a new plan by drawing insights from historical, high-quality plans. While atlas-based KBP methods identify the most analogous patients in the plan database to ascertain the optimal starting parameters for inverse planning, model-based KBP methods deploy various computational models trained on previous plans to predict ideal parameters for the new case [123, 124, 125]. The parameters generated by KBP can streamline the optimization process, cutting down on the number of trial-and-error adjustments, as they are inherently more aligned with ideal outcomes than those derived solely from planners’ experience. When it comes to describing the geometric relationship between organs-at-risk (OAR) and the planning target volumes (PTV), various metrics such as the overlap volume histogram (OVH), distance-to-target histogram (DTH), and out-of-field volume are commonly employed. These metrics can inform DVH predictions through machine learning techniques, including support vector regression (SVR) [126, 127, 128]. One commercial embodiment of this DVH-KBP model is Varian’s RapidPlan [129]. Contrary to DVH prediction alone, 3D-dose prediction preserves spatial specificity. Given that the DVH serves as a one-dimensional representation of the three-dimensional dose distribution, it inherently loses some spatial information during the translation. Consequently, identical DVH curves could originate from divergent dose distributions, potentially resulting in clinically meaningful disparities. Physicians typically utilize both DVH and raw dose data to assess the quality of treatment plans. Therefore, dose prediction could offer certain advantages that DVH prediction alone may not capture. Currently, deep learning algorithms have been proven to achieve expert-level 3D-dose prediction accuracy but with enhanced efficiency, even in advanced therapeutic modalities like helical tomotherapy and proton radiotherapy [130, 131, 132, 133, 89]. Figure 4 shows the workflow of deep learning-based dose prediction and clinical deployment for pencil beam scanning proton therapy [89]. Moreover, predicted spatial dose distribution can be operationalized to generate clinically executable plans via dose-mimicking algorithms [134, 135].

---

In recent years, large foundational models such as ChatGPT and SAM have garnered substantial attention. As the burgeoning frontrunners in the AGI arena, these models offer tremendous promise for revolutionizing dose prediction in radiation therapy [38, 78]. Current dose prediction paradigms, whether employing convolutional neural networks (CNN) or traditional machine-learning algorithms, often specialize in either particular types of clinical cases or specific anatomical locations [136]. This specialization complicates their deployment and broad applicability in clinical environments. In contrast, AGI frameworks possess an innate versatility, enabling a singular model to tackle a multitude of tasks [137]. This capability could facilitate the development of a unified AGI-based model for comprehensive dose prediction, substantially reducing the complexity of its clinical integration. Moreover, AGI exhibits robust cross-modality learning potential; for example, SAM can transfer the segmentation knowledge acquired from natural images to medical image analyses. A recent study on SAM in clinical radiation therapy shows that the natural images trained SAM can achieve clinically acceptable Dice score  $>0.7$  for most OARs segmentation across four disease sites, demonstrating super generalization capabilities across different disease sites and different modalities that make it feasible to develop a generic auto-segmentation model in radiotherapy with SAM [18]. Given the often limited availability of training data for specific clinical scenarios or innovative techniques in clinical practice, AGI offers a robust solution for learning from a broad variety of knowledge. It permits fine-tuning based on a minimal dataset, generating reliable dose predictions. This is especially beneficial in cases of rare conditions or specialized techniques, such as MRI-based dose prediction, underscoring AGI's transformative potential in radiation therapy treatment planning [138].

## 5 Synthetic CT Generation Using Deep Learning

Computed tomography (CT) is the primary imaging modality in the current practice of radiation therapy. It provides three-dimensional structural information of the patient for treatment planning, enabling electron density calibration required for dose calculation [139]. Synthetic CT offers the possibility of reducing the additional dose of CT scan and expanding the usage of other imaging modalities in adaptive radiotherapy, such as MR, CBCT, and MVCT [140][141][142][105, 143]. Magnetic resonance imaging (MRI) simulation has superior soft-tissue contrast compared with CT and delivers no ionizing radiation, which plays a vital role in target and organs-at-risk delineation. To create a treatment plan, MRI images have to be registered onto CT scans for dose calculation. Traditional synthetic CT generation methods can be grouped into three categories: bulk density override, atlas-based, and voxel-based methods [144]. In recent years, many deep learning approaches have been proposed to predict the synthetic CT for MR-only radiotherapy [145][146][147][148][149][150]. There are mainly two categories, generator-only model, and generator and discriminator model. Han first proposed a deep convolutional neural network (DCNN) to model the MR-to-CT mapping using eighteen brain tumor patients [151]. Other generator-only architectures, such as U-net [152][153], Res-Net[154], multiple deep CNNs [155], deep embedding CNN [156], patch-based CNN [157], also have been used to estimate the electron density maps. The application of these methods extended from brain cancer [151] to head and neck cancer [157], and prostate cancer [156].

Compared with the generator-only model, generative adversarial network (GAN) architecture involves two sub-models: a generator model to generate plausible data, and a discriminator model to determine fake from real data. Most researchers used the co-registered/paired MR and CT images as the training data of GAN and its variant models, such as GAN [158], conditional GAN [150], residual transformer conditional GAN [159], compensation cycle GAN [160]. To overcome the limitation of precisely aligned MR and CT images, some GAN models were developed to synthesize images from




Figure 5: Synthetic CT generation using deep learning

weakly paired, or unpaired data [161][162][163][164][165]. Cycle-consistent GAN model was used in the training of weakly paired CT and MR images [161]. Zeng explored a hybrid GAN consisting of a 3D generator network and a 2D discriminator network, in which 3D generator was believed to model the 3D spatial information across slices [163]. Augmented cycle GAN was proposed to generate synthetic CT using unpaired data from multiple scanners in different centers [165]. The model provided improved generalization performance and produced clinically acceptable synthetic CTs [165]. Although the robustness and generalizability of the generation models were considered in recent studies [166][167], the question of these techniques in clinical application is how to evaluate the quality of synthetic CT without acquiring the ground truth CT [168]. As far as we know, there are no studies using AGI to generate synthetic CT. The large vision models may be trained using diverse datasets from multiple centers, disease sites, and patient populations, which may ensure the model's robustness and generalizability.

## 6 AGI for Image Registration

Image registration, which aims to find the spatial relationship between two or multiple sets of images, is usually formalized as the optimization of a function balancing the similarity between images in terms of intensity, topology, or both [169]. Compared to rigid image registration (RIR), the deformable image registration (DIR), which attempts to find the voxel-specific spatial relationship between two or multiple sets of images, has far more flexibility than RIR, thus, it can be used in more complicated clinical scenarios such as images with large anatomical structure changes. DIR has been extensively used in radiation therapy such as automatic segmentation [170, 171], mathematical modeling [172, 173, 174, 175], functional imaging [176, 177, 178], and dose deformation [179, 180, 181, 182, 183, 184, 185]. Over the years, many conventional DIR approaches have been developed and adopted clinically. The conventional DIR approaches can be broadly categorized into two categories: parametric [174, 175, 186] and non-parametric [187, 188, 189, 190] models. The parametric model generates deformable vector fields (DVF) as a linear combination of its basic functions. The B-spline model is an example of such parametric models and it can handle the local change of a voxel by linear regression from nearby voxels within a certain distance. This property significantly reduces the computation time and memory required. Yet, the results can only be used for CTs with some strict conditions, such as breath-holding or respiratory gating, which limits its wide applications in clinics. In contrast, non-parametric models such as demons-based methods calculate transformation vectors of all voxels, thus achieving more accurate DVF, but requiring

---

more computation time and memory than the parametric models.

Recently, several deep learning-based methods have been developed to speed up DIR in medical image analysis. According to the supervision used in model training, it can be broadly categorized as, supervised and unsupervised learning-based DIR approaches. For the supervised learning-based DIR methods, the ground-truth DVF is needed as the supervision. Yang et al. [191] proposed a two-steps deep learning framework for predicting the momentum parameterization for the large deformation diffeomorphic metric mapping (LDDMM) model. The proposed deep learning framework consists of two auto-encoder networks with the same architecture, in which the first auto-encoder is used to estimate the initial patch-wise momentum and the second one further tunes the initial patch-wise momentum. Although the proposed method is much faster comparing to the conventional DIR approaches, the computational complexity is higher than a typical single-step deep learning network. Besides, since it has two cascade networks, the symmetrical error may accumulate as the layers go deeper. The supervised learning-based DIR for other disease sites have also been studied [192, 193, 194]. Nevertheless, the generation of the ground-truth DVFs that are used for model training can be time-consuming as well, besides, the computed ground-truth DVFs may also be different from the real DVFs, thus introducing unexpected errors. Therefore, the unsupervised learning-based DIR approach which learns the similarity between the ground-truth image and wrapped image is more practical and favorable in clinical applications. Balakrishnan et al. [195] proposed a UNet-like model termed as VoxelMorph to learn the DVFs from pairs of magnetic resonance images (MRIs) (i.e., moving images and fixed images), then the generated DVFs and moving images go through a non-learnable spatial transformation to form the final generated warped images that resemble the fixed images. The VoxelMorph can achieve comparable performance as the state-of-the-art conventional DIR methods, whereas it is orders of magnitude faster. Thus, it has been widely used in medical image analysis. Most of these methods have been proposed for MRIs, which typically have high-resolution and rich anatomical information, whereas in radiation therapy the commonly used image modality is CT with a relatively low resolution. Recently, [21] have extended the unsupervised learning-based DIR to CT modality and yielded fast and accurate results.

With the advent of large multimodal foundation models, there has been a surge in the development of AGI-based image processing models, most notably represented by SAM, which has consequently accelerated research into the capabilities of AGI for image registration tasks. The general-purpose nature of AGI models makes them particularly well-suited for complex tasks like multi-modal image registration. Preliminary research reveals that SAM-enabled systems can perform real-time, accurate tracking and mapping of reference points in deformable images. For example, SAM has been shown to be effective in tracking respiratory motion within lung images. This capacity for continuous, point-to-point tracking has promising implications for its broader adoption in medical image registration scenarios [196].

## 7 Discussion and Conclusion

In this review, we have outlined how AGI can revolutionize radiation oncology, enhancing healthcare standards. The key insight lies in AGI's capacity to leverage large-scale multimodal clinical data. Among the six stages of the radiotherapy workflow, initial consultation, simulation, treatment planning, treatment delivery, treatment verification, and patient follow-up all involve visual and linguistic information. The processing and comprehension abilities of AGI in relation to visual and linguistic information have the potential to provide support throughout every stage of radiotherapy.




Figure 6: AGI for Image registration

This could result in improved radiotherapy safety and precision, enhanced efficiency, and favorable patient outcomes. However, the realization of AGI’s potential in radiotherapy necessitates seamless integration with existing medical systems, and the performance of AGI is limited by its dependence on domain-specific knowledge. To surpass this limitation, future efforts should prioritize broadening the scope of knowledge by integrating diverse and comprehensive clinical datasets, utilizing interdisciplinary approaches, and encouraging interdisciplinary collaborations with clinical experts in radiation oncology.

As far as we know, LLMs, like GPT-3, are trained on text from multiple sources which include web pages, internet-based books, and Wikipedia. The LVMs are trained on the natural image datasets. Vision-language models are typically trained on multi-modal datasets harvested from the web in the form of matching image/video and text pairs. Although AGI generalized human cognitive abilities faced with unfamiliar tasks, most of the AGI models were not designed to provide high-quality clinical applications. To fine-tune these LLMs, LVMs, or vision-language models with high-quality medical data holds promising potential. By incorporating such data, AGI models can benefit from a broader range of clinical scenarios, leading to improved accuracy and performance in the applications of radiation oncology.

One of the challenges is the data standardization. The inconsistency of structure names in radiotherapy poses a significant challenge when employing automated methods. As AAPM task group 263 reported, it is important to follow a standardized target and organs-at-risk naming rules for AGI training, data sharing between multi-centers, and quality assurance [197]. Data sharing presents an additional challenge in building a large, high-quality medical training dataset due to factors like privacy concerns, legal and ethical restrictions, and standardization obstacles to disseminating individual patient records. Data sharing has become increasingly prevalent in some fields of medical research, especially among genomics researchers and groups conducting systematic reviews and meta-analyses. Investigators still have concerns about sharing individual patient data from clinical trials [198]. Data interpretation in radiotherapy poses a notable challenge that warrants careful consideration to ensure meaningful analysis. This challenge arises from the inherent complexities of the clinical data, treatment uncertainties, and errors/variations in data recording. Integration of heterogeneous data in radiotherapy, including patient consultation, clinical examinations, medical imaging, treatment plans, verification records, machine logs, and patient outcomes, presents difficulties due to variations in formats, resolutions, etc.

Interdisciplinary collaboration teams play a vital role in ensuring the clinical applications of AGI models. These teams offer diverse perspectives and expertise, enabling comprehensive feedback

---

throughout the development and validation stages. Involving end-users, such as radiation oncologists, and medical physicists, in the development process through user-centered design methodologies[49]. It ensures that the AGI models meet the domain-specific needs and preferences, enhancing the usability and acceptance of the models. The involvement of professionals from diverse disciplines ensures a comprehensive approach to address precision, ethical considerations, and regulatory compliance. In addition, developing AGI models that collaborate with medical professionals, rather than replacing them, offers several advantages. This approach allows for the maximization of AGI benefits while avoiding the resistance of medical professionals and optimizing their workload. With the development of AI, radiotherapy staffs prioritize continued education using AI to preserve their skills, such as manual segmentation ability[199]. The introduction of AGI models will shape the evolving role of medical professionals in the radiotherapy department. With AGI handling micro/macro processes, the focus of radiotherapy staffs' work will shift towards verifying the model performance quality. As a result, the continued education of radiotherapy staffs will need to adapt accordingly.

AGI holds great promise for radiotherapy, offering advancements in patient consultation, image registration, structure segmentation, radiation dose prediction, auto-treatment planning, patient outcomes prediction, etc. However, challenges remain regarding clinical datasets, regulation, and interdisciplinary collaboration considerations. Despite these challenges, the opportunities presented by AGI in enhancing radiation oncology and improving clinical automation are significant. This review aims to serve as a reference and catalyst for further exploration in this rapidly evolving field, stimulating advancements and discussions for the benefit of healthcare.

## References

- [1] Subir Nag, David Beyer, Jay Friedland, Peter Grimm, and Ravinder Nath. American brachytherapy society (abs) recommendations for transperineal permanent brachytherapy of prostate cancer. *International Journal of Radiation Oncology\* Biology\* Physics*, 44(4):789–799, 1999.
- [2] JS Lott and Ivan H Smith. Cobalt-60 beam therapy in carcinoma of the esophagus. *Radiology*, 71(3):321–326, 1958.
- [3] JV Siebers, JK Gardner, JJ Gordon, S Wang, and JD Ververs. Quantification of exit fluence variations and implications for exit fluence-based dose reconstruction based. *International Journal of Radiation Oncology, Biology, Physics*, 72(1):S552, 2008.
- [4] Radhe Mohan and David Grosshans. Proton therapy—present and future. *Advanced drug delivery reviews*, 109:26–44, 2017.
- [5] Steven E Schild, William G Rule, Jonathan B Ashman, Sujay A Vora, Sameer Keole, Aman Anand, Wei Liu, and Martin Bues. Proton beam therapy for locally advanced lung cancer: A review. *World journal of clinical oncology*, 5(4):568, 2014.
- [6] Wei Deng, Yunze Yang, Chenbin Liu, Martin Bues, Radhe Mohan, William W Wong, Robert H Foote, Samir H Patel, and Wei Liu. A critical review of let-based intensity-modulated proton therapy plan evaluation and optimization for head and neck cancer management. *International Journal of Particle Therapy*, 8(1):36–49, 2021.
- [7] Wei Liu. System and method for robust intensity-modulated proton therapy planning, August 2019. US Patent 10,369,381.

- 
- [8] Jason W Chan, Nicole Hohenstein, Colin Carpenter, Adam J Pattison, Olivier Morin, Gilmer Valdes, Cristina Tolentino Sanchez, Jennifer Perkins, Timothy D Solberg, and Sue S Yom. Artificial intelligence-guided prediction of dental doses before planning of radiation therapy for oropharyngeal cancer: Technical development and initial feasibility of implementation. *Advances in radiation oncology*, 7(2):100886, 2022.
  - [9] Wei Liu, Zhongxing Liao, Steven E Schild, Zhong Liu, Heng Li, Yupeng Li, Peter C Park, Xiaoqiang Li, Joshua Stoker, Jiajian Shen, et al. Impact of respiratory motion on worst-case scenario optimized intensity modulated proton therapy for lung cancers. *Practical radiation oncology*, 5(2):e77–e86, 2015.
  - [10] Steve Webb and Alan Effraim Nahum. A model for calculating tumour control probability in radiotherapy including the effects of inhomogeneous distributions of dose and clonogenic cell density. *Physics in medicine & biology*, 38(6):653, 1993.
  - [11] Maryam Zaghian, Gino Lim, Wei Liu, and Radhe Mohan. An automatic approach for satisfying dose-volume constraints in linear fluence map optimization for impt. *Journal of cancer therapy*, 5(2):198, 2014.
  - [12] Maryam Zaghian, Wenhua Cao, Wei Liu, Laleh Kardar, Sharmalee Randeniya, Radhe Mohan, and Gino Lim. Comparison of linear and nonlinear programming approaches for “worst case dose” and “minmax” robust optimization of intensity-modulated proton therapy dose distributions. *Journal of applied clinical medical physics*, 18(2):15–25, 2017.
  - [13] James E Younkin, Martin Bues, Terence T Sio, Wei Liu, Xiaoning Ding, Sameer R Keole, Joshua B Stoker, and Jiajian Shen. Multiple energy extraction reduces beam delivery time for a synchrotron-based proton spot-scanning system. *Advances in Radiation Oncology*, 3(3):412–420, 2018.
  - [14] James E Younkin, Jiajian Shen, Martin Bues, Daniel G Robertson, Daniel W Mundy, Edward Clouser, Wei Liu, Xiaoning Ding, and Joshua B Stoker. an efficient daily qa procedure for proton pencil beam scanning. *Medical physics*, 45(3):1040–1049, 2018.
  - [15] Y Yu Nathan, Todd A DeWees, Chenbin Liu, Thomas B Daniels, Jonathan B Ashman, Staci E Beamer, Dawn E Jaroszewski, Helen J Ross, Harshita R Paripati, Jean-Claude M Rwigema, et al. Early outcomes of patients with locally advanced non-small cell lung cancer treated with intensity-modulated proton therapy versus intensity-modulated radiation therapy: the mayo clinic experience. *Advances in Radiation Oncology*, 5(3):450–458, 2020.
  - [16] Nathan Y Yu, Todd A DeWees, Molly M Voss, William G Breen, Jennifer S Chiang, Julia X Ding, Thomas B Daniels, Dawn Owen, Kenneth R Olivier, Yolanda I Garces, et al. Cardiopulmonary toxicity following intensity-modulated proton therapy (impt) versus intensity-modulated radiation therapy (imrt) for stage iii non-small cell lung cancer. *Clinical Lung Cancer*, pages S1525–7304, 2022.
  - [17] Anjali Balagopal, Howard Morgan, Michael Dohopolski, Ramsey Timmerman, Jie Shan, Daniel F Heitjan, Wei Liu, Dan Nguyen, Raquibul Hannan, Aurelie Garant, et al. Psa-net: Deep learning-based physician style-aware segmentation network for postoperative prostate cancer clinical target volumes. *Artificial Intelligence in Medicine*, 121:102195, 2021.
  - [18] Lian Zhang, Zhengliang Liu, Lu Zhang, Zihao Wu, Xiaowei Yu, Jason Holmes, Hongying Feng, Haixing Dai, Xiang Li, Quanzheng Li, et al. Segment anything model (sam) for radiation oncology. *arXiv preprint arXiv:2306.11730*, 2023.
-

- 
- [19] Shujun Liang, Xiuyu Dong, Kaifan Yang, Zhiqin Chu, Fan Tang, Feng Ye, Bei Chen, Jian Guan, and Yu Zhang. A multi-perspective information aggregation network for automated t-staging detection of nasopharyngeal carcinoma. *Physics in Medicine & Biology*, 67(24):245007, 2022.
  - [20] Xiaohuan Cao, Jianhua Yang, Jun Zhang, Qian Wang, Pew-Thian Yap, and Dinggang Shen. Deformable image registration using a cue-aware deep regression network. *IEEE Transactions on Biomedical Engineering*, 65(9):1900–1911, 2018.
  - [21] Yuzhen Ding, Hongying Feng, Yunze Yang, Jason Holmes, Zhengliang Liu, David Liu, William W Wong, Nathan Y Yu, Terence T Sio, Steven E Schild, et al. Deep-learning based fast and accurate 3d ct deformable image registration in lung cancer. *Medical Physics*, 2023.
  - [22] Mingqing Wang, Qilin Zhang, Saikit Lam, Jing Cai, and Ruijie Yang. A review on application of deep learning algorithms in external beam radiotherapy automated treatment planning. *Frontiers in oncology*, 10:580919, 2020.
  - [23] Liesbeth Vandewincke, Michaël Claessens, Anna Dinkla, Charlotte Brouwer, Wouter Crijns, Dirk Verellen, and Wouter van Elmpt. Overview of artificial intelligence-based applications in radiotherapy: Recommendations for implementation and quality assurance. *Radiotherapy and Oncology*, 153:55–66, 2020.
  - [24] Wei Liu, Samir H Patel, Daniel P Harrington, Yanle Hu, Xiaoning Ding, Jiajian Shen, Michele Y Halyard, Steven E Schild, William W Wong, Gary E Ezzell, et al. Exploratory study of the association of volumetric modulated arc therapy (vmat) plan robustness with local failure in head and neck cancer. *Journal of applied clinical medical physics*, 18(4):76–83, 2017.
  - [25] Bin Wang, Huayong Jiang, Tingyang Zhou, Ning Ma, Wei Liu, Yajie Wang, and Li Zuo. Expression of icosl is associated with decreased survival in invasive breast cancer. *PeerJ*, 7:e6903, 2019.
  - [26] Yunze Yang, Samir H Patel, Jidapa Bridhikitti, William W Wong, Michele Y Halyard, Lisa A McGee, Jean-Claude M Rwigema, Steven E Schild, Sujay A Vora, Tianming Liu, et al. Exploratory study of seed spots analysis to characterize dose and linear-energy-transfer effect in adverse event initialization of pencil-beam-scanning proton therapy. *Medical physics*, 49(9):6237–6252, 2022.
  - [27] Yunze Yang, Olivia M Muller, Satomi Shiraishi, Matthew Harper, Adam C Amundson, William W Wong, Lisa A McGee, Jean-Claude M Rwigema, Steven E Schild, Martin Bues, et al. Empirical relative biological effectiveness (rbe) for mandible osteoradionecrosis (orn) in head and neck cancer patients treated with pencil-beam-scanning proton therapy (pbspt): a retrospective, case-matched cohort study. *Frontiers in Oncology*, 12:843175, 2022.
  - [28] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural information processing systems*, 30, 2017.
  - [29] Stephen Gilbert, Hugh Harvey, Tom Melvin, Erik Vollebregt, and Paul Wicks. Large language model ai chatbots require approval as medical devices. *Nature Medicine*, pages 1–3, 2023.
  - [30] Karan Singhal, Shekoofeh Azizi, Tao Tu, S Sara Mahdavi, Jason Wei, Hyung Won Chung, Nathan Scales, Ajay Tanwani, Heather Cole-Lewis, Stephen Pfohl, et al. Large language models encode clinical knowledge. *Nature*, pages 1–9, 2023.
-

- 
- [31] Bichen Wu, Chenfeng Xu, Xiaoliang Dai, Alvin Wan, Peizhao Zhang, Zhicheng Yan, Masayoshi Tomizuka, Joseph Gonzalez, Kurt Keutzer, and Peter Vajda. Visual transformers: Token-based image representation and processing for computer vision, 2020.
- [32] Limin Wang, Bingkun Huang, Zhiyu Zhao, Zhan Tong, Yinan He, Yi Wang, Yali Wang, and Yu Qiao. Videomae v2: Scaling video masked autoencoders with dual masking. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pages 14549–14560, 2023.
- [33] Xuran Pan, Tianzhu Ye, Dongchen Han, Shiji Song, and Gao Huang. Contrastive language-image pre-training with knowledge graphs. *Advances in Neural Information Processing Systems*, 35:22895–22910, 2022.
- [34] Jacob Devlin, Ming-Wei Chang, Kenton Lee, and Kristina Toutanova. Bert: Pre-training of deep bidirectional transformers for language understanding. *arXiv preprint arXiv:1810.04805*, 2018.
- [35] Ce Zhou, Qian Li, Chen Li, Jun Yu, Yixin Liu, Guangjing Wang, Kai Zhang, Cheng Ji, Qiben Yan, Lifang He, et al. A comprehensive survey on pretrained foundation models: A history from bert to chatgpt. *arXiv preprint arXiv:2302.09419*, 2023.
- [36] Alec Radford, Karthik Narasimhan, Tim Salimans, Ilya Sutskever, et al. Improving language understanding by generative pre-training. 2018.
- [37] Alec Radford, Jeffrey Wu, Rewon Child, David Luan, Dario Amodei, Ilya Sutskever, et al. Language models are unsupervised multitask learners. *OpenAI blog*, 1(8):9, 2019.
- [38] Tom Brown, Benjamin Mann, Nick Ryder, Melanie Subbiah, Jared D Kaplan, Prafulla Dhariwal, Arvind Neelakantan, Pranav Shyam, Girish Sastry, Amanda Askell, et al. Language models are few-shot learners. *Advances in neural information processing systems*, 33:1877–1901, 2020.
- [39] Long Ouyang, Jeff Wu, Xu Jiang, Diogo Almeida, Carroll L Wainwright, Pamela Mishkin, Chong Zhang, Sandhini Agarwal, Katarina Slama, Alex Ray, et al. Training language models to follow instructions with human feedback. *arXiv preprint arXiv:2203.02155*, 2022.
- [40] Rohan Anil, Andrew M Dai, Orhan Firat, Melvin Johnson, Dmitry Lepikhin, Alexandre Passos, Siamak Shakeri, Emanuel Taropa, Paige Bailey, Zhifeng Chen, et al. Palm 2 technical report. *arXiv preprint arXiv:2305.10403*, 2023.
- [41] Niklas Muennighoff, Thomas Wang, Lintang Sutawika, Adam Roberts, Stella Biderman, Teven Le Scao, M Saiful Bari, Sheng Shen, Zheng-Xin Yong, Hailey Schoelkopf, et al. Crosslingual generalization through multitask finetuning. *arXiv preprint arXiv:2211.01786*, 2022.
- [42] Hugo Touvron, Thibaut Lavril, Gautier Izacard, Xavier Martinet, Marie-Anne Lachaux, Timothée Lacroix, Baptiste Rozière, Naman Goyal, Eric Hambro, Faisal Azhar, et al. Llama: Open and efficient foundation language models. *arXiv preprint arXiv:2302.13971*, 2023.
- [43] Hugo Touvron, Louis Martin, Kevin Stone, Peter Albert, Amjad Almahairi, Yasmine Babaee, Nikolay Bashlykov, Soumya Batra, Prajjwal Bhargava, Shruti Bhosale, et al. Llama 2: Open foundation and fine-tuned chat models. *arXiv preprint arXiv:2307.09288*, 2023.
- [44] Zhengliang Liu, Aoxiao Zhong, Yiwei Li, Longtao Yang, Chao Ju, Zihao Wu, Chong Ma, Peng Shu, Cheng Chen, Sekeun Kim, et al. Radiology-gpt: A large language model for radiology. *arXiv preprint arXiv:2306.08666*, 2023.

- 
- [45] Kai Zhang, Jun Yu, Zhiling Yan, Yixin Liu, Eashan Adhikarla, Sunyang Fu, Xun Chen, Chen Chen, Yuyin Zhou, Xiang Li, et al. Biomedgpt: A unified and generalist biomedical generative pre-trained transformer for vision, language, and multimodal tasks. *arXiv preprint arXiv:2305.17100*, 2023.
- [46] Zhengliang Liu, Zihao Wu, Mengxuan Hu, Bokai Zhao, Lin Zhao, Tianyi Zhang, Haixing Dai, Xianyan Chen, Ye Shen, Sheng Li, et al. Pharmacygpt: The ai pharmacist. *arXiv preprint arXiv:2307.10432*, 2023.
- [47] Saed Rezayi, Haixing Dai, Zhengliang Liu, Zihao Wu, Akarsh Hebbar, Andrew H Burns, Lin Zhao, Dajiang Zhu, Quanzheng Li, Wei Liu, et al. Clinicalradiobert: Knowledge-infused few shot learning for clinical notes named entity recognition. In *Machine Learning in Medical Imaging: 13th International Workshop, MLMI 2022, Held in Conjunction with MICCAI 2022, Singapore, September 18, 2022, Proceedings*, pages 269–278. Springer, 2022.
- [48] Chong Ma, Zihao Wu, Jiaqi Wang, Shaochen Xu, Yaonai Wei, Zhengliang Liu, Lei Guo, Xiaoyan Cai, Shu Zhang, Tuo Zhang, et al. Impressiongpt: An iterative optimizing framework for radiology report summarization with chatgpt. *arXiv preprint arXiv:2304.08448*, 2023.
- [49] Xiang Li, Lu Zhang, Zihao Wu, Zhengliang Liu, Lin Zhao, Yixuan Yuan, Jun Liu, Gang Li, Dajiang Zhu, Pingkuan Yan, et al. Artificial general intelligence for medical imaging. *arXiv preprint arXiv:2306.05480*, 2023.
- [50] Zihan Guan, Zihao Wu, Zhengliang Liu, Dufan Wu, Hui Ren, Quanzheng Li, Xiang Li, and Ninghao Liu. Cohortgpt: An enhanced gpt for participant recruitment in clinical study. *arXiv preprint arXiv:2307.11346*, 2023.
- [51] Haixing Dai, Zhengliang Liu, Wenxiong Liao, Xiaoke Huang, Zihao Wu, Lin Zhao, Wei Liu, Ninghao Liu, Sheng Li, Dajiang Zhu, et al. Chataug: Leveraging chatgpt for text data augmentation. *arXiv preprint arXiv:2302.13007*, 2023.
- [52] Mengyue Zhou, Xu Liu, David Liu, Zihao Wu, Zhengliang Liu, Lin Zhao, Dajiang Zhu, Lei Guo, Junwei Han, Tianming Liu, et al. Fine-grained artificial neurons in audio-transformers for disentangling neural auditory encoding. In *Findings of the Association for Computational Linguistics: ACL 2023*, pages 7943–7956, 2023.
- [53] Zihao Wu, Lu Zhang, Chao Cao, Xiaowei Yu, Haixing Dai, Chong Ma, Zhengliang Liu, Lin Zhao, Gang Li, Wei Liu, et al. Exploring the trade-offs: Unified large language models vs local fine-tuned models for highly-specific radiology nli task. *arXiv preprint arXiv:2304.09138*, 2023.
- [54] Zhengliang Liu, Xiaowei Yu, Lu Zhang, Zihao Wu, Chao Cao, Haixing Dai, Lin Zhao, Wei Liu, Dinggang Shen, Quanzheng Li, et al. Deid-gpt: Zero-shot medical text de-identification by gpt-4. *arXiv preprint arXiv:2303.11032*, 2023.
- [55] Wenxiong Liao, Zhengliang Liu, Haixing Dai, Shaochen Xu, Zihao Wu, Yiyang Zhang, Xiaoke Huang, Dajiang Zhu, Hongmin Cai, Tianming Liu, et al. Differentiate chatgpt-generated and human-written medical texts. *arXiv preprint arXiv:2304.11567*, 2023.
- [56] Yiheng Liu, Tianle Han, Siyuan Ma, Jiayue Zhang, Yuanyuan Yang, Jiaming Tian, Hao He, Antong Li, Mengshen He, Zhengliang Liu, et al. Summary of chatgpt/gpt-4 research and perspective towards the future of large language models. *arXiv preprint arXiv:2304.01852*, 2023.

- 
- [57] Zhengliang Liu, Tianyang Zhong, Yiwei Li, Yutong Zhang, Yi Pan, Zihao Zhao, Peixin Dong, Chao Cao, Yuxiao Liu, Peng Shu, et al. Evaluating large language models for radiology natural language processing. *arXiv preprint arXiv:2307.13693*, 2023.
- [58] Jason Holmes, Zhengliang Liu, Lian Zhang, Yuzhen Ding, Terence T Sio, Lisa A McGee, Jonathan B Ashman, Xiang Li, Tianming Liu, Jiajian Shen, et al. Evaluating large language models on a highly-specialized topic, radiation oncology physics. *arXiv preprint arXiv:2304.01938*, 2023.
- [59] Guoyu Lu, Sheng Li, Gengchen Mai, Jin Sun, Dajiang Zhu, Lilong Chai, Haijian Sun, Xianqiao Wang, Haixing Dai, Ninghao Liu, et al. Agi for agriculture. *arXiv preprint arXiv:2304.06136*, 2023.
- [60] Gengchen Mai, Weiming Huang, Jin Sun, Suhang Song, Deepak Mishra, Ninghao Liu, Song Gao, Tianming Liu, Gao Cong, Yingjie Hu, et al. On the opportunities and challenges of foundation models for geospatial artificial intelligence. *arXiv preprint arXiv:2304.06798*, 2023.
- [61] Ehsan Latif, Gengchen Mai, Matthew Nyaaba, Xuansheng Wu, Ninghao Liu, Guoyu Lu, Sheng Li, Tianming Liu, and Xiaoming Zhai. Artificial general intelligence (agi) for education. *arXiv preprint arXiv:2304.12479*, 2023.
- [62] Lin Zhao, Lu Zhang, Zihao Wu, Yuzhong Chen, Haixing Dai, Xiaowei Yu, Zhengliang Liu, Tuo Zhang, Xintao Hu, Xi Jiang, et al. When brain-inspired ai meets agi. *arXiv preprint arXiv:2303.15935*, 2023.
- [63] Elizabeth Huynh, Ahmed Hosny, Christian Guthier, Danielle S Bitterman, Steven F Petit, Daphne A Haas-Kogan, Benjamin Kann, Hugo JW Aerts, and Raymond H Mak. Artificial intelligence in radiation oncology. *Nature Reviews Clinical Oncology*, 17(12):771–781, 2020.
- [64] Ali Haidar, Matthew Field, Vikneswary Batumalai, Kirrily Cloak, Daniel Al Mouiee, Phillip Chlap, Xiaoshui Huang, Vicky Chin, Farhannah Aly, Martin Carolan, Jonathan Sykes, Shalini K. Vinod, Geoffrey P. Delaney, and Lois Holloway. Standardising breast radiotherapy structure naming conventions: A machine learning approach. *Cancers*, 15(3), 2023.
- [65] Thilo Schuler, John Kipritidis, Thomas Eade, George Hruby, Andrew Kneebone, Mario Perez, Kylie Grimberg, Kylie Richardson, Sally Evill, Brooke Evans, and Blanca Gallego. Big data readiness in radiation oncology: An efficient approach for relabeling radiation therapy structures with their tg-263 standard name in real-world data sets. *Advances in Radiation Oncology*, 4(1):191–200, 2019.
- [66] William C. Sleeman IV, Joseph Nalluri, Khajamoinuddin Syed, Preetam Ghosh, Bartosz Krawczyk, Michael Hagan, Jatinder Palta, and Rishabh Kapoor. A machine learning method for relabeling arbitrary dicom structure sets to tg-263 defined labels. *Journal of Biomedical Informatics*, 109:103527, 2020.
- [67] David Livermore, Thomas Trappenberg, and Alasdair Syme. Machine learning for contour classification in tg-263 noncompliant databases. *Journal of Applied Clinical Medical Physics*, 23(9):e13662, 2022.
- [68] Khajamoinuddin Syed, William Sleeman IV, Kevin Ivey, Michael Hagan, Jatinder Palta, Rishabh Kapoor, and Preetam Ghosh. Integrated natural language processing and machine learning models for standardizing radiotherapy structure names. *Healthcare*, 8(2), 2020.

- 
- [69] Priyankar Bose, Pratip Rana, William C. Sleeman, Sriram Srinivasan, Rishabh Kapoor, Jatinder Palta, and Preetam Ghosh. Multimodal deep learning methods on image and textual data to predict radiotherapy structure names. *BioMedInformatics*, 3(3):493–513, 2023.
- [70] Charles S. Mayo, Jean M. Moran, Walter Bosch, Ying Xiao, Todd McNutt, Richard Popple, Jeff Michalski, Mary Feng, Lawrence B. Marks, Clifton D. Fuller, Ellen Yorke, Jatinder Palta, Peter E. Gabriel, Andrea Molineu, Martha M. Matuszak, Elizabeth Covington, Kathryn Masi, Susan L. Richardson, Timothy Ritter, Tomasz Morgas, Stella Flampouri, Lakshmi Santanam, Joseph A. Moore, Thomas G. Purdie, Robert C. Miller, Coen Hurkmans, Judy Adams, Qing-Rong Jackie Wu, Colleen J. Fox, Ramon Alfredo Siochi, Norman L. Brown, Wilko Verbakel, Yves Archambault, Steven J. Chmura, Andre L. Dekker, Don G. Eagle, Thomas J. Fitzgerald, Theodore Hong, Rishabh Kapoor, Beth Lansing, Shruti Jolly, Mary E. Napolitano, James Percy, Mark S. Rose, Salim Siddiqui, Christof Schadt, William E. Simon, William L. Straube, Sara T. St. James, Kenneth Ulin, Sue S. Yom, and Torunn I. Yock. American association of physicists in medicine task group 263: Standardizing nomenclatures in radiation oncology. *International Journal of Radiation Oncology\*Biology\*Physics*, 100(4):1057–1066, 2018.
- [71] Wenxiong Liao, Zhengliang Liu, Haixing Dai, Zihao Wu, Yiyang Zhang, Xiaoke Huang, Yuzhong Chen, Xi Jiang, Dajiang Zhu, Tianming Liu, et al. Mask-guided bert for few shot text classification. *arXiv preprint arXiv:2302.10447*, 2023.
- [72] Lin Zhao, Zihao Wu, Haixing Dai, Zhengliang Liu, Xintao Hu, Tuo Zhang, Dajiang Zhu, and Tianming Liu. A generic framework for embedding human brain function with temporally correlated autoencoder. *Medical Image Analysis*, page 102892, 2023.
- [73] Saed Rezayi, Zhengliang Liu, Zihao Wu, Chandra Dhakal, Bao Ge, Haixing Dai, Gengchen Mai, Ninghao Liu, Chen Zhen, Tianming Liu, et al. Exploring new frontiers in agricultural nlp: Investigating the potential of large language models for food applications. *arXiv preprint arXiv:2306.11892*, 2023.
- [74] Hongmin Cai, Xiaoke Huang, Zhengliang Liu, Wenxiong Liao, Haixing Dai, Zihao Wu, Dajiang Zhu, Hui Ren, Quanzheng Li, Tianming Liu, et al. Exploring multimodal approaches for alzheimer’s disease detection using patient speech transcript and audio data. *arXiv preprint arXiv:2307.02514*, 2023.
- [75] Zhengliang Liu, Xinyu He, Lei Liu, Tianming Liu, and Xiaoming Zhai. Context matters: A strategy to pre-train language model for science education. *arXiv preprint arXiv:2301.12031*, 2023.
- [76] Yuchou Chang, Zhiqiang Li, Gulfam Saju, Hui Mao, and Tianming Liu. Meta-radiology. *Meta*, 1:100001, 2023.
- [77] Tianyang Zhong, Yaonai Wei, Li Yang, Zihao Wu, Zhengliang Liu, Xiaozheng Wei, Wenjun Li, Junjie Yao, Chong Ma, Xiang Li, et al. Chatabl: Abductive learning via natural language interaction with chatgpt. *arXiv preprint arXiv:2304.11107*, 2023.
- [78] Alexander Kirillov, Eric Mintun, Nikhila Ravi, Hanzi Mao, Chloe Rolland, Laura Gustafson, Tete Xiao, Spencer Whitehead, Alexander C Berg, Wan-Yen Lo, et al. Segment anything. *arXiv preprint arXiv:2304.02643*, 2023.
- [79] Hao Zhang, Feng Li, Shilong Liu, Lei Zhang, Hang Su, Jun Zhu, Lionel M Ni, and Heung-Yeung Shum. Dino: Detr with improved denoising anchor boxes for end-to-end object detection. *arXiv preprint arXiv:2203.03605*, 2022.

- 
- [80] Maxime Oquab, Timothée Darcet, Théo Moutakanni, Huy Vo, Marc Szafraniec, Vasil Khalidov, Pierre Fernandez, Daniel Haziza, Francisco Massa, Alaaeldin El-Nouby, et al. Dinov2: Learning robust visual features without supervision. *arXiv preprint arXiv:2304.07193*, 2023.
- [81] Alexey Dosovitskiy, Lucas Beyer, Alexander Kolesnikov, Dirk Weissenborn, Xiaohua Zhai, Thomas Unterthiner, Mostafa Dehghani, Matthias Minderer, Georg Heigold, Sylvain Gelly, et al. An image is worth 16x16 words: Transformers for image recognition at scale. *arXiv preprint arXiv:2010.11929*, 2020.
- [82] Ze Liu, Yutong Lin, Yue Cao, Han Hu, Yixuan Wei, Zheng Zhang, Stephen Lin, and Baining Guo. Swin transformer: Hierarchical vision transformer using shifted windows. In *Proceedings of the IEEE/CVF international conference on computer vision*, pages 10012–10022, 2021.
- [83] Zhenxiang Xiao, Yuzhong Chen, Lu Zhang, Junjie Yao, Zihao Wu, Xiaowei Yu, Yi Pan, Lin Zhao, Chong Ma, Xinyu Liu, et al. Instruction-vit: Multi-modal prompts for instruction learning in vit. *arXiv preprint arXiv:2305.00201*, 2023.
- [84] Xiaowei Yu, Lu Zhang, Haixing Dai, Yanjun Lyu, Lin Zhao, Zihao Wu, David Liu, Tianming Liu, and Dajiang Zhu. Core-periphery principle guided redesign of self-attention in transformers. *arXiv preprint arXiv:2303.15569*, 2023.
- [85] Chong Ma, Lin Zhao, Yuzhong Chen, David Weizhong Liu, Xi Jiang, Tuo Zhang, Xintao Hu, Dinggang Shen, Dajiang Zhu, and Tianming Liu. Rectify vit shortcut learning by visual saliency. *arXiv preprint arXiv:2206.08567*, 2022.
- [86] Xiaowei Yu, Lu Zhang, Lin Zhao, Yanjun Lyu, Tianming Liu, and Dajiang Zhu. Disentangling spatial-temporal functional brain networks via twin-transformers. *arXiv preprint arXiv:2204.09225*, 2022.
- [87] Haixing Dai, Chong Ma, Zhengliang Liu, Yiwei Li, Peng Shu, Xiaozheng Wei, Lin Zhao, Zihao Wu, Dajiang Zhu, Wei Liu, et al. Samaug: Point prompt augmentation for segment anything model. *arXiv preprint arXiv:2307.01187*, 2023.
- [88] Haixing Dai, Lu Zhang, Lin Zhao, Zihao Wu, Zhengliang Liu, David Liu, Xiaowei Yu, Yanjun Lyu, Changying Li, Ninghao Liu, et al. Hierarchical semantic tree concept whitening for interpretable image classification. *arXiv preprint arXiv:2307.04343*, 2023.
- [89] Lian Zhang, Jason M Holmes, Zhengliang Liu, Sujay A Vora, Terence T Sio, Carlos E Vargas, Nathan Y Yu, Sameer R Keole, Steven E Schild, Martin Bues, et al. Beam mask and sliding window-facilitated deep learning-based accurate and efficient dose prediction for pencil beam scanning proton therapy. *arXiv preprint arXiv:2305.18572*, 2023.
- [90] Xia-An Bi, Ke Chen, Siyu Jiang, Sheng Luo, Wenyan Zhou, Zhaoxu Xing, Luyun Xu, Zhengliang Liu, and Tianming Liu. Community graph convolution neural network for alzheimer’s disease classification and pathogenetic factors identification. *IEEE Transactions on Neural Networks and Learning Systems*, 2023.
- [91] Shu Zhang, Enze Shi, Lin Wu, Ruoyang Wang, Sigang Yu, Zhengliang Liu, Shaochen Xu, Tianming Liu, and Shijie Zhao. Differentiating brain states via multi-clip random fragment strategy-based interactive bidirectional recurrent neural network. *Neural Networks*, 165:1035–1049, 2023.
- [92] Y Ding, Z Liu, H Feng, J Holmes, Y Yang, N Yu, T Sio, S Schild, B Li, and W Liu. Accurate and efficient deep neural network based deformable image registration method in lung cancer.

---

In *MEDICAL PHYSICS*, volume 49, pages E148–E148. WILEY 111 RIVER ST, HOBOKEN 07030-5774, NJ USA, 2022.

- [93] Yiheng Liu, Enjie Ge, Mengshen He, Zhengliang Liu, Shijie Zhao, Xintao Hu, Dajiang Zhu, Tianming Liu, and Bao Ge. Discovering dynamic functional brain networks via spatial and channel-wise attention. *arXiv preprint arXiv:2205.09576*, 2022.
- [94] Haixing Dai, Qing Li, Lin Zhao, Liming Pan, Cheng Shi, Zhengliang Liu, Zihao Wu, Lu Zhang, Shijie Zhao, Xia Wu, et al. Graph representation neural architecture search for optimal spatial/temporal functional brain network decomposition. In *International Workshop on Machine Learning in Medical Imaging*, pages 279–287. Springer, 2022.
- [95] Nanyi Fei, Zhiwu Lu, Yizhao Gao, Guoxing Yang, Yuqi Huo, Jingyuan Wen, Haoyu Lu, Ruihua Song, Xin Gao, Tao Xiang, et al. Towards artificial general intelligence via a multimodal foundation model. *Nature Communications*, 13(1):3094, 2022.
- [96] Haotian Liu, Chunyuan Li, Qingsyang Wu, and Yong Jae Lee. Visual instruction tuning. *arXiv preprint arXiv:2304.08485*, 2023.
- [97] Wenhui Wang, Zhe Chen, Xiaokang Chen, Jiannan Wu, Xizhou Zhu, Gang Zeng, Ping Luo, Tong Lu, Jie Zhou, Yu Qiao, et al. Visionllm: Large language model is also an open-ended decoder for vision-centric tasks. *arXiv preprint arXiv:2305.11175*, 2023.
- [98] Feilong Chen, Minglun Han, Haozhi Zhao, Qingsyang Zhang, Jing Shi, Shuang Xu, and Bo Xu. X-llm: Bootstrapping advanced large language models by treating multi-modalities as foreign languages. *arXiv preprint arXiv:2305.04160*, 2023.
- [99] Zhiyang Xu, Ying Shen, and Lifu Huang. Multiinstruct: Improving multi-modal zero-shot learning via instruction tuning. *arXiv preprint arXiv:2212.10773*, 2022.
- [100] Deyao Zhu, Jun Chen, Xiaoqian Shen, Xiang Li, and Mohamed Elhoseiny. Minigpt-4: Enhancing vision-language understanding with advanced large language models. *arXiv preprint arXiv:2304.10592*, 2023.
- [101] Zijia Zhao, Longteng Guo, Tongtian Yue, Sihan Chen, Shuai Shao, Xinxin Zhu, Zehuan Yuan, and Jing Liu. Chatbridge: Bridging modalities with large language model as a language catalyst. *arXiv preprint arXiv:2305.16103*, 2023.
- [102] KunChang Li, Yinan He, Yi Wang, Yizhuo Li, Wenhui Wang, Ping Luo, Yali Wang, Limin Wang, and Yu Qiao. Videochat: Chat-centric video understanding. *arXiv preprint arXiv:2305.06355*, 2023.
- [103] Rui Yang, Lin Song, Yanwei Li, Sijie Zhao, Yixiao Ge, Xiu Li, and Ying Shan. Gpt4tools: Teaching large language model to use tools via self-instruction. *arXiv preprint arXiv:2305.18752*, 2023.
- [104] Philippe Lambin, Ruud GPM Van Stiphout, Maud HW Starmans, Emmanuel Rios-Velazquez, Georgi Nalbantov, Hugo JW Aerts, Erik Roelofs, Wouter Van Elmpt, Paul C Boutros, Pierluigi Granone, et al. Predicting outcomes in radiation oncology—multifactorial decision support systems. *Nature reviews Clinical oncology*, 10(1):27–40, 2013.
- [105] Hongying Feng, Samir H Patel, William W Wong, James E Younkin, Gregory P Penoncello, Danairis Hernandez Morales, Joshua B Stoker, Daniel G Robertson, Mirek Fatyga, Martin

- 
- Bues, et al. Gpu-accelerated monte carlo-based online adaptive proton therapy: A feasibility study. *Medical physics*, 49(6):3550–3563, 2022.
- [106] Sunan Cui, Andrew Hope, Thomas J Dilling, Laura A Dawson, Randall Ten Haken, and Issam El Naqa. Artificial intelligence for outcome modeling in radiotherapy. In *Seminars in Radiation Oncology*, volume 32, pages 351–364. Elsevier, 2022.
- [107] Khadijeh Saednia, Sami Tabbarah, Andrew Lagree, Tina Wu, Jonathan Klein, Eduardo Garcia, Michael Hall, Edward Chow, Eileen Rakovitch, Charmaine Childs, et al. Quantitative thermal imaging biomarkers to detect acute skin toxicity from breast radiation therapy using supervised machine learning. *International Journal of Radiation Oncology\* Biology\* Physics*, 106(5):1071–1083, 2020.
- [108] Sunan Cui, Randall K Ten Haken, and Issam El Naqa. Integrating multiomics information in deep learning architectures for joint actuarial outcome prediction in non-small cell lung cancer patients after radiation therapy. *International Journal of Radiation Oncology\* Biology\* Physics*, 110(3):893–904, 2021.
- [109] Lise Wei, Dawn Owen, Benjamin Rosen, Xinzhou Guo, Kyle Cuneo, Theodore S Lawrence, Randall Ten Haken, and Issam El Naqa. A deep survival interpretable radiomics model of hepatocellular carcinoma patients. *Physica Medica*, 82:295–305, 2021.
- [110] Thomas Bortfeld. Imrt: a review and preview. *Physics in Medicine & Biology*, 51(13):R363, 2006.
- [111] M Kara Bucci, Alison Bevan, and Mack Roach III. Advances in radiation therapy: conventional to 3d, to imrt, to 4d, and beyond. *CA: a cancer journal for clinicians*, 55(2):117–134, 2005.
- [112] Wei Liu, Xiaodong Zhang, Yupeng Li, and Radhe Mohan. Robust optimization of intensity modulated proton therapy. *Medical physics*, 39(2):1079–1091, 2012.
- [113] Wei Liu, Steven E Schild, Joe Y Chang, Zhongxing Liao, Yu-Hui Chang, Zhifei Wen, Jiajian Shen, Joshua B Stoker, Xiaoning Ding, Yanle Hu, et al. Exploratory study of 4d versus 3d robust optimization in intensity modulated proton therapy for lung cancer. *International Journal of Radiation Oncology\* Biology\* Physics*, 95(1):523–533, 2016.
- [114] Wei Liu, Yupeng Li, Xiaoqiang Li, Wenhua Cao, and Xiaodong Zhang. Influence of robust optimization in intensity-modulated proton therapy with different dose delivery techniques. *Medical physics*, 39(6Part1):3089–3101, 2012.
- [115] Wei Liu, Steven J Frank, Xiaoqiang Li, Yupeng Li, Ron X Zhu, and Radhe Mohan. Ptv-based impt optimization incorporating planning risk volumes vs robust optimization. *Medical physics*, 40(2):021709, 2013.
- [116] Dan Nguyen, Mu-Han Lin, David Sher, Weiguo Lu, Xun Jia, and Steve Jiang. Advances in automated treatment planning. In *Seminars in Radiation Oncology*, volume 32, pages 343–350. Elsevier, 2022.
- [117] Jason Matney, Peter C Park, Jaques Bluett, Yi Pei Chen, Wei Liu, Laurence E Court, Zhongxing Liao, Heng Li, and Radhe Mohan. Effects of respiratory motion on passively scattered proton therapy versus intensity modulated photon therapy for stage iii lung cancer: are proton plans more sensitive to breathing motion? *International Journal of Radiation Oncology\* Biology\* Physics*, 87(3):576–582, 2013.

- 
- [118] Chenbin Liu, Steven E Schild, Joe Y Chang, Zhongxing Liao, Shawn Korte, Jiajian Shen, Xiaoning Ding, Yanle Hu, Yixiu Kang, Sameer R Keole, et al. Impact of spot size and spacing on the quality of robustly optimized intensity modulated proton therapy plans for lung cancer. *International Journal of Radiation Oncology\* Biology\* Physics*, 101(2):479–489, 2018.
- [119] Yu An, Jie Shan, Samir H Patel, William Wong, Steven E Schild, Xiaoning Ding, Martin Bues, and Wei Liu. Robust intensity-modulated proton therapy to reduce high linear energy transfer in organs at risk. *Medical physics*, 44(12):6138–6147, 2017.
- [120] Chenbin Liu, Samir H Patel, Jie Shan, Steven E Schild, Carlos E Vargas, William W Wong, Xiaoning Ding, Martin Bues, and Wei Liu. Robust optimization for intensity modulated proton therapy to redistribute high linear energy transfer from nearby critical organs to tumors in head and neck cancer. *International Journal of Radiation Oncology\* Biology\* Physics*, 107(1):181–193, 2020.
- [121] Olga L Green, Lauren E Henke, and Geoffrey D Hugo. Practical clinical workflows for online and offline adaptive radiation therapy. In *Seminars in radiation oncology*, volume 29, pages 219–227. Elsevier, 2019.
- [122] Qingrong Jackie Wu, Taoran Li, Qiuwen Wu, and Fang-Fang Yin. Adaptive radiation therapy: technical components and clinical applications. *The cancer journal*, 17(3):182–189, 2011.
- [123] Shu Haseai, Hidetaka Arimura, Kaori Asai, Tadamasa Yoshitake, and Yoshiyuki Shioyama. Similar-cases-based planning approaches with beam angle optimizations using water equivalent path length for lung stereotactic body radiation therapy. *Radiological physics and technology*, 13:119–127, 2020.
- [124] Chris McIntosh and Thomas G Purdie. Contextual atlas regression forests: multiple-atlas-based automated dose prediction in radiation therapy. *IEEE transactions on medical imaging*, 35(4):1000–1012, 2015.
- [125] Yang Sheng, Taoran Li, You Zhang, W Robert Lee, Fang-Fang Yin, Yaorong Ge, and Q Jackie Wu. Atlas-guided prostate intensity modulated radiation therapy (imrt) planning. *Physics in Medicine & Biology*, 60(18):7277, 2015.
- [126] Xiaofeng Zhu, Yaorong Ge, Taoran Li, Danhai Thongphiew, Fang-Fang Yin, and Q Jackie Wu. A planning quality evaluation tool for prostate adaptive imrt based on machine learning. *Medical physics*, 38(2):719–726, 2011.
- [127] Justin J Boutilier, Taewoo Lee, Tim Craig, Michael B Sharpe, and Timothy CY Chan. Models for predicting objective function weights in prostate cancer imrt. *Medical physics*, 42(4):1586–1595, 2015.
- [128] Jiahua Zhang, Q Jackie Wu, Tianyi Xie, Yang Sheng, Fang-Fang Yin, and Yaorong Ge. An ensemble approach to knowledge-based intensity-modulated radiation therapy planning. *Frontiers in oncology*, 8:57, 2018.
- [129] Changsheng Ma and Fujing Huang. Assessment of a knowledge-based rapidplan model for patients with postoperative cervical cancer. *Precision Radiation Oncology*, 1(3):102–107, 2017.
- [130] Jiawei Fan, Jiazhou Wang, Zhi Chen, Chaosu Hu, Zhen Zhang, and Weigang Hu. Automatic treatment planning based on three-dimensional dose distribution predicted from deep learning technique. *Medical physics*, 46(1):370–381, 2019.

- 
- [131] Zhiqiang Liu, Jiawei Fan, Minghui Li, Hui Yan, Zhihui Hu, Peng Huang, Yuan Tian, Junjie Miao, and Jianrong Dai. A deep learning method for prediction of three-dimensional dose distribution of helical tomotherapy. *Medical physics*, 46(5):1972–1983, 2019.
- [132] Dan Nguyen, Xun Jia, David Sher, Mu-Han Lin, Zohaib Iqbal, Hui Liu, and Steve Jiang. 3d radiotherapy dose prediction on head and neck cancer patients with a hierarchically densely connected u-net deep learning architecture. *Physics in medicine & Biology*, 64(6):065020, 2019.
- [133] Roya Norouzi Kandalan, Dan Nguyen, Nima Hassan Rezaeian, Ana M Barragán-Montero, Sebastiaan Breedveld, Kamesh Namuduri, Steve Jiang, and Mu-Han Lin. Dose prediction with deep learning for prostate cancer radiation therapy: model adaptation to different treatment planning practices. *Radiotherapy and Oncology*, 153:228–235, 2020.
- [134] Roel GJ Kierkels, Albin Fredriksson, Stefan Both, Johannes A Langendijk, Daniel Scandurra, and Erik W Korevaar. Automated robust proton planning using dose-volume histogram-based mimicking of the photon reference dose and reducing organ at risk dose optimization. *International Journal of Radiation Oncology\* Biology\* Physics*, 103(1):251–258, 2019.
- [135] Philippe Meyer, M-C Biston, C Khamphan, T Marghani, J Mazurier, V Bodez, L Fezzani, PA Rigaud, G Sidorski, L Simon, et al. Automation in radiotherapy treatment planning: Examples of use in clinical practice and future trends for a complete automated workflow. *Cancer/Radiothérapie*, 25(6-7):617–622, 2021.
- [136] Berkman Sahiner, Aria Pezeshk, Lubomir M Hadjiiski, Xiaosong Wang, Karen Drukker, Kenny H Cha, Ronald M Summers, and Maryellen L Giger. Deep learning in medical imaging and radiation therapy. *Medical physics*, 46(1):e1–e36, 2019.
- [137] Ben Goertzel. Artificial general intelligence: concept, state of the art, and future prospects. *Journal of Artificial General Intelligence*, 5(1):1, 2014.
- [138] Julianne M Pollard, Zhifei Wen, Ramaswamy Sadagopan, Jihong Wang, and Geoffrey S Ibbott. The future of image-guided radiotherapy will be mr guided. *The British journal of radiology*, 90(1073):20160667, 2017.
- [139] Joao Seco and Phil M Evans. Assessing the effect of electron density in photon dose calculations. *Medical physics*, 33(2):540–552, 2006.
- [140] Yingzi Liu, Yang Lei, Tonghe Wang, Yabo Fu, Xiangyang Tang, Walter J Curran, Tian Liu, Pretesh Patel, and Xiaofeng Yang. Cbct-based synthetic ct generation using deep-attention cyclegan for pancreatic adaptive radiotherapy. *Medical physics*, 47(6):2472–2483, 2020.
- [141] Yingzi Liu, Yang Lei, Tonghe Wang, Oluwatosin Kayode, Sibo Tian, Tian Liu, Pretesh Patel, Walter J Curran, Lei Ren, and Xiaofeng Yang. Mri-based treatment planning for liver stereotactic body radiotherapy: validation of a deep learning-based synthetic ct generation method. *The British Journal of Radiology*, 92(1100):20190067, 2019.
- [142] Jun Zhao, Zhi Chen, Jiazhou Wang, Fan Xia, Jiayuan Peng, Yiwen Hu, Weigang Hu, and Zhen Zhang. Mv cbct-based synthetic ct generation using a deep learning method for rectal cancer adaptive radiotherapy. *Frontiers in Oncology*, 11:655325, 2021.
- [143] Jie Shan, Hongying Feng, Danairis Hernandez Morales, Samir H Patel, William W Wong, Mirek Fatyga, Martin Bues, Steven E Schild, Robert L Foote, and Wei Liu. Virtual particle monte carlo: A new concept to avoid simulating secondary particles in proton therapy dose calculation. *Medical physics*, 49(10):6666–6683, 2022.

- 
- [144] Emily Johnstone, Jonathan J. Wyatt, Ann M. Henry, Susan C. Short, David Sebag-Montefiore, Louise Murray, Charles G. Kelly, Hazel M. McCallum, and Richard Speight. Systematic review of synthetic computed tomography generation methodologies for use in magnetic resonance imaging-only radiation therapy. *International Journal of Radiation Oncology\*Biology\*Physics*, 100(1):199–217, 2018.
- [145] Davide Cusumano, Jacopo Lenkowicz, Claudio Votta, Luca Boldrini, Lorenzo Placidi, Francesco Catucci, Nicola Dinapoli, Marco Valerio Antonelli, Angela Romano, Viola De Luca, et al. A deep learning approach to generate synthetic ct in low field mr-guided adaptive radiotherapy for abdominal and pelvic cases. *Radiotherapy and Oncology*, 153:205–212, 2020.
- [146] Matteo Maspero, Laura G Bentvelzen, Mark HF Savenije, Filipa Guerreiro, Enrica Seravalli, Geert O Janssens, Cornelis AT van den Berg, and Marielle EP Philippens. Deep learning-based synthetic ct generation for paediatric brain mr-only photon and proton radiotherapy. *Radiotherapy and Oncology*, 153:197–204, 2020.
- [147] David Bird, Michael G Nix, Hazel McCallum, Mark Teo, Alexandra Gilbert, Nathalie Casanova, Rachel Cooper, David L Buckley, David Sebag-Montefiore, Richard Speight, et al. Multicentre, deep learning, synthetic-ct generation for ano-rectal mr-only radiotherapy treatment planning. *Radiotherapy and Oncology*, 156:23–28, 2021.
- [148] Jacopo Lenkowicz, Claudio Votta, Matteo Nardini, Flaviovincenzo Quaranta, Francesco Catucci, Luca Boldrini, Marica Vagni, Sebastiano Menna, Lorenzo Placidi, Angela Romano, et al. A deep learning approach to generate synthetic ct in low field mr-guided radiotherapy for lung cases. *Radiotherapy and Oncology*, 176:31–38, 2022.
- [149] Ghazal Shafai-Erfani, Tonghe Wang, Yang Lei, Sibo Tian, Pretesh Patel, Ashesh B Jani, Walter J Curran, Tian Liu, and Xiaofeng Yang. Dose evaluation of mri-based synthetic ct generated using a machine learning method for prostate cancer radiotherapy. *Medical Dosimetry*, 44(4):e64–e70, 2019.
- [150] Matteo Maspero, Mark HF Savenije, Anna M Dinkla, Peter R Seevinck, Martijn PW Intven, Ina M Jurgenliemk-Schulz, Linda GW Kerkmeijer, and Cornelis AT van den Berg. Dose evaluation of fast synthetic-ct generation using a generative adversarial network for general pelvis mr-only radiotherapy. *Physics in Medicine & Biology*, 63(18):185001, 2018.
- [151] Xiao Han. Mr-based synthetic ct generation using a deep convolutional neural network method. *Medical physics*, 44(4):1408–1419, 2017.
- [152] Fang Liu, Poonam Yadav, Andrew M Baschnagel, and Alan B McMillan. Mr-based treatment planning in radiation therapy using a deep learning approach. *Journal of applied clinical medical physics*, 20(3):105–114, 2019.
- [153] Lianli Liu, Adam Johansson, Yue Cao, Janell Dow, Theodore S Lawrence, and James M Balter. Abdominal synthetic ct generation from mr dixon images using a u-net trained with ‘semi-synthetic’ct data. *Physics in Medicine & Biology*, 65(12):125001, 2020.
- [154] Abbas Bahrami, Alireza Karimian, and Hossein Arabi. Comparison of different deep learning architectures for synthetic ct generation from mr images. *Physica Medica*, 90:99–107, 2021.
- [155] Maria Francesca Spadea, Giampaolo Pileggi, Paolo Zaffino, Patrick Salome, Ciprian Catana, David Izquierdo-Garcia, Francesco Amato, and Joao Seco. Deep convolution neural network (dcnn) multiplane approach to synthetic ct generation from mr images—application in brain

- 
- proton therapy. *International Journal of Radiation Oncology\* Biology\* Physics*, 105(3):495–503, 2019.
- [156] Lei Xiang, Qian Wang, Dong Nie, Lichi Zhang, Xiayao Jin, Yu Qiao, and Dinggang Shen. Deep embedding convolutional neural network for synthesizing ct image from t1-weighted mr image. *Medical image analysis*, 47:31–44, 2018.
- [157] Anna M Dinkla, Mateusz C Florkow, Matteo Maspero, Mark HF Savenije, Frank Zijlstra, Patricia AH Doornaert, Marijn van Stralen, Marielle EP Philippens, Cornelis AT van den Berg, and Peter R Seevinck. Dosimetric evaluation of synthetic ct for head and neck radiotherapy generated by a patch-based three-dimensional convolutional neural network. *Medical physics*, 46(9):4095–4104, 2019.
- [158] Bin Tang, Fan Wu, Yuchuan Fu, Xianliang Wang, Pei Wang, Lucia Clara Orlandini, Jie Li, and Qing Hou. Dosimetric evaluation of synthetic ct image generated using a neural network for mr-only brain radiotherapy. *Journal of Applied Clinical Medical Physics*, 22(3):55–62, 2021.
- [159] Bo Zhao, Tingting Cheng, Xueren Zhang, Jingjing Wang, Hong Zhu, Rongchang Zhao, Dengwang Li, Zijian Zhang, and Gang Yu. Ct synthesis from mr in the pelvic area using residual transformer conditional gan. *Computerized Medical Imaging and Graphics*, 103:102150, 2023.
- [160] Yao Zhao, He Wang, Cenji Yu, Laurence E Court, Xin Wang, Qianxia Wang, Tinsu Pan, Yao Ding, Jack Phan, and Jinzhong Yang. Compensation cycle consistent generative adversarial networks (comp-gan) for synthetic ct generation from mr scans with truncated anatomy. *Medical physics*, 2023.
- [161] Seung Kwan Kang, Hyun Joon An, Hyeongmin Jin, Jung-in Kim, Eui Kyu Chie, Jong Min Park, and Jae Sung Lee. Synthetic ct generation from weakly paired mr images using cycle-consistent gan for mr-guided radiotherapy. *Biomedical engineering letters*, 11(3):263–271, 2021.
- [162] Alaa Abu-Srhan, Israa Almallahi, Mohammad AM Abushariah, Waleed Mahafza, and Omar S Al-Kadi. Paired-unpaired unsupervised attention guided gan with transfer learning for bidirectional brain mr-ct synthesis. *Computers in Biology and Medicine*, 136:104763, 2021.
- [163] Guodong Zeng and Guoyan Zheng. Hybrid generative adversarial networks for deep mr to ct synthesis using unpaired data. In *Medical Image Computing and Computer Assisted Intervention—MICCAI 2019: 22nd International Conference, Shenzhen, China, October 13–17, 2019, Proceedings, Part IV 22*, pages 759–767. Springer, 2019.
- [164] Jiayuan Wang, QM Jonathan Wu, and Farhad Pourpanah. Dc-cyclegan: bidirectional ct-to-mr synthesis from unpaired data. *Computerized Medical Imaging and Graphics*, page 102249, 2023.
- [165] Kévin ND Brou Boni, John Klein, Akos Gulyban, Nick Reynaert, and David Pasquier. Improving generalization in mr-to-ct synthesis in radiotherapy by using an augmented cycle generative adversarial network with unpaired data. *Medical physics*, 48(6):3003–3010, 2021.
- [166] Anders B Olin, Christopher Thomas, Adam E Hansen, Jacob H Rasmussen, Georgios Krokos, Teresa Guerrero Urbano, Andriana Michaelidou, Björn Jakoby, Claes N Ladefoged, Anne K Berthelsen, et al. Robustness and generalizability of deep learning synthetic computed tomography for positron emission tomography/magnetic resonance imaging–based radiation therapy planning of patients with head and neck cancer. *Advances in radiation oncology*, 6(6):100762, 2021.

- 
- [167] Shu-Hui Hsu, Zhaohui Han, Jonathan E Leeman, Yue-Houng Hu, Raymond H Mak, and Atchar Sudhyadhom. Synthetic ct generation for mri-guided adaptive radiotherapy in prostate cancer. *Frontiers in Oncology*, 12:969463, 2022.
- [168] Emily Johnstone, Jonathan J Wyatt, Ann M Henry, Susan C Short, David Sebag-Montefiore, Louise Murray, Charles G Kelly, Hazel M McCallum, and Richard Speight. Systematic review of synthetic computed tomography generation methodologies for use in magnetic resonance imaging–only radiation therapy. *International Journal of Radiation Oncology\* Biology\* Physics*, 100(1):199–217, 2018.
- [169] Seungjong Oh and Siyong Kim. Deformable image registration in radiation therapy. *Radiation oncology journal*, 35(2):101, 2017.
- [170] Maria Thor, Jørgen BB Petersen, Lise Bentzen, Morten Høyer, and Ludvig Paul Muren. Deformable image registration for contour propagation from ct to cone-beam ct scans in radiotherapy of prostate cancer. *Acta Oncologica*, 50(6):918–925, 2011.
- [171] Gilion Hautvast, Steven Lobregt, Marcel Breeuwer, and Frans Gerritsen. Automatic contour propagation in cine cardiac magnetic resonance images. *IEEE transactions on medical imaging*, 25(11):1472–1482, 2006.
- [172] M Söhn, M Birkner, D Yan, and M Alber. Modelling individual geometric variation based on dominant eigenmodes of organ deformation: implementation and evaluation. *Physics in Medicine & Biology*, 50(24):5893, 2005.
- [173] T-N Nguyen, JL Moseley, LA Dawson, DA Jaffray, and KK Brock. Adapting liver motion models using a navigator channel technique. *Medical physics*, 36(4):1061–1073, 2009.
- [174] E Budiarto, M Keijzer, PR Storchi, MS Hoogeman, L Bondar, TF Mutanga, HCJ de Boer, and AW Heemink. A population-based model to describe geometrical uncertainties in radiotherapy: applied to prostate cases. *Physics in Medicine & Biology*, 56(4):1045, 2011.
- [175] Seungjong Oh, David Jaffray, and Young-Bin Cho. A novel method to quantify and compare anatomical shape: application in cervix cancer radiotherapy. *Physics in Medicine & Biology*, 59(11):2687, 2014.
- [176] Thomas Guerrero, Kevin Sanders, Josue Noyola-Martinez, Edward Castillo, Yin Zhang, Richard Tapia, Rudy Guerra, Yerko Borghero, and Ritsuko Komaki. Quantification of regional ventilation from treatment planning ct. *International Journal of Radiation Oncology\* Biology\* Physics*, 62(3):630–634, 2005.
- [177] Brian P Yaremko, Thomas M Guerrero, Josue Noyola-Martinez, Rudy Guerra, David G Lege, Linda T Nguyen, Peter A Balter, James D Cox, and Ritsuko Komaki. Reduction of normal lung irradiation in locally advanced non–small-cell lung cancer patients, using ventilation images for functional avoidance. *International Journal of Radiation Oncology\* Biology\* Physics*, 68(2):562–571, 2007.
- [178] Tokihiro Yamamoto, Sven Kabus, Jens Von Berg, Cristian Lorenz, and Paul J Keall. Impact of four-dimensional computed tomography pulmonary ventilation imaging-based functional avoidance for lung cancer radiotherapy. *International Journal of Radiation Oncology\* Biology\* Physics*, 79(1):279–288, 2011.
- [179] X Sharon Qi, Anand Santhanam, John Neylon, Yugang Min, Tess Armstrong, Ke Sheng, Robert J Staton, Jason Pukala, Andrew Pham, Daniel A Low, et al. Near real-time assessment of

- 
- anatomic and dosimetric variations for head and neck radiation therapy via graphics processing unit-based dose deformation framework. *International Journal of Radiation Oncology\* Biology\* Physics*, 92(2):415–422, 2015.
- [180] Manju Sharma, Elisabeth Weiss, and Jeffrey V Siebers. Dose deformation-invariance in adaptive prostate radiation therapy: implication for treatment simulations. *Radiotherapy and oncology*, 105(2):207–213, 2012.
- [181] Michael Velec, Joanne L Moseley, Cynthia L Eccles, Tim Craig, Michael B Sharpe, Laura A Dawson, and Kristy K Brock. Effect of breathing motion on radiotherapy dose accumulation in the abdomen using deformable registration. *International Journal of Radiation Oncology\* Biology\* Physics*, 80(1):265–272, 2011.
- [182] Di Yan, Frank Vicini, John Wong, and Alvaro Martinez. Adaptive radiation therapy. *Physics in Medicine & Biology*, 42(1):123, 1997.
- [183] B Schaly, JA Kempe, GS Bauman, JJ Battista, and J Van Dyk. Tracking the dose distribution in radiation therapy by accounting for variable anatomy. *Physics in Medicine & Biology*, 49(5):791, 2004.
- [184] Gary E Christensen, Blake Carlson, KS Clifford Chao, Pen Yin, Perry W Grigsby, Kim Nguyen, James F Dempsey, Fritz A Lerma, Kyongtae T Bae, Michael W Vannier, et al. Image-based dose planning of intracavitary brachytherapy: registration of serial-imaging studies using deformable anatomic templates. *International Journal of Radiation Oncology\* Biology\* Physics*, 51(1):227–243, 2001.
- [185] Lian Zhang, Zhi Wang, Chengyu Shi, Tengfei Long, and X George Xu. The impact of robustness of deformable image registration on contour propagation and dose accumulation for head and neck adaptive radiotherapy. *Journal of applied clinical medical physics*, 19(4):185–194, 2018.
- [186] Di Yan, DA Jaffray, and JW Wong. A model to accumulate fractionated dose in a deforming organ. *International Journal of Radiation Oncology\* Biology\* Physics*, 44(3):665–675, 1999.
- [187] Tom Vercauteren, Xavier Pennec, Aymeric Perchant, and Nicholas Ayache. Non-parametric diffeomorphic image registration with the demons algorithm. In *International Conference on Medical Image Computing and Computer-Assisted Intervention*, pages 319–326. Springer, 2007.
- [188] Hualiang Zhong, Jinkoo Kim, Haisen Li, Teamour Nurushev, Benjamin Movsas, and Indrin J Chetty. A finite element method to correct deformable image registration errors in low-contrast regions. *Physics in Medicine & Biology*, 57(11):3499, 2012.
- [189] Xuejun Gu, Bin Dong, Jing Wang, John Yordy, Loren Mell, Xun Jia, and Steve B Jiang. A contour-guided deformable image registration algorithm for adaptive radiotherapy. *Physics in Medicine & Biology*, 58(6):1889, 2013.
- [190] Sajendra Nithiananthan, Sebastian Schafer, Daniel J Mirota, J Webster Stayman, Wojciech Zbijewski, Douglas D Reh, Gary L Gallia, and Jeffrey H Siewerdsen. Extra-dimensional demons: a method for incorporating missing tissue in deformable image registration. *Medical physics*, 39(9):5718–5731, 2012.
- [191] Xiao Yang, Roland Kwitt, and Marc Niethammer. Fast predictive image registration. In *Deep Learning and Data Labeling for Medical Applications: First International Workshop, LABELS 2016, and Second International Workshop, DLMIA 2016, Held in Conjunction with MICCAI 2016, Athens, Greece, October 21, 2016, Proceedings 1*, pages 48–57. Springer, 2016.

- 
- [192] Xiaohuan Cao, Jianhuan Yang, Li Wang, Zhong Xue, Qian Wang, and Dinggang Shen. Deep learning based inter-modality image registration supervised by intra-modality similarity. In *Machine Learning in Medical Imaging: 9th International Workshop, MLMI 2018, Held in Conjunction with MICCAI 2018, Granada, Spain, September 16, 2018, Proceedings* 9, pages 55–63. Springer, 2018.
- [193] Jorge Onieva Onieva, Berta Martí-Fuster, María Pedrero de la Puente, and Raúl San José Estépar. Diffeomorphic lung registration using deep cnns and reinforced learning. In *International Workshop on Reconstruction and Analysis of Moving Body Organs*, pages 284–294. Springer, 2018.
- [194] Jun Lv, Ming Yang, Jue Zhang, and Xiaoying Wang. Respiratory motion correction for free-breathing 3d abdominal mri using cnn-based image registration: a feasibility study. *The British journal of radiology*, 91(XXXX):20170788, 2018.
- [195] Guha Balakrishnan, Amy Zhao, Mert R Sabuncu, John Guttag, and Adrian V Dalca. Voxel-morph: a learning framework for deformable medical image registration. *IEEE transactions on medical imaging*, 38(8):1788–1800, 2019.
- [196] Frano Rajič, Lei Ke, Yu-Wing Tai, Chi-Keung Tang, Martin Danelljan, and Fisher Yu. Segment anything meets point tracking. *arXiv preprint arXiv:2307.01197*, 2023.
- [197] Thilo Schuler, John Kipritidis, Thomas Eade, George Hruby, Andrew Kneebone, Mario Perez, Kylie Grimberg, Kylie Richardson, Sally Evill, Brooke Evans, et al. Big data readiness in radiation oncology: an efficient approach for relabeling radiation therapy structures with their tg-263 standard name in real-world data sets. *Advances in radiation oncology*, 4(1):191–200, 2019.
- [198] Joseph S Ross, Richard Lehman, and Cary P Gross. The importance of clinical trial data sharing: toward more open science. *Circulation: Cardiovascular Quality and Outcomes*, 5(2):238–240, 2012.
- [199] Wing-Yan Ip, Fu-Ki Yeung, Shang-Peng Felix Yung, Hong-Cheung Jeffrey Yu, Tsz-Him So, and Varut Vardhanabhuti. Current landscape and potential future applications of artificial intelligence in medical physics and radiotherapy. *Artificial Intelligence in Medical Imaging*, 2021.