

Algebra  
from the context of the course  
MTH 418H: Honors Algebra

Kaedon Cleland-Host

December 15, 2021

# Contents

<b>1</b>	<b>Group Theory</b>	<b>2</b>
1.1	Groups . . . . .	2
1.2	Subgroups . . . . .	2
1.2.1	Greatest Common Divisor . . . . .	2
1.2.2	Least Common Multiple . . . . .	3
1.2.3	Cyclic Groups . . . . .	3
1.3	Homomorphisms . . . . .	3
1.4	Cosets . . . . .	4
1.4.5	Counting Formula . . . . .	4
1.4.6	Lagrange's Theorem . . . . .	4
1.5	Normal Subgroups . . . . .	4
1.6	Quotient Groups . . . . .	4
1.6.3	Correspondence Theorem . . . . .	4
1.7	Product Groups . . . . .	4
1.7.3	Multiplication Isomorphism . . . . .	4
1.7.5	First Isomorphism Theorem . . . . .	5
1.8	Group Actions . . . . .	5
1.8.6	Orbit Stabilizer Theorem . . . . .	5
1.9	Conjugation . . . . .	6
1.10	p-Groups . . . . .	6
1.10.3	Fixed Point Theorem . . . . .	6
1.10.7	First Sylow Theorem . . . . .	6
1.10.9	Second Sylow Theorem . . . . .	6
1.10.11	Third Sylow Theorem . . . . .	6
<b>2</b>	<b>Ring Theory</b>	<b>7</b>
2.1	Rings . . . . .	7
2.2	Ring Homomorphisms . . . . .	7
2.3	Ideals . . . . .	7
2.4	Quotient Rings . . . . .	7
2.5	Maximal Ideals . . . . .	7
2.6	Algebraic Geometry . . . . .	7

# Chapter 1

## Group Theory

### 1.1 Groups

**Definition 1.1.1.** A **law of composition** is a map  $S^2 \rightarrow S$ .

*Remark.* We will use the notation  $ab$  for the elements of  $S$  obtained as  $a, b \rightarrow ab$ . This element is the product of  $a$  and  $b$ .

**Definition 1.1.2.** A **group** is a set  $G$  together with a law of composition that has the following three properties:

1. **Identity** There exists an element  $1 \in G$  such that  $1a = a1 = A$  for all  $a \in G$ .
2. **Associativity**  $(ab)c = a(bc)$  for all  $a, b, c \in G$ .
3. **Inverse** For any  $a \in G$ , there exists  $a^{-1} \in G$  such that  $aa^{-1} = a^{-1}a = 1$ .

**Definition 1.1.3.** An **abelian group** is a group with a commutative law of composition. That is for any  $a, b \in G$ ,  $ab = ba$ .

**Definition 1.1.4.** The **order** of a group  $G$  is the cardinality of the set.

**Proposition 1.1.5. Cancellation Law** For  $a, b, c \in G$  if  $ab = ac$  then  $b = c$ .

**Proposition 1.1.6.** Let  $S$  be a set with an associative law of composition and an identity. The subset of elements of  $S$  that are invertible forms a group.

### 1.2 Subgroups

**Definition 1.2.1.** A group  $H$  is a **Subgroup** of  $G$  if  $H$  is subset of  $G$ ,  $H$  has the same law of composition as  $G$ , and  $H$  is also a group. In other words  $H$  a group if it is a subset of  $G$  with the following properties:

1. **Closure**  $a, b \in H$  then  $ab \in H$ .
2. **Identity**  $1 \in H$ .
3. **Inverse** For all  $a \in H$ ,  $a^{-1} \in H$ .

**Definition 1.2.2.** A subgroup  $S$  of  $G$  is a **proper subgroup** if  $S \neq G$  and  $S \neq \{\mathbb{I}\}$ .

**Proposition 1.2.3.** If  $H$  and  $K$  are subgroup of  $G$ , then  $H \cap K$  is a subgroup.

**Theorem 1.2.4.** If  $S$  is a subgroup of  $\mathbb{Z}^+$ , then either

- $S = \{0\}$
- $S = \mathbb{Z}a$ , where  $a$  is the smallest elements of  $S$ .

**Definition 1.2.5.** For two integers  $a, b \in \mathbb{Z}$  we say that  $a$  **divides**  $b$  if  $\frac{b}{a} \in \mathbb{Z}$  denoted  $a|b$ .

#### 1.2.1 Greatest Common Divisor

**Definition 1.2.6.** The **greatest common divisor** of two integers  $a, b \in \mathbb{Z}$  is the integer  $d \in \mathbb{Z}$  such that

$$\mathbb{Z}d = \mathbb{Z}a + \mathbb{Z}b = \{n \in \mathbb{Z} | n = ra + sb \forall r, s \in \mathbb{Z}\}$$

**Proposition 1.2.7. Properties of the greatest common divisor** Let  $a, b \in \mathbb{Z}$ , not both zero, and let  $d$  be the greatest common divisor. Then

1. There are integers  $r, s \in \mathbb{Z}$  such that  $d = ra + sb$ .
2.  $d|a$  and  $d|b$ .
3. If  $e \in \mathbb{Z}$  such that  $e|a$  and  $e|b$  then  $e|d$ .

**Definition 1.2.8.** Two integers  $a, b \in \mathbb{Z}$  are **relatively prime** if  $\gcd(a, b) = 1$ .

**Corollary 1.2.9.** A pair  $a, b \in \mathbb{Z}$  is relatively prime if and only if there are integers  $r, s \in \mathbb{Z}$  such that  $ra + sb = 1$ .

**Corollary 1.2.10.** Let  $p$  be a prime integer. If  $p$  divides a product  $ab$  of integers, then at least one of  $p|a$  or  $p|b$  holds.

### 1.2.2 Least Common Multiple

**Definition 1.2.11.** The **least common multiple** of two integers  $a, b \in \mathbb{Z}$  is the integer  $m \in \mathbb{Z}$  such that

$$\mathbb{Z}m = \mathbb{Z}a \cap \mathbb{Z}b$$

**Proposition 1.2.12. Properties of least common multiple** Let  $a, b$  be non-zero integers and let  $m$  be there least common multiple. Then

1.  $a|m$  and  $b|m$ .
2. If  $n \in \mathbb{Z}$  such that  $b|n$  and  $a|n$ , then  $m|n$ .

**Corollary 1.2.13.** For  $d = \gcd(a, b)$  and  $m = \text{lcm}(a, b)$  then  $ab = dm$ .

### 1.2.3 Cyclic Groups

**Definition 1.2.14.** Let  $G$  be a group and  $x \in G$ . The **cyclic subgroup** generated by  $x$  denoted  $\langle x \rangle$  is

$$\langle x \rangle = \{\dots, x^{-2}, x^{-1}, 1, x^1, x^2, \dots\}$$

*Remark.* For any subgroup  $S$  that contains  $x$  we have  $S \subset \langle x \rangle$ .

**Definition 1.2.15.** The **order of an element**  $x \in G$  is the order of the group  $\langle x \rangle$ . This is the smallest positive integer  $n$  such that  $x^n = 1$ .

**Proposition 1.2.16.** Let  $\langle x \rangle \subset G$  and consider the set  $S = \{k \in \mathbb{Z} | x^k = 1\}$

1. The set  $S$  is a subgroup of  $\mathbb{Z}^+$
2.  $x^r = x^s$  ( $r \geq s$ ) if and only if  $x^{r-s} = 1$ .
3. If  $S \neq \{0\}$ , then  $S = \mathbb{Z}n$  for some positive  $n \in \mathbb{Z}$  and  $\langle x \rangle = \{1, x^1, x^2, \dots, x^{n-1}\}$

**Proposition 1.2.17.** Let  $x$  be an element of finite order  $n$  in a group and let  $k \in \mathbb{Z}$ . Let  $k = nq + r$ , where  $q, r \in \mathbb{Z}$  and  $0 \leq r < n$ . Then

1.  $x^k = x^r$
2.  $x^k = 1$  if and only if  $r = 0$ .
3. The order of  $x^k$  is  $n/\gcd(k, n)$ .

## 1.3 Homomorphisms

**Definition 1.3.1.** A **homomorphism**  $\varphi : G \rightarrow G'$  is a map from a group  $G$  to a group  $G'$  such that for any  $a, b \in G$  we have

$$\varphi(ab) = \varphi(a)\varphi(b)$$

**Proposition 1.3.2.** Let  $\varphi : G \rightarrow G'$  be a homomorphism

1.  $\varphi(1) = 1$
2.  $\varphi(a^{-1}) = \varphi(a)^{-1}$  for any  $a \in G$

**Definition 1.3.3.** A homomorphism  $\varphi : G \rightarrow G'$  is **injective** if  $\varphi(x) = \varphi(u) \Rightarrow x = u$

**Definition 1.3.4.** A homomorphism  $\varphi : G \rightarrow G'$  is **surjective** if for every  $b \in G'$ , there exists  $a \in G$  such that  $\varphi(a) = b$ .

**Definition 1.3.5.** Let  $\varphi : G \rightarrow G'$  be a homomorphism

1. The **kernal** of  $\varphi$  denoted  $\ker(\varphi)$  is the set

$$\ker(\varphi) = \{a \in G | \varphi(a) = 1\}$$

2. The **image** of  $\varphi$  denoted  $\text{Im}(\varphi)$  is the set

$$\text{im}(\varphi) = \{b \in G' | \exists a \in G, \varphi(a) = b\}$$

**Corollary 1.3.6.** A homomorphism  $\varphi : G \rightarrow G'$  is injective if  $\ker(\varphi) = \{1\}$

**Corollary 1.3.7.** A homomorphism  $\varphi : G \rightarrow G'$  is surjective if  $\text{Im}(\varphi) = G'$

**Proposition 1.3.8.** Let  $\varphi : G \rightarrow G'$  be a homomorphism the  $\ker(\varphi)$  and  $\text{Im}(\varphi)$  are subgroups of  $G$  and  $G'$

**Definition 1.3.9.** An **isomorphism** is a **bijective** homomorphism. A homomorphism is **bijective** if it is both injective and surjective.

**Proposition 1.3.10.** If  $\varphi : G \rightarrow G'$  is an isomorphism, then  $\varphi^{-1} : G' \rightarrow G$  is also an isomorphism.

**Definition 1.3.11.** Two groups  $G$  and  $G'$  are **isomorphic** if there is an isomorphism  $\varphi : G \rightarrow G'$ .

**Definition 1.3.12.** An **automorphism** is an isomorphism  $\varphi : G \rightarrow G$ .

## 1.4 Cosets

**Definition 1.4.1.** Let  $H$  be a subgroup of  $G$ . The **left coset** of  $H$  induced by an element  $a \in G$  is the set

$$aH = \{ah | h \in H\}$$

The **right coset** of  $H$  induced by an element  $a \in G$  is the set

$$Ha = \{ha | h \in H\}$$

**Proposition 1.4.2.** Let  $H$  be a subgroup of  $G$ . The left cosets partition  $G$ . The right cosets partition  $G$ .

**Definition 1.4.3.** For a subgroup  $H$  of  $G$ . The **index of  $H$  in  $G$**  denoted  $[G : H]$  is the number of left cosets of  $H$  in  $G$ .

**Lemma 1.4.4.** All left cosets  $aH$  and all right cosets  $Ha$  of a subgroup  $H$  of a group  $G$  have the same order.

**Lemma 1.4.5. Counting Formula.** For a subgroup  $H$  of  $G$  we have

$$|G| = |H|[G : H]$$

**Theorem 1.4.6. Lagrange's Theorem.** Let  $H$  be a subgroup of a finite group  $G$ . The order of  $H$  divides the order of  $G$ .

**Corollary 1.4.7.** The order of an element of a finite group divides the order of the group.

**Corollary 1.4.8.** If  $G$  is a group of prime order then for  $a \in G$  where  $a \neq \mathbb{I}$ , we have  $G = \langle a \rangle$ .

**Corollary 1.4.9.** If  $\varphi : G \rightarrow G'$  is a homomorphism of finite groups then

$$|G| = |\ker(\varphi)||\text{Im}(\varphi)|$$

## 1.5 Normal Subgroups

**Definition 1.5.1.** A subgroup  $N$  of a group  $G$  is **normal** if for every  $a \in N$  and  $g \in G$ ,  $gag^{-1} \in N$ .

**Proposition 1.5.2.** For any homomorphism  $\varphi : G \rightarrow G'$  the  $\ker(\varphi)$  is a normal subgroup of  $G$ .

**Proposition 1.5.3.** Let  $H \subset G$  be a subgroup. Then the following are equivalent

1.  $H$  is a normal subgroup.
2. For all  $g \in G$ ,  $gHg^{-1} = H$
3. For all  $G \in G$ ,  $gH = Hg$
4. Every left coset of  $H$  in  $G$  is a right coset of  $H$  in  $G$ .

**Corollary 1.5.4.** If a group  $G$  has just one subgroup of order  $n$ , then that subgroup is normal.

## 1.6 Quotient Groups

**Definition 1.6.1.** If  $H \subset G$  is a subgroup. The **Quotient** is defined  $G/H = \{\text{left cosets of } H\}$ .

**Proposition 1.6.2.** If  $H \subset G$  is a normal subgroup, then  $G/H$  is a group with law of composition  $[aH][bH] = [abH]$ .

**Theorem 1.6.3. Correspondence Theorem** Let  $\varphi : G \rightarrow G'$  be a surjective homomorphism with kernel  $K$ . There is a bijective correspondence between subgroups of  $G'$  and subgroups of  $G$  that contain  $K$ .

$$\{\text{subgroups of } G \text{ that contain } K\} \leftrightarrow G/K$$

## 1.7 Product Groups

**Definition 1.7.1.** If  $G$  and  $G'$  are groups,  $G \times G'$  is the **product group** defined

$$G \times G' = \{(g, g') | g \in G, g' \in G'\}$$

with the law of composition

$$(a, a')(b, b') = (ab, a'b')$$

**Proposition 1.7.2.** Let  $G$  be a cyclic group of order  $mn$  where  $\gcd(m, n) = 1$  then  $G \cong C_m \times C_n$ .

**Proposition 1.7.3.** Let  $H, K$  be subgroups of a group  $G$ . Consider the multiplication map

$$f : H \times K \rightarrow G$$

given by  $f(h, k) = hk$ . Then

1.  $f$  is a homomorphism if and only if  $kh = hk$  for all  $h \in H$  and  $k \in K$
2.  $f$  is injective if and only if  $H \cap K = \{1\}$
3. if  $H$  is normal the image  $HK$  of  $f$  is a subgroup of  $G$ .

In particular,  $G \cong H \times K$  under  $f$  if and only if  $H \cap K = \{1\}$ ,  $HK = G$  and  $K$  and  $H$  are both normal.

**Proposition 1.7.4.** The map  $\pi : G \rightarrow G/N$  defined by  $\pi(x) = [aN]$  such that  $x \in aN$  is a surjective homomorphism with kernel  $N$ .

**Theorem 1.7.5. First Isomorphism Theorem** Let  $\varphi : G \rightarrow G'$  be a surjective homomorphism and let  $N$  be its kernel.

$$G' \cong G/N$$

## 1.8 Group Actions

**Definition 1.8.1.** An **action** of a group  $G$  on a set  $S$  is a map

$$\begin{aligned} G \times S &\rightarrow S \\ (g, s) &\mapsto g * s \end{aligned}$$

such that

1.  $1 * s = s$  for all  $s \in S$ .
2. **Associativity:**  $(gg') * s = g * (g' * s)$  for all  $g, g' \in G$  and  $s \in S$ .

**Definition 1.8.2.** Given an action of a group  $G$  on the set  $S$ , the **orbit**  $O_s$  of an element  $s \in S$  is

$$O_s = \{gs \in S \mid g \in G\}$$

**Definition 1.8.3.** An action of  $G$  on  $S$  is **transitive** if  $S = O_s$  for some  $s \in S$ .

**Definition 1.8.4.** The **stabilizer**  $G_s$  of an element  $s \in S$  is

$$G_s = \{g \in G \mid gs = s\}$$

**Proposition 1.8.5.** Let  $G$  be a subgroup of a group  $G$ .

1. The action of  $G$  on  $G/H$  is transitive.
2. The stabilizer  $G_{[H]}$  of  $[H]$  is the subgroup  $H$ .

**Theorem 1.8.6. Orbit Stabilizer Theorem** Let  $G$  be a group action on a set  $S$ . For any  $s \in S$ , there is a bijection

$$\begin{aligned} \epsilon : G/G_s &\leftrightarrow O_s \\ [aG_s] &\mapsto as \end{aligned}$$

such that  $\epsilon(g[C]) = g\epsilon([C])$  for all  $g \in G$  and  $[C] \in G/G_s$

**Corollary 1.8.7.** Let  $G$  be a group acting on a finite set  $S$ . Then for any  $s \in S$

$$|G| = |O_s| |G_s|$$

## 1.9 Conjugation

**Definition 1.9.1.** The **conjugate** of  $a \in G$  by  $g \in G$  is  $gag^{-1}$ .

**Definition 1.9.2.** The **conjugation action** is the action of a group  $G$  defined by  $G \times G \rightarrow G$  with  $(g, x) \mapsto gxg^{-1}$ .

**Lemma 1.9.3.**  $G$  is abelian  $\Leftrightarrow$  conjugation map is the identity

**Definition 1.9.4.** The **centralizer** of  $x$  is the stabilizer of  $x$  under conjugation.

$$Z(x) = \{g \in G | gxg^{-1} = x\} = \{g \in G | gx = xg\}$$

**Definition 1.9.5.** The conjugacy class of  $x$  is the orbit of  $x$  under conjugation.

$$C(x) = \{gxg^{-1} \in G | g \in G\}$$

**Definition 1.9.6.** The **center** of a group  $G$  is the subgroup

$$Z = \{z \in G | zg = gz \text{ for all } g \in G\}$$

**Corollary 1.9.7.** The center of a group is a normal subgroup.

**Corollary 1.9.8.** Every centralizer contains the center.

**Proposition 1.9.9. The Class Equation** The orbits of of conjugation partition the group.

$$|G| = \sum_{\text{conjugacy classes } C} |C|$$

## 1.10 p-Groups

**Definition 1.10.1.** A  $p$ -group is a group of order  $p^n$  for some prime  $p$ .

**Proposition 1.10.2.** The center of a  $p$ -group is non-trivial.

**Theorem 1.10.3. Fixed Point Theorem** Let  $G$  be a  $p$ -group action on a finite set  $S$ . If  $|S|$  is not divisible by  $p$ , then there is a fixed point for the action of  $G$  on  $S$ .

**Proposition 1.10.4.** Every group of order  $p^2$  is abelian.

**Corollary 1.10.5.** A group of order  $p^2$  is either cyclic or a product of two cyclic groups

**Definition 1.10.6.** A subgroup  $H \subset G$  of order  $p^e$  is called a **Sylow  $p$ -subgroup**.

**Theorem 1.10.7. First Sylow Theorem** A finite group whose order is divisible by a prime  $p$  contains a Sylow  $p$ -subgroup.

**Corollary 1.10.8.** A group whose order is divisible by a prime  $p$  contains a Sylow  $p$ -subgroup.

**Theorem 1.10.9. Second Sylow Theorem** Let  $G$  be a finite group whose order is divisible by a prime  $p$ .

1. The Sylow  $p$ -subgroups of  $G$  are conjugate subgroups.
2. Every subgroup of  $G$  that is a  $p$ -group is contained in a Sylow  $p$ -subgroup.

**Corollary 1.10.10.** A group  $G$  has just one Sylow  $p$ -subgroup  $H$  if and only if  $H$  is normal.

**Theorem 1.10.11. Third Sylow Theorem** Let  $G$  be a finite group whose order  $n = p^e m$ , with  $p$  prime and  $p$  not dividing  $m$ . Let  $s$  be the number of Sylow  $p$ -subgroups of  $G$ . Then  $s$  divides  $m$  and  $s \equiv 1 \pmod{p}$ .