

Probability Notes

May 27, 2020

Contents

1	Probability Theorems	1
1.1	Set Theorems	1
1.2	Basic Probability Rules	1
1.2.1	Total Probability Theorem	1
1.3	Independence	1
1.4	Joint Probability Distributions	1
1.5	Expected Value	2
1.5.1	Total Expectation Theorem	2
1.6	Variance	3
1.7	Cumulative Probability Distribution	3
2	Covariance and Correlation	3
3	Iterated Expectation and Variance	3
4	Random number of Random Variables	4
5	Convolutions	4
6	Moment Generating Function	5
6.1	Moment Generating Function for Sum of Independent RV	6
7	Binomial Random Variable	6
7.1	Mean and Variance	6
8	Continuous Uniform Random Variable	6
8.1	Mean and Variance	7
9	Gaussian Distribution	7
10	Counting Process	7
10.1	Independent Increments	7
10.2	Stationary Increments	8
11	Renewal Process	8
11.1	Laplace Transform	9
11.2	Calculating the Expectation	9
11.3	Limit Theorems for Renewal Processes	10

12 Bernoulli Process	11
12.1 Mean and Variance	11
12.2 Interarrival Times (Geometric Random Variable)	11
12.3 Sum of Interarrival times	11
13 Exponential Distribution	12
13.1 Mean and Variance	12
13.2 Memoryless Property	13
14 Poisson Process	13
14.1 Poisson Random Variable	13
14.2 Mean and Variance	13
14.3 Poisson Process	14
14.4 A Special Counting Process	14
14.5 Derivation from Bernoulli Process	14
14.6 Time till kth arrival	15
14.7 Time of 1st Arrival	15
14.8 Renewal Process	15
14.9 Merging of Poisson Processes	15
14.10 Splitting of Poisson Process	16
14.11 Random Incidence for Poisson	16
14.12 Non Homogenous Poisson Process	17
15 Gamma Distribution	17
15.1 Mean and Variance	17
15.2 Sum of Gamma Distributions	18
16 Chi-Square Distribution	18
16.1 Relation between Chi-Square and Gamma Distribution	19
16.2 Mean and Variance	20
17 t-Distribution	20
17.1 Mean and Variance	20
18 F-Distribution	21
19 Logistics Distribution	21
19.1 Mean	22
20 Markov Process	22
20.1 Recurring and Transient States	22
20.2 Steady State Probabilities	22
20.3 Birth Death Process	23
20.4 Absorption Probabilities	23
21 Central Limit Theorem	24
21.1 Weak Law of Large Numbers	24
21.2 Markov Inequality/Chebychev Inequality	24
21.3 Central Limit Theorem	25

22 Distribution of Sample Mean and Variance	26
22.1 Sample Mean	26
22.2 Sample Variance	27
22.3 Distributions for a Normal Population	27
23 Parameter Estimation	29
23.1 Maximum Likelihood Estimator	29
23.1.1 MLE for Bernoulli Variable	29
23.1.2 MLE for Poisson Variable	30
23.1.3 MLE for Normal Variable	30
23.1.4 MLE for Uniform Random Variable	31
23.2 Interval Estimates	31
23.2.1 Confidence interval for Mean of Normal Distribution when Variance is Known	32
23.2.2 Confidence interval for Mean of Normal Distribution when Variance is Unknown	33
23.2.3 Confidence interval for Variance of Normal Distribution when Mean is Unknown	33
23.2.4 Estimating Difference in Means of Two Normal Populations	34
23.2.5 Confidence Interval for Mean of Bernoulli Random Variable	36
24 Linear Regression	36
25 Exercises	37
25.1 Problems	37
25.2 Solutions	42

1 Probability Theorems

1.1 Set Theorems

For any three sets, the following hold true

$$\begin{aligned}A &= (A \cap B) \cup (A \cap B^c) \text{ where } B \text{ and } B^c \text{ are disjoint} \\A \cap (B \cup C) &= (A \cap B) \cup (A \cap C) \\A \cup (B \cap C) &= (A \cup B) \cap (A \cup C)\end{aligned}$$

1.2 Basic Probability Rules

$$\begin{aligned}\text{If } A \cap B &= \phi, \text{ then } P(A \cup B) = P(A) + P(B) \\P(A|B)P(B) &= P(B|A)P(A) = P(A \cap B) && \text{Bayes' Theorem} \\P(A) &= P(A \cap B) + P(A \cap B^c) = P(A|B)P(B) + P(A|B^c)P(B^c) \\P(A \cap B \cap C) &= P(A)P(B|A)P(C|B, A) && \text{Chain Rule}\end{aligned}$$

1.2.1 Total Probability Theorem

Let A_1, A_2, \dots, A_n be n disjoint events that completely cover the event space, and B be another event, then

$$\begin{aligned}P(B) &= P(B|A_1)P(A_1) + P(B|A_2)P(A_2) + \dots + P(B|A_n)P(A_n) \\ \text{or, } P(B) &= \sum_{i=1}^n P(B|A_i)P(A_i)\end{aligned}$$

1.3 Independence

Two events A and B are independent iff

$$P(A \cap B) = P(A)P(B)$$

Note that *independence* is not the same as *disjoint*

$$A \cap B = \phi \Rightarrow P(A \cap B) = 0 \text{ but } P(A) \neq P(B) \neq 0$$

Multiple events A_1, A_2, \dots, A_n are independent iff

$$P(A_i \cap A_j \cap \dots \cap A_k) = P(A_i)P(A_j) \dots P(A_k) \quad \forall i, j, \dots, k \mid i, j, \dots, k \in 1, 2, \dots, n$$

Conditional Independence is similar to the above equation. For an event C ,

$$P(A_i \cap A_j \cap \dots \cap A_k | C) = P(A_i | C)P(A_j | C) \dots P(A_k | C) \quad \forall i, j, \dots, k \mid i, j, \dots, k \in 1, 2, \dots, n$$

1.4 Joint Probability Distributions

Joint Probability Distributions are defined for two or more than two variables. In this section, we only consider two variables. The formal definition is

$$P_{XY}(x, y) = P(X = x \text{ and } Y = y)$$

Based on this definition, the following theorems follow

$$\sum_x \sum_y P_{XY}(x, y) = 1$$

$$P_X(x) = \sum_y P_{XY}(x, y) \quad \text{Marginal Probability}$$

$$P_{X|Y}(x|y) = P_{X|Y}(X = x|Y = y) = \frac{P_{XY}(x, y)}{P_Y y}$$

$$\sum_x P_{X|Y}(x|y) = 1 \quad \text{Since Y is fixed and we sum over all X's}$$

$$P_{XYZ}(x, y, z) = P_X(x)P_{Y|X}(y|x)P_{Z|X,Y}(z|x, y) \quad \text{Chain Rule}$$

1.5 Expected Value

Before going to expected value, let's define a Random Variable

Random Variable X is a linear map : $\mathbb{R} \rightarrow \mathbb{R}$. The value taken by the variable is denoted by x . X will have an associated probability distribution, i.e., $P_X(X = x)$. Using these quantities, we have

$$E[X] = \sum_x x P_X(X = x) \quad \text{Expected Value}$$

Based on this definition, the following theorems for expected value follow

$$E[\alpha] = \alpha E[X] \quad = \alpha E[X]$$

$$E[\alpha X + \beta] = \alpha E[X] + \beta$$

$$E[g(X)] = \sum_x g(x) P_X(X = x)$$

$$E[X^2] = \sum_x x^2 P_X(X = x) \quad \text{Also called Second Moment}$$

$$E[X|A] = \sum_x x P_{X|A}(X|A)$$

$$E[g(X)|A] = \sum_x g(x) P_{X|A}(X|A)$$

$$E[X + Y + Z] = E[X] + E[Y] + E[Z] \quad \text{Linearity of Expectation}$$

$$E[XY] = \sum_X \sum_Y xy P_{XY}(x, y)$$

$$E[g(X, Y)] = \sum_X \sum_Y g(xy) P_{XY}(x, y)$$

$$E[XY] = E[X]E[Y] \quad \text{if X and Y are independent}$$

where $\alpha, \beta \in \mathbb{R}$, $g(X) : \mathbb{R} \rightarrow \mathbb{R}$, and A is an event, X, Y, Z are Random Variables

1.5.1 Total Expectation Theorem

The *Total Expectation Theorem* is the natural extension of the *Total Probability Theorem*. Let A_1, A_2, \dots, A_n be n disjoint events that completely cover the event space, and X be random variable, then

$$E[X] = E[X|A_1]P(A_1) + E[X|A_2]P(A_2) + \dots + E[X|A_n]P(A_n)$$

$$\text{or, } E[X] = \sum_{i=1}^n E[X|A_i]P(A_i)$$

1.6 Variance

The formal definition of variance is

$$Var(X) = E[(X - \bar{X})^2] = E[X^2] - E[X]^2$$

Using this definition, the following theorems follow

$$E[X^2] = E[X]^2 + Var(X)$$

$$Var(\alpha) = 0$$

$$Var(\alpha X + \beta) = \alpha^2 Var(X)$$

$$Var(X + Y) = Var(X) + Var(Y) \text{ if } X \text{ and } Y \text{ are independent random variables}$$

1.7 Cumulative Probability Distribution

Cumulative probability distribution is defined for both discrete and continuous variables

$$F_x(X) = P(X \leq x) = \begin{cases} \int_{-\infty}^x p_X(t) dt & X \text{ is a discrete random variable} \\ \sum_{k \leq x} P_X(k) & X \text{ is a continuous random variable} \end{cases}$$

2 Covariance and Correlation

For any two random variables X and Y,

$$Cov(X, Y) = E[(X - \bar{X})(Y - \bar{Y})] = E[XY] - E[X]E[Y]$$

$$Cov(X, X) = Var(X)$$

$$\begin{aligned} Corr(X, Y) &= E\left[\left(\frac{X - \bar{X}}{\sigma_X}\right)\left(\frac{Y - \bar{Y}}{\sigma_Y}\right)\right] \\ &= \frac{Cov(X, Y)}{\sigma_X \sigma_Y} \end{aligned}$$

Key points to note

- Independence $\Rightarrow Cov(X, Y) = Corr(X, Y) = 0$, but the converse is **not** true
- Correlation is dimensionless and $-1 \leq Corr(X, Y) \leq 1$ with value close to 0 implying minimal relation and values close to $-1, 1$ implying perfect relation

3 Iterated Expectation and Variance

The law of iterated expectation tells the following about expectation and variance

$$E[E[X|Y]] = E[X]$$

$$Var(X) = E[Var(X|Y)] + Var(E[X|Y])$$

Proof for Iterated Expectation

$$\begin{aligned} P(X) &= \sum_y P(X|Y)P(Y) \\ \Rightarrow E[X] &= \sum_x xP(X) = \sum_x \sum_y xyP(X|Y)P(Y) \\ &= \sum_y P(Y) \sum_x xP(X|Y) = \sum_y P(Y)E[X|Y] \\ \text{or, } E[X] &= E[E[X|Y]] \quad E[X|Y] \text{ is a function of } Y \text{ and not } X \end{aligned}$$

Proof for Variance

$$\begin{aligned}
Var(X) &= E[X^2] - E[X]^2 \\
Var(X|Y) &= E[(X - \bar{X})^2|Y] = E[X^2|Y] - E[X|Y]^2 & 1 \\
Var[E(X|Y)] &= E[E(X|Y)^2] - E[E(X|Y)]^2 \\
&= E[E[(X|Y)]^2] - E[X]^2 & 2 \\
E[Var(X|Y)] &= E[E[X^2|Y]] - E[E[X|Y]^2] & \text{from 1} \\
&= E[X^2] - E[E[X|Y]^2] & 3 \\
E[Var(X|Y)] + Var(E[X|Y]) &= E[X^2] - E[X]^2 & \text{adding 2 and 3} \\
&= Var(X)
\end{aligned}$$

4 Random number of Random Variables

Let X_i be independent identically distributed Random Variables and let $Y = \sum_{i=1}^N X_i$ be the sum of N such random variables where N itself is a random variable. Then,

$$\begin{aligned}
Y &= X_1 + X_2 + \cdots + X_N \\
E[Y|N = n] &= \sum_{i=1}^n E[X_i] \\
&= NE[X] \\
E[Y] &= E[E[Y|N]] = E[NE[X]] \\
&= E[N]E[X] & \text{since } E[X] \text{ will be a number} \\
Var(Y) &= E[Var(Y|N)] + Var(E[Y|N]) \\
&= E[NVar(X)] + Var(NE[X]) \\
&= E[N]Var(X) + E[X]^2Var(N)
\end{aligned}$$

5 Convolutions

Convolution operations are defined for both CDF and PDF/PMFs. Let X and Y be random independent variables, then

$$\begin{aligned}
F_{X+Y}(x) &= F_X * F_Y = \int_{\mathbb{R}} F_X(x-y) dF_Y(y) \\
p_{X+Y}(x) &= p_X * p_Y = \int_{\mathbb{R}} p_X(x-y) p_Y(y) dy
\end{aligned}$$

We can extend the idea to n independent variables as

$$F_X^{n*} = F_X * \cdots * F_X \text{ } n \text{ times}$$

It has the following properties for positive random variable X_i s

1.

$$F_X^{n*}(x) \leq F_X^n(x)$$

This can be proven as

$$\begin{aligned}
P(X_1 + \dots + X_n \leq x) &\leq P(X_1 \leq x, \dots, X_n \leq x) \\
P(X_1 + \dots + X_n \leq x) &\leq \prod_{i=1}^n P(X_i \leq x) \text{ by independence} \\
\text{or, } F_X^{n*}(x) &\leq F_X^n(x)
\end{aligned}$$

2.

$$F_X^{n*}(x) \geq F_X^{n+1}(x)$$

which follows immediately from the fact that

$$P(X_1 + \dots + X_n \leq x) \geq P(X_1 \leq x, \dots, X_{n+1} \leq x)$$

since the volume of the regions denoting the sums will be lower in the higher dimensions. This can be quickly verified by considering $X_1 \leq 1$ and $X_1 + X_2 \leq 1$.

6 Moment Generating Function

Moment generating function is defined as the following for all values of t

$$\phi(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p_X(x) & \text{for discrete case} \\ \int_{-\infty}^{\infty} e^{tx} f_X(x) & \text{for continuous case} \end{cases}$$

This function is called the moment generating function because all the moments of the random variable X can be obtained by successively differentiating the function $\phi(t)$.

$$\begin{aligned}
\phi'(t) &= \frac{d}{dt} E[e^{tX}] \\
&= E\left[\frac{d}{dt} e^{tX}\right] \\
&= E[X e^{tX}] \\
\text{mean} &= E[X] \\
&= \phi'(0)
\end{aligned}$$

Continuing in a similar fashion,

$$\begin{aligned}
\phi''(t) &= \frac{d}{dt} E[X e^{tX}] \\
&= E\left[\frac{d}{dt} X e^{tX}\right] \\
&= E[X^2 e^{tX}] \\
\text{variance} &= \phi''(0) \\
&= E[X^2]
\end{aligned}$$

In general, for any $n > 0$, the n^{th} derivative will give the n^{th} moment

$$\phi^n(0) = E[X^n]$$

There exists a one to one correspondence between the moment generating function and the distribution function of a random variable, similar to Lagrangian multipliers.

6.1 Moment Generating Function for Sum of Independent RV

An important property is in the context of sum of two or more random variables. The **moment generating of sum of two independent random variables is simply the product of the moment generating functions of the two individual random variables**

$$\begin{aligned}\phi_{X+Y}(t) &= E[e^{t(X+Y)}] \\ &= E[e^{tX}e^{tY}] \\ &= E[e^{tX}]E[e^{tY}]\end{aligned}$$

$$\boxed{\phi_{X+Y}(t) = \phi_X(t)\phi_Y(t)} \text{ for independent random variables}$$

7 Binomial Random Variable

Binomial Random Variable X is defined as the number of successes in an experiment with n independent trials, where each trial can only have two outcomes, *success* or *failure*.

Let X_i denote the Random Variable corresponding to the individual trials, with probability of success p . Then we have the following

$$X_i = \begin{cases} 1 & \text{if success in trial } i \\ 0 & \text{otherwise} \end{cases} \quad \text{indicator variable}$$

$$X = X_1 + X_2 + \cdots + X_n = \sum_{i=1}^n X_i$$

$$P(X = k) = \sum_{k=0}^n \binom{n}{k} p^k (1-p)^{n-k}$$

7.1 Mean and Variance

First let's calculate the mean and variance for a single trial X_i

$$\begin{aligned}E[X_i] &= 1 * p + 0 * (1-p) = p \\ \text{Var}(X_i) &= (1-p)^2 p + (0-p)^2 (1-p) = p(1-p)\end{aligned}$$

We know that all X'_i s are independent. Hence, the mean and variance for X become

$$\begin{aligned}E[X] &= E\left[\sum_{i=1}^n X_i\right] = \sum_{i=1}^n E[X_i] = np \\ \text{Var}(X) &= \text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i) = np(1-p)\end{aligned}$$

8 Continuous Uniform Random Variable

A uniform random variable is defined as follows

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

8.1 Mean and Variance

$$\begin{aligned} E[X] &= \int_a^b x \frac{1}{b-a} dx = \left[\frac{x^2}{2(b-a)} \right]_a^b \\ &= \frac{a+b}{2} \\ \text{Var}(X) &= \int_a^b \left(x - \frac{a+b}{2}\right)^2 \frac{1}{b-a} dx \\ &= \frac{(b-a)^2}{12} \end{aligned}$$

9 Gaussian Distribution

The gaussian distribution (or normal distribution) is defined between $-\infty$ and ∞ . It is parametrized by mean μ and variance σ , $X \sim \mathcal{N}(\mu, \sigma^2)$

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

As already described,

$$\begin{aligned} E[X] &= \mu \\ \text{Var}(X) &= \sigma^2 \end{aligned}$$

A *Standard Normal* is defined as a normal distribution with $\mu = 0$ and $\sigma^2 = 1$. Any normal distribution can be converted to a standard normal as $X = \frac{X-\mu}{\sigma}$. If $Y = aX + b$, then $Y \sim \mathcal{N}(a\mu + b, a^2\sigma^2)$

10 Counting Process

Counting process is used in scenarios when we want to count the occurrence of a certain event. N_t denotes the number of events till time t starting from 0. It is assumed that $N_0 = 0$. Formal definition is

A random process $\{N_t, t \in [0, \infty)\}$ is said to be a counting process if N_t is the number of events from time $t = 0$ upto time t . For a counting process, we assume

1. $N_0 = 0$
2. $N_t \in \{0, 1, 2, \dots\}$ for all $t \in [0, \infty)$
3. for $0 \leq s < t$, $N_t - N_s$ shows the number of events that occur in the interval $(s, t]$

10.1 Independent Increments

We say that a continuous time counting process N_t has independent increments if for all $0 \leq t_1 < t_2 < \dots < t_n$, the random variables

$$N_{t_2} - N_{t_1}, N_{t_3} - N_{t_2}, \dots, N_{t_n} - N_{t_{n-1}}$$

are independent.

Note that these differences are nothing but the number of arrivals in a given time interval. Thus, we are equivalently saying that **the number of arrivals in any two disjoint intervals are**

independent.

A very simple consequence of this property is:

Suppose we wish to find the probability of 2 arrivals in the interval $(1, 2]$ and 3 arrivals in the interval $(3, 5]$. Then,

$$P(2 \text{ arrivals in } (1, 2] \text{ and } 3 \text{ arrivals in } (3, 5]) = P(2 \text{ arrivals in } (1, 2])P(3 \text{ arrivals in } (3, 5])$$

since the arrivals in disjoint intervals are independent.

10.2 Stationary Increments

We say that a continuous time counting process N_t has stationary increments if for all $t_2 > t_1 \geq 0$ and for all $r > 0$, $N_{t_2} - N_{t_1}$ and $N_{t_2+r} - N_{t_1+r}$ are independent.

In other words, **the number of arrivals in a given time interval is invariant to its location.** Note that the number of arrivals in the time interval between t_1 and t_2 is nothing but $N_{t_2} - N_{t_1}$. By the above statement, if the process has stationary increments, then this quantity is same as $N_{t_2-t_1}$, which is the distribution of the counting process itself.

11 Renewal Process

This is a fundamental stochastic process useful in modelling arrivals and interarrival times. Some definitions will make the usage clear.

Let S_i denote the i th renewal time or the time when the i th arrival takes place. By definition, $S_0 = 0$. We can also define

$$\begin{aligned} S_n &= S_{n-1} + \xi_n \\ S_n &= \xi_1 + \xi_2 + \dots + \xi_n \end{aligned}$$

where ξ_i are positive ($P(\xi > 0) = 1$) independent identically distributed variables representing the interarrival times. We also define

$$\begin{aligned} N_t &= \arg\max_k \{S_k \leq t\} \\ \{S_n > t\} &= \{N_t < n\} \end{aligned}$$

or, N_t is simply the number of arrivals till the time t .

Define the following quantity

$$\begin{aligned} F^{n*} &= F_\xi * \dots * F_\xi \text{ } n \text{ times} \\ u(t) &= \sum_{i=1}^{\infty} F^{i*}(t) \end{aligned}$$

It can be shown that the function $u(t)$ converges. The expectation of N_t then becomes

$$\begin{aligned}
E[N_t] &= E[\text{number of } n \text{ such that } S_n \leq t] \\
&= E\left[\sum_{n=1}^{\infty} I(S_n \leq t)\right] && \text{sum of Indicators will equal } n \\
&= \sum_{n=1}^{\infty} P(S_n \leq t) && \text{since } E[\text{Indicator}] \text{ is just the function inside indicator} \\
&= \sum_{n=1}^{\infty} F^{n*}(t) && \text{by defining cumulative as sum of } \xi\text{s} \\
&= u(t)
\end{aligned}$$

11.1 Laplace Transform

For a density function f defined from $\mathbb{R}^{\geq 0} \rightarrow \mathbb{R}$, Laplace transform is

$$L_f(s) = \int_{\mathbb{R}^{\geq 0}} e^{-sx} f(x) dx$$

The following properties hold for this transform

1. If f is a probability density function, then

$$E[e^{-sx}] = L_f(s)$$

2. if f_1 and f_2 are two probability density functions, then

$$L_{f_1 * f_2}(s) = L_{f_1}(s) L_{f_2}(s)$$

3. If F is the cumulative probability distribution for X and p is the probability density function, then

$$L_{F_X}(s) = \frac{L_{p_X}(s)}{s}$$

which can be proven using integration by parts as follows

$$L_{F_X}(s) = \int_{\mathbb{R}^{\geq 0}} F_X(x) \frac{d(e(-sx))}{s} = 0 + \frac{1}{s} \int_{\mathbb{R}^{\geq 0}} p_X(x) e^{-sx} dx$$

11.2 Calculating the Expectation

Armed with the concept of a Laplace transform, we make the following observation first

$$\begin{aligned}
u(t) &= \sum_{i=1}^{\infty} F^{i*}(t) = F(t) + \sum_{i=2}^{\infty} F^{i*}(t) \\
&= F(t) + \left(\sum_{i=1}^{\infty} F^{i*}(t) \right) * F(t) \\
&= F(t) + u(t) * F(t) \\
u(t) &= F(t) + u(t) * p(t)
\end{aligned}$$

where p is the probability density function and the last line stems from the fact that $\int u * F = \int u(x-y) dF(y) = \int u(x-y) p(y) dy$. Taking Laplace transform on both sides,

$$\begin{aligned}
L_u(s) &= L_F(s) + L_u(s)L_p(s) \\
L_u(s) &= \frac{L_p(s)}{s} + L_u(s)L_p(s) \text{ from 3} \\
L_u(s) &= \frac{L_p(s)}{s(1 - L_p(s))}
\end{aligned}$$

The last equation can be used to calculate the laplace transform of $u(t)$ and consecutively guess the functional form of $u(t)$.

11.3 Limit Theorems for Renewal Processes

The following two theorems hold true for Renewal processes

1. If $E[\xi] = \mu < \infty$, then

$$\lim_{t \rightarrow \infty} \frac{N_t}{t} = \frac{1}{\mu}$$

which is analogous to the strong law of large numbers. This can be proven as follows

$$\begin{aligned}
S_{N_t} \leq t \leq S_{N_t+1} & \text{ from the definition of } N_t \\
\text{or, } \frac{N_t}{S_{N_t+1}} \leq \frac{N_t}{t} \leq \frac{N_t}{S_{N_t}}
\end{aligned}$$

we can calculate the limits on the two bounds as

$$\lim_{t \rightarrow \infty} \frac{N_t}{S_{N_t}} = \lim_{n \rightarrow \infty} \frac{n}{S_n} = \frac{1}{\mu}$$

from the strong law of large numbers applied to $\lim_{n \rightarrow \infty} \frac{S_n}{n}$. Similarly, one can show

$$\lim_{t \rightarrow \infty} \frac{N_t}{S_{N_t+1}} = \lim_{t \rightarrow \infty} \frac{N_t}{N_t + 1} \lim_{t \rightarrow \infty} \frac{N_t + 1}{S_{N_t+1}} = 1 * \frac{1}{\mu}$$

2. If $Var(\xi) = \sigma^2 < \infty$, then

$$\lim_{t \rightarrow \infty} \frac{N_t - t/\mu}{\sigma\sqrt{t}/\mu^{3/2}} = \mathcal{N}(0, 1)$$

which is analogous to the central limit theorem. It can be proven by considering the CLT on ξ s

$$\begin{aligned}
\lim_{n \rightarrow \infty} P\left(\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq x\right) &= \text{CDF of } \mathcal{N}(0, 1) \\
\text{or, } \lim_{n \rightarrow \infty} P(S_n \leq n\mu + \sigma\sqrt{n}x) &= \text{CDF of } \mathcal{N}(0, 1) \\
\text{or, } \lim_{n \rightarrow \infty} P(N_t \geq n) &= \text{CDF of } \mathcal{N}(0, 1) \text{ from definition of } N_t, \text{ where } t = n\mu + \sigma\sqrt{n}x
\end{aligned}$$

We substitute $n\mu = t$ for very large value of n , since the total time will become total variables into the expected time for one ξ when n is large. Hence,

$$\begin{aligned}
n &= \frac{t}{\mu} - \frac{\sigma\sqrt{t}}{\mu^{3/2}}x \\
\lim_{n \rightarrow \infty} P(N_t \geq n) &= \lim_{n \rightarrow \infty} P\left(\frac{N_t - t/\mu}{\sigma\sqrt{t}/\mu^{3/2}} \leq x\right) = \text{CDF of } \mathcal{N}(0, 1)
\end{aligned}$$

12 Bernoulli Process

Bernoulli process falls under the family of random processes, which are random variables continuously evolving over time. Bernoulli process can be described as a sequence of independent Bernoulli trials, where each trial has only two outcomes : success with $P(\text{success}) = p$ and failure.

$$P_{X_t}(x_t) = \begin{cases} p & \text{if } X_t = 1 \\ 1 - p & \text{if } X_t = 0 \end{cases}$$

$$E[X_t] = p$$

$$Var(X_t) = p(1 - p)$$

12.1 Mean and Variance

Number of successes S in n time slots

$$P(S = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

$$E[S] = np$$

$$Var(S) = np(1 - p)$$

12.2 Interarrival Times (Geometric Random Variable)

Let T_1 denote the number of trials till the first success

$$P(T_1 = t) = (1 - p)^{t-1} p \quad t \in 1, 2, \dots$$

$$E[T_1] = \frac{1}{p}$$

$$Var(T_1) = \frac{1 - p}{p^2}$$

This process is memoryless as all future coin flips are independent of whatever has happened till now. Also, the distribution is a **Geometric Random Variable**.

12.3 Sum of Interarrival times

We are interested in the total time till k arrivals. Let this random variable be Y_k

$$Y_k = T_1 + T_2 + \dots + T_k \quad \text{where } T_i \text{'s are i.i.d geometric with parameter } p$$

$$P(Y_k = t) = P(k - 1 \text{ arrivals between } t = 1 \text{ to } t = t \text{ and last arrival at time } t)$$

$$= \binom{t-1}{k-1} p^k (1 - p)^{t-k} \quad \forall t \geq k$$

$$E[Y_k] = \sum_{i=1}^k k E[T_i]$$

$$= \frac{k}{p}$$

$$Var(Y_k) = \sum_{i=1}^k Var(T_i)$$

$$= \frac{k(1 - p)}{p^2}$$

13 Exponential Distribution

Exponential distribution is characterized by the parameter λ and has the following probability distribution

$$f_X(x) = \begin{cases} 0 & \text{if } x < 0 \\ \lambda e^{-\lambda x} & \text{otherwise} \end{cases}$$

Exponential distribution is used to represent the interarrival time probability distribution in the context of Poisson Process. The cumulative distribution is given by

$$\begin{aligned} F_X(x) &= \begin{cases} 0 & \text{if } x < 0 \\ 1 - e^{-\lambda x} & \text{otherwise} \end{cases} \\ P(X > x) &= \int_x^{\infty} \lambda e^{-\lambda x} dx \\ &= e^{-\lambda x} \end{aligned}$$

13.1 Mean and Variance

The mean of the distribution is given by

$$\begin{aligned} E[x] &= \int_0^{\infty} \lambda x e^{-\lambda x} dx \\ &= [-x e^{-\lambda x}]_0^{\infty} + \int_0^{\infty} e^{-\lambda x} dx = \frac{1}{\lambda} \end{aligned}$$

$$\boxed{E[X] = \frac{1}{\lambda}}$$

where we used integration by parts, $\int uv' = uv - \int u'v$ and substituted $u = x$ and $v = -e^{-\lambda x}/\lambda$.

For variance, we first calculate the value of $E[x^2]$

$$\begin{aligned} E[x^2] &= \int_0^{\infty} \lambda x^2 e^{-\lambda x} dx \\ &= [-x^2 e^{-\lambda x}]_0^{\infty} + \int_0^{\infty} 2x e^{-\lambda x} dx \\ &= \left[\frac{-2x e^{-\lambda x}}{\lambda} \right]_0^{\infty} - \left[\frac{2e^{-\lambda x}}{\lambda^2} \right]_0^{\infty} \\ &= \frac{2}{\lambda^2} \\ \text{Var}(X) &= E[X^2] - E[X]^2 \end{aligned}$$

$$\boxed{\text{Var}(X) = \frac{1}{\lambda^2}}$$

The above property can be generalized for the n th power as well

$$E[X^n] = \frac{n!}{\lambda^n}$$

13.2 Memoryless Property

A fundamental mathematical property of the exponential distribution is the memoryless property. In summary, this means that whatever has transpired till now will not affect the future distribution. Mathematically $P(T > t + s | T > t)$ is independent of t

$$\begin{aligned} P(T > t + s | T > t) &= \frac{P(T > t + s \text{ and } T > t)}{P(T > t)} \\ &= \frac{P(T > t + s)}{P(T > t)} \\ &= \frac{e^{-\lambda(t+s)}}{e^{-\lambda t}} \\ &= e^{-\lambda s} \end{aligned}$$

$$\boxed{P(T > t + s | T > t) = P(T > s)}$$

14 Poisson Process

14.1 Poisson Random Variable

A random variable X is said to be $Poisson(\mu)$ if it has the following probability distribution

$$p_X(x = k) = \begin{cases} e^{-\mu} \frac{\mu^k}{k!} & \text{for all } x = \{0, 1, 2, \dots\} \\ 0 & \text{otherwise} \end{cases}$$

The sum of n independent Poisson variables is also Poisson

$$X_1 + X_2 + \dots + X_n \sim Poisson(\mu_1 + \mu_2 + \dots + \mu_n)$$

14.2 Mean and Variance

Expected value is calculated as follows

$$\begin{aligned} E[X] &= \sum_{k=0}^{\infty} k e^{-\mu} \frac{\mu^k}{k!} = \mu e^{-\mu} \sum_{k=1}^{\infty} \frac{\mu^{k-1}}{(k-1)!} \\ &= \mu e^{-\mu} \sum_{k=0}^{\infty} \frac{\mu^k}{k!} \end{aligned}$$

$$\boxed{E[X] = \mu}$$

Variance can be calculated using $Var(X) = E[X^2] - E[X]^2$

$$\begin{aligned} E[X^2] &= \sum_{k=0}^{\infty} k^2 e^{-\mu} \frac{\mu^k}{k!} = \mu e^{-\mu} \sum_{k=1}^{\infty} k \frac{\mu^{k-1}}{(k-1)!} \\ &= \mu e^{-\mu} \sum_{k=0}^{\infty} (k+1) \frac{\mu^k}{k!} \\ &= \mu e^{-\mu} \left(\mu \sum_{k=1}^{\infty} \frac{\mu^{k-1}}{(k-1)!} + \sum_{k=0}^{\infty} \frac{\mu^k}{k!} \right) \\ &= \mu e^{-\mu} (\mu e^{\mu} + e^{\mu}) \end{aligned}$$

$$Var(X) = E[X^2] - E[X]^2$$

$$\boxed{Var(X) = \mu}$$

Thus, mean and variance is the same for a Poisson variable.

14.3 Poisson Process

Poisson process also falls in the realm of random processes but is different from Bernoulli process as it is a continuous time process. This process is very commonly used to model arrival times and number of arrivals in a given time interval.

$$P(k, \tau) = \text{Probability of } k \text{ arrivals in interval of duration } \tau$$

$$\sum_k P(k, \tau) = 1 \quad \text{for a given } \tau$$

Assumptions

- The Probability is dependent only on τ and not the *location* of the interval
- Number of arrivals in disjoint time intervals are *independent*

14.4 A Special Counting Process

A counting process $N_t : t \in [0, \infty)$ is a Poisson process with rate λ if

1. $N_0 = 0$
2. N_t is composed of independent and stationary increments
3. The number of arrivals in any time interval $\tau > 0$ has $Poisson(\lambda\tau)$ distribution

Hence, for a Poisson process, the number of arrivals in any interval is dependent only on the length of that interval and not the location. Further, the number of arrivals in the interval will follow a Poisson distribution.

14.5 Derivation from Bernoulli Process

For a very small interval δ ,

$$P(k, \delta) = \begin{cases} 1 - \lambda\delta & k = 0 \\ \lambda\delta & k = 1 + O(\delta^2) \\ 0 & k > 2 \end{cases}$$

$$\lambda = \lim_{\delta \rightarrow 0} \frac{P(1, \delta)}{\delta} \quad \text{arrival rate per unit time}$$

$$E[k] = (\lambda\delta) * 1 + (1 - \lambda\delta) * 0$$

$$= \lambda\delta$$

$$\tau = n\delta$$

The last equation clearly implies that we can approximate the whole process as a Bernoulli process where we have n miniscule time intervals with at most one arrival per interval.

$$P(k \text{ arrivals}) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$= \binom{n}{k} \left(\frac{\lambda\delta}{n}\right)^k \left(1 - \frac{\lambda\delta}{n}\right)^{n-k}$$

$$\lambda\tau = np \quad \text{or, arrival rate * time} = E[\text{arrivals}]$$

$$Poisson = \lim_{\delta \rightarrow 0, n \rightarrow \infty} Bernoulli$$

$$\text{or, } P(k, \tau) = \frac{(\lambda\tau)^k e^{-\lambda\tau}}{k!} \quad k = 0, 1, \dots, \text{ for a given } \tau$$

$$\text{where, } \sum_k P(k, \tau) = 1 \quad \text{for a given } \tau$$

Let N_t denote the no of arrivals till time t , then

$$E[N_t] = \lambda t$$

$$Var(N_t) = \lambda t$$

14.6 Time till k th arrival

Suppose the k^{th} arrival happens at a time t . Then we are saying that there have been $k - 1$ arrivals till time t and the k^{th} arrival happens at time t (precisely in an interval of $[t, t + \delta]$). Let Y_k be the required time,

$$\begin{aligned} f_{Y_k}(t)\delta &= P(t \leq Y_k \leq t + \delta) \\ &= P(k - 1 \text{ arrivals in } [0, t])(\lambda\delta) \\ &= \frac{(\lambda t)^{k-1}}{(k-1)!} e^{-\lambda t} (\lambda\delta) \\ f_{Y_k}(t) &= \frac{\lambda^k t^{k-1}}{(k-1)!} e^{-\lambda t} \end{aligned} \quad \text{Erlang Distribution}$$

14.7 Time of 1st Arrival

Using the Erlang Distribution described above, we have

$$f_{Y_1}(t) = \lambda e^{-\lambda t}$$

$Y_k = T_1 + T_2 + \dots + T_k$ where all T_i are independent and exponential distributions.

14.8 Renewal Process

Poisson process can be seen as a special case of a renewal process, when the interarrival times are all exponentially distributed.

$$\begin{aligned} \text{Interarrival time } \xi_i &= \lambda e^{-\lambda t} \\ \text{Number of arrivals } P(N_t = n) &= \frac{(\lambda t)^n}{n!} e^{-\lambda t} \\ \text{Time till } n\text{th arrival } P(S_n = t) &= \lambda \frac{(\lambda t)^{n-1}}{(n-1)!} e^{-\lambda t} \text{ for } t > 0 \\ \text{Cumulative distribution } P(S_n \leq t) &= \begin{cases} 1 - e^{-\lambda t} \sum_{k=1}^{n-1} \frac{(\lambda t)^k}{k!} & \text{if } t > 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

14.9 Merging of Poisson Processes

Merging of two Poisson processes is also a Poisson process. Consider two flasbulbs of Red and Green colours, flashing as Possion processes with rates λ_1 and λ_2 . Then the process denoting the combined flashing of the two bulbs is also Poisson.

Consider a very small interval of time δ . In this small interval, any of the individual bulbs can have at most one flashes (since we ignore higher order terms). Thus, the following four possibilities arise

Thus, the combined process is Poisson with parameter $\lambda_1 + \lambda_2$

$$P(\text{arrival happened from first process}) = \frac{\lambda_1 \delta}{\lambda_1 \delta + \lambda_2 \delta} = \frac{\lambda_1}{\lambda_1 + \lambda_2}$$

0	<i>Red</i>	\overline{Red}
<i>Green</i>	$\lambda_1 \delta \lambda_2 \delta$	$(1 - \lambda_1 \delta) \lambda_2 \delta$
\overline{Green}	$\lambda_1 \delta (1 - \lambda_2 \delta)$	$(1 - \lambda_1 \delta) (1 - \lambda_2 \delta)$

Table 1: Base Probabilities for flashes

0	<i>Red</i>	\overline{Red}
<i>Green</i>	0	$\lambda_2 \delta$
\overline{Green}	$\lambda_1 \delta$	$(1 - (\lambda_1 + \lambda_2) \delta)$

Table 2: Probabilities after ignoring δ^2 terms

14.10 Splitting of Poisson Process

Suppose we have a Poisson process with parameter λ which we split into two processes up and down, with probabilities p and $1 - p$. The two resulting processes are also Poisson with different parameters.

Consider a small time slot of length δ . Then,

$$\begin{aligned} P(\text{arrival in this time slot}) &= \lambda \delta \\ P(\text{arrival in up slot}) &= \lambda \delta p \\ P(\text{arrival in down slot}) &= \lambda \delta (1 - p) \end{aligned}$$

Thus, up and down are themselves Poisson with parameters λp and $\lambda(1 - p)$ respectively.

14.11 Random Indcidence for Poisson

Suppose we have a Poisson process with parameter λ running forever. We show up at a random time instant. What is the length of the chosen interarrival time (the total of the time from the last arrival to the next arrival).

Let T'_1 denote the time that has elapsed since the last arrival and T_1 be the time till the next arrival. Note that the reverse process is also Poisson with the same parameter. Thus,

$$E[\text{interarrival time}] = E[T'_1 + T_1] = \frac{1}{\lambda} + \frac{1}{\lambda} = \frac{2}{\lambda}$$

This may seem paradoxical since the time difference between any two arrivals in a Poisson process is same and it's expected length is $\frac{1}{\lambda}$, whereas we got an interval twice this length. The paradox is resolved by considering the fact that when we choose a random point in time, it is more likely to fall in an interval of larger size than the smaller ones (since probability will be proportional to the length of the interval).

Consider a separate example where we want to compare two values $E[\text{size of a family}]$ and $E[\text{size of a family of a given person}]$.

The two value will be different due to the underlying nature of the way experiment is conducted. For the first, we randomly choose families and average their sizes. Here, family of any size is equally likely to be picked. In the second case, we first pick a person from the population, get their family size, and then average the sizes of the families. Note that, this experiment is biased since the we are more likely to select people from larger families (or equivalently, it is more likely that we pick a person from a large family since the probability of picking is proportional to the family size). Hence, the second value will likely be larger and the two quantities are not equal.

14.12 Non Homogenous Poisson Process

Sometimes, it may not be accurate to use a simple Poisson process to model arrival. For example, a restaurant will not have the same rate of influx throughout the day. This rate itself is a function of time. In such cases, we model the arrivals as Non Homogenous Poisson Process.

For such a process, we have $\lambda(t) : [0, \infty) \rightarrow [0, \infty)$ and the counting process N_t is non homogenous if the following hold

1. $N_0 = 0$
2. The increments to N_t are **independent but not stationary**
3. For any small time interval δ , the probability of more than 1 arrival in the interval is zero

The distribution of arrivals in a time interval is still Poisson, but the Poisson parameter is now dependent on the location of the interval itself (since the process does not have stationary increments)

$$N_{t+s} - N_t \sim \text{Poisson}\left(\int_t^{t+s} \lambda(\alpha) d\alpha\right)$$

15 Gamma Distribution

A random variable is said to have a Gamma distribution if for parameters $\lambda > 0, \alpha > 0$ it has the following probability distribution

$$p_X(x) = \begin{cases} \frac{\lambda e^{-\lambda x} (\lambda x)^{\alpha-1}}{\Gamma(\alpha)} & \text{if } x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

The denominator in the above formula acts as nothing but a normalization constant and is defined as

$$\begin{aligned} \Gamma(\alpha) &= \int_0^{\infty} \lambda e^{-\lambda x} (\lambda x)^{\alpha-1} dx \\ &= \int_0^{\infty} e^{-y} y^{\alpha-1} dy && \text{by letting } \lambda x = y \\ &= (\alpha - 1) \int_0^{\infty} e^{-y} y^{\alpha-2} dy && \text{using integration by parts} \\ &= (\alpha - 1) \Gamma(\alpha - 1) \end{aligned}$$

Note that at $\alpha = 1$, $\Gamma(1) = \int_0^{\infty} \lambda e^{-\lambda x} = 1$. Hence, if α is an integer, $\Gamma(\alpha) = \alpha!$ using the recursion relation derived above.

For a fixed λ , as the value of α becomes large, the distribution takes the form of a normal distribution.

15.1 Mean and Variance

Mean and variance are easily obtainable for this using the moment generating function. Recall

$$\begin{aligned} \phi(t) &= E[e^{tX}] \\ \phi^n(t) &= E[X^n] \end{aligned}$$




Figure 1: Gamma distribution for $\lambda = 1$ and different values of α . The bottom figure shows the distribution for $\alpha = 50$.

For the current distribution,

$$\begin{aligned}\phi(t) &= \frac{\lambda^\alpha}{\Gamma(\alpha)} \int_0^\infty e^{tx} e^{-\lambda x} x^{\alpha-1} dx \\ &= \left(\frac{\lambda}{\lambda - t}\right)^\alpha\end{aligned}$$

Differentiating,

$$\begin{aligned}\phi'(t) &= \frac{\alpha \lambda^\alpha}{(\lambda - t)^{\alpha+1}} \\ \phi''(t) &= \frac{\alpha(\alpha + 1) \lambda^\alpha}{(\lambda - t)^{\alpha+2}} E[X] = \phi'(0) \\ \boxed{E[X] = \frac{\alpha}{\lambda}} \\ \text{Var}(X) &= \phi''(0) \\ \boxed{\text{Var}(X) = \frac{\alpha}{\lambda^2}}\end{aligned}$$

15.2 Sum of Gamma Distributions

Let X_1, X_2, \dots, X_n be n random variables that are gamma distributed with parameters $(\alpha_1, \lambda), (\alpha_2, \lambda), \dots, (\alpha_n, \lambda)$. Then the distribution of the sum of these random variables is itself a gamma distribution with the parameters $\alpha' = \sum_{i=1}^n \alpha_i$ and $\lambda' = \lambda$

16 Chi-Square Distribution

If Z_1, Z_2, \dots, Z_n are n independent standard normal variables, then the random variable X

$$\begin{aligned}X &= Z_1^2 + Z_2^2 + \dots + Z_n^2 \\ \text{then, } X &\sim \chi_n^2\end{aligned}$$

i.e., X follows the chi-square distribution with n degrees of freedom.

If we add two chi-square distributed variables with degrees of freedom n_1 and n_2 , then the resultant variable itself is chi-square distributed with $n_1 + n_2$ degrees of freedom. This simply follows from the fact that the sum of the two random variables is nothing but sum of $n_1 + n_2$

standard normal squared variables which is nothing but a chi-square variable with $n_1 + n_2$ degrees of freedom.

If $X \sim \chi_n^2$, then $\chi_{\alpha,n}^2$ is

$$P(X \geq \chi_{\alpha,n}^2) = \alpha$$

This quantity is usually listed in mathematical tables since they are heavily used in hypothesis testing.

16.1 Relation between Chi-Square and Gamma Distribution

Consider the moment generating function for a chi-square random variable with $n = 1$ degrees of freedom

$$\begin{aligned} E[e^{tX}] &= E[e^{tZ^2}] \quad Z \sim \mathcal{N}(0,1) \\ &= \int_{-\infty}^{\infty} e^{tx^2} f_Z(x) dx \quad \text{since } E[g(x)] = \sum_x g(x)p(x) \\ &= \frac{1}{\sqrt{2\pi}} \sum_{-\infty}^{\infty} e^{tx^2} e^{-x^2/2} \\ &= \frac{1}{\sqrt{2\pi}} \sum_{-\infty}^{\infty} e^{-x^2(1/2-t)} \\ \text{Using } \int_{-\infty}^{\infty} e^{-a(x+b)^2} &= \sqrt{\frac{\pi}{a}} \\ E[e^{tX}] &= \frac{1}{\sqrt{2\pi}} \sqrt{\frac{\pi}{1/2-t}} \\ &= \frac{1}{\sqrt{1-2t}} \end{aligned}$$

Extending this idea to the case of n degrees of freedom,

$$\begin{aligned} E[e^{tX}] &= E[e^{t(Z_1^2 + Z_2^2 + \dots + Z_n^2)}] \\ &= E\left[\prod_{i=1}^n e^{tZ_i^2}\right] \\ &= \prod_{i=1}^n E[e^{tZ_i^2}] \quad \text{since } Z_i \text{ are independent} \\ &= (1-2t)^{-n/2} \quad \text{from the derivation above} \end{aligned}$$

But, the quantity just derived is nothing but the moment generating function of the Gamma distribution with parameters $(n/2, 1/2)$. Hence, by the uniqueness of the moment generating function, we are forced to conclude that the **probability density function of a chi-square variable with n degrees is same as that of a Gamma distribution with parameters $(n/2, 1/2)$.**

Thus,

$$f_X(x) = \frac{\frac{1}{2} e^{-x/2} \left(\frac{x}{2}\right)^{(n/2)-1}}{\Gamma\left(\frac{n}{2}\right)} \quad x > 0$$

16.2 Mean and Variance

Since the distribution of a chi-square variable is identical to a Gamma distribution,

$$E[X] = n$$

$$Var(x) = 2n$$

17 t-Distribution

Let Z be a standard normal random variable and let χ_n^2 be a chi-square random variable. Assuming these two random variables are independent, the random variable T_n is

$$T_n = \frac{Z}{\sqrt{\chi_n^2/n}}$$

is said to have a t-distribution with n degrees of freedom.

This distribution is symmetric around the normal, and as n increases, the distribution becomes more and more like the standard normal distribution.




Figure 2: t-distribution for different degrees of freedom, and comparison with standard normal

From figure 2, we see that t-distribution is heavier tailed than a standard normal. Translation, this means that a larger value is more likely to occur under a t-distribution than a standard normal. Furthermore, the heavy tails imply more variance than the standard normal.

For α between 0 and 1, let $t_{\alpha,n}$ be such that

$$P(T_n \geq t_{\alpha,n}) = \alpha$$

By symmetry around the origin,

$$P(T_n \leq -t_{\alpha,n}) = \alpha$$

$$\text{or, } P(T_n \geq -t_{\alpha,n}) = 1 - \alpha$$

$$\text{and, } -t_{\alpha,n} = t_{1-\alpha,n}$$

These standard values are available in math charts since they form the basis of the t test.

17.1 Mean and Variance

The following are stated without proof

$$E[T_n] = 0$$

$$Var(T_n) = \frac{n}{n-2}$$




Figure 3: visual representation of $t_{\alpha, n}$

In the limit of large n , the variance is close to 1, which is consistent with the fact that the distribution resembles a standard normal in that limit.

18 F-Distribution

If χ_n^2 and χ_m^2 are two independent chi-squared distributions with n and m degrees of freedom respectively, then the variable $F_{n, m}$ defined as

$$F_{n, m} = \frac{\chi_n^2/n}{\chi_m^2/m}$$

is said to have an F-distribution with n and m degrees of freedom.

For any α between 0 and 1, we define $F_{\alpha, n, m}$ as

$$P(F_{n, m} > F_{\alpha, n, m}) = \alpha$$

These values are available in standard tables for different combinations of α, n and m .

Consider

$$\begin{aligned} \alpha &= P\left(\frac{\chi_n^2/n}{\chi_m^2/m} > F_{\alpha, n, m}\right) \\ &= P\left(\frac{\chi_m^2/m}{\chi_n^2/n} < \frac{1}{F_{\alpha, n, m}}\right) \\ &= 1 - P\left(\frac{\chi_m^2/m}{\chi_n^2/n} \geq \frac{1}{F_{\alpha, n, m}}\right) \\ \text{or, } P\left(\frac{\chi_m^2/m}{\chi_n^2/n} \geq \frac{1}{F_{\alpha, n, m}}\right) &= 1 - \alpha \\ \text{or, } P(F_{m, n} \geq F_{1-\alpha, m, n}) &= 1 - \alpha \\ \text{Hence, } \frac{1}{F_{\alpha, n, m}} &= F_{1-\alpha, m, n} \end{aligned}$$

19 Logistics Distribution

A random variable X is said to have a logistics distribution with parameters μ and v if its cumulative density function is of the form

$$F_X(x) = \frac{e^{(x-\mu)/v}}{1 + e^{(x-\mu)/v}}, \quad \forall -\infty < x < \infty$$

Differentiating to get the density function

$$f_X(x) = \frac{e^{(x-\mu)/v}}{v(1 + e^{(x-\mu)/v})^2}, \quad \forall -\infty < x < \infty$$

19.1 Mean

$$\boxed{E[X] = \mu}$$

$v =$ dispersion parameter

20 Markov Process

Markov Process is a discrete time process that is not memoryless. Here the random variable takes several possible states, and the probability distribution is defined in such a way that $P(\text{transition from state 1 to state 2})$ is dependent on state 1.

Let X_n be the random variable denoting the state after n transitions and X_0 will represent the starting state (which can be given or random). Markov assumption states that *Given the current state, past does not matter*. Armed with these,

$$\begin{aligned} p_{ij} &= P(\text{next state } j \mid \text{current state } i) \\ p_{ij} &= P(X_{n+1} = j \mid X_n = i) = P(X_{n+1} = j \mid X_n = i, X_{n-1}, \dots, X_0) \\ r_{ij}(n) &= P(X_n = j \mid X_0 = i) && \text{or, in state } j \text{ after } n \text{ steps} \\ r_{ij}(n) &= \sum_{k=1}^m r_{ik}(n-1)p_{kj} \end{aligned}$$

20.1 Recurring and Transient States

A state i is called *recurrent* if, starting from i , and travelling anywhere, it is always possible to return to i . If a state is not recurrent, it is *transient*. States in a recurrent class are periodic if they can be grouped into $d > 1$ groups so that all transitions from one group lead to the next group.

20.2 Steady State Probabilities

Do $r_{ij}(n)$ converge to some π_j (independent of i) ?

Yes if,

- recurrent states are all in a single class
- single recurrent class is not periodic (otherwise oscillations are possible)

Assuming yes,


$$\begin{aligned} r_{ij}(n) &= \sum_k r_{ik}(n-1)p_{kj} \\ \lim_{n \rightarrow \infty} r_{ij}(n) &= \sum_k r_{ik}(n-1)p_{kj} \\ \pi_{ij} &= \sum_k \pi_{ik}p_{kj} && \text{balance equations} \\ \sum_i \pi_i &= 1 \end{aligned}$$

frequency of transitions $k \rightarrow j = \pi_k p_{kj}$ in one step


frequency of transitions into $j = \sum_k \pi_k p_{kj}$ influx from all connected states

20.3 Birth Death Process

Consider the checkout counter example. The states are represented by the number of people currently being processed, and we always move n to $[n-1, n, n+1]$, i.e., either the people in the queue decrease by one, remain same or increase by one. Let the probability for moving up be p and moving down be q .



Let's estimate the steady state probabilities. Consider the following diagram splitting the chain into two parts through the two adjacent states



In this case, to maintain steady state, long term frequency of left-right transition should be same as right left transition, i.e., $\pi_i p_i = \pi_{i+1} q_i$

In the special case of $p_i = p$ and $q_i = q \forall i$,

$$\rho = \frac{p}{q} \quad \text{load factor}$$

$$\pi_{i+1} = \pi_i \frac{p}{q} = \pi_i \rho$$

$$\pi_i = \pi_0 \rho^i \quad i = 0, \dots, m$$

$$\text{Using } \sum_{i=0}^m \pi_0 \rho^i = 1,$$

$$\pi_0 = \frac{1}{\sum_{i=0}^m \rho^i}$$

if $p < q$ and $m \rightarrow \infty$,

$$\pi_0 = 1 - \rho$$

$$\pi_i = (1 - \rho) \rho^i$$

$$E[X_n] = \frac{\rho}{1 - \rho} \quad \text{Exponential Distribution}$$

When $\rho = 1$ or $p = q$, then all states are equally likely - symmetric random walk.

20.4 Absorption Probabilities

let a_i denote the probability of absorption and μ_i denote the expected no of steps until absorption starting from state i . Then,

$$a_i = \sum_j a_j p_{ij} \quad \text{outflux to the possible states}$$

$$\mu_i = 1 + \sum_j \mu_j p_{ij}$$

For multiple absorption states, we can possibly consider them together as a group and calculate the relevant quantities.

For a given state s ,

$$\begin{aligned} E[\text{steps to first time reach } s \text{ from } i] &= t_i \\ t_i &= E[\min\{n \geq 0 \text{ such that } X_n = s\}] \\ t_s &= 0 \\ t_i &= 1 + \sum_j t_j p_{ij} \quad \text{outflux to all possible states} \end{aligned}$$

Mean recurrence time (mean time to reach back a state) for s

$$\begin{aligned} t_s^* &= E[\min\{n \geq 1 \text{ such that } X_n = s\} | X_0 = s] \\ t_s^* &= 1 + \sum_j t_j p_{sj} \end{aligned}$$

21 Central Limit Theorem

21.1 Weak Law of Large Numbers

Suppose we want to know the mean height of penguins in the world. The absolutely correct answer can be obtained by taking the average of the entire population. But this is not practical, and often we will have to resort to estimating the quantity through a sample. Let there be n penguins in the sample and X_1, X_2, \dots, X_n be the random variables denoting their heights. Then,

$$\begin{aligned} M_n &= \frac{X_1 + X_2 + \dots + X_n}{n} \\ \lim_{n \rightarrow \infty} E[M_n] &= E[X] = \text{The true mean} \end{aligned}$$

21.2 Markov Inequality/Chebychev Inequality

For nonnegative random variable X ,

$$\begin{aligned} E[X] &= \sum_x x p_X(x) \geq \sum_{x \geq a} x p_X(x) && \text{discrete case} \\ &= \int_x x p_X(x) \geq \int_{x \geq a} x p_X(x) && \text{continuous case} \end{aligned}$$

Applying the above set of inequalities to the variable $X - \mu$

$$E[(X - \mu)^2] \geq a^2 P((X - \mu)^2 \geq a^2)$$

$$\text{or, } \text{Var}(X) \geq a^2 P(|X - \mu| \geq a)$$

For continuous case,

$$\begin{aligned} \sigma^2 &= \int_{-\infty}^{\infty} (x - \mu)^2 f_X(x) dx \\ &\geq \int_{-\infty}^{\mu-c} (x - \mu)^2 f_X(x) dx + \int_{\mu+c}^{\infty} (x - \mu)^2 f_X(x) dx \\ &\geq c^2 P(|X - \mu| \geq c) \end{aligned}$$

Hence,

$$P(|X - \mu| \geq c^2) \leq \frac{\sigma^2}{c^2}$$

or,

$$\boxed{P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}} \quad \text{where } c = k\sigma$$

Going back to the problem of estimating the mean,

$$M_n = \frac{X_1 + X_2 + \cdots + X_n}{n}$$

$$E[M_n] = \frac{1}{n} \sum_{i=1}^n E[X_i] = \mu \quad \text{expectation of expectation}$$

$$\text{Var}(M_n) = \sum_{i=1}^n \text{Var}\left(\frac{X_i}{n}\right) = \frac{\sigma^2}{n} \quad \text{since } X_i \text{ are independent}$$

$$\boxed{P(|M_n - \mu| \geq \epsilon) \leq \frac{\sigma^2}{n\epsilon^2}}$$

or, as $n \rightarrow \infty$, $M_n - \mu \rightarrow 0$, ϵ is the error bound/confidence.

21.3 Central Limit Theorem

Chebychev's inequality gives a loose bound. We can do better with CLT. Let X be a random variable with mean μ and variance σ^2 , and let X_i be independent identically distributed random variables with the same distribution as X . Then,

$$S_n = X_1 + X_2 + \cdots + X_n$$

$$\begin{aligned} Z_n &= \frac{S_n - E[S_n]}{\sigma_n} \quad \text{random variable with mean 0 and variance 1} \\ &= \frac{S_n - nE[X]}{\sqrt{n}\sigma} \end{aligned}$$

$$\text{or, } S_n = \sqrt{n}\sigma Z_n + nE[X]$$

$$\text{In } \lim_{n \rightarrow \infty} Z_n \rightarrow Z \quad (\text{standard normal})$$

$$\text{or, } \boxed{Z = \frac{S_n - nE[X]}{\sqrt{n}\sigma}} \quad \text{only for CDF (no comment on PDF/PMF)}$$

$$\text{Thus, } \boxed{P(Z > c) = P\left(\frac{S_n - nE[X]}{\sqrt{n}\sigma} > c\right)}$$

By defining the confidence on how close we desire S_n to the actual mean, we can calculate the required value of the n using standard normal CDF tables. However, we need to have an estimate of variance of the distribution in order to do the estimate of n .

22 Distribution of Sample Mean and Variance

Let X_1, X_2, \dots, X_n be independent random variables from a distribution having mean μ and variance σ^2 . From the central limit theorem,

$$\frac{X_1 + X_2 + \dots + X_n - n\mu}{\sigma\sqrt{n}} \sim \mathcal{N}(0, 1)$$

or, the sum of the random variables follows the distribution of a standard normal as the value of n becomes large. Typically, the property starts to manifest as soon as n becomes around 30.

22.1 Sample Mean

Let the random variable \bar{X} denote the sample mean and is defined as

$$\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

Note that any scaled version of a normal distribution is also normal. Thus, **The mean of n independent random variables coming from the same distribution also follows a normal distribution for sufficiently large n .**

Note that sample mean is itself a random variable and thus has a distribution. This happens because the quantity itself is the average of several random variables, which are instances of the same probability distribution.

$$E[\bar{X}] = E\left[\frac{X_1 + X_2 + \dots + X_n}{n}\right] \quad (1)$$

$$= \frac{1}{n} \sum_{i=1}^n E[X_i] \quad (2)$$

$$\boxed{E[\bar{X}] = \mu} \quad (3)$$

Similarly, the variance of the sample mean can be computed as follows

$$\begin{aligned} Var(\bar{X}) &= Var\left(\frac{X_1 + X_2 + \dots + X_n}{n}\right) \\ &= Var\left(\frac{X_1}{n}\right) + Var\left(\frac{X_2}{n}\right) + \dots + Var\left(\frac{X_n}{n}\right) \text{ using independence} \\ &= n \frac{\sigma^2}{n^2} \text{ using } Var(aX) = a^2 Var(X) \end{aligned}$$

$$\boxed{Var(\bar{X}) = \frac{\sigma^2}{n}}$$

Hence, for a population of mean μ and variance σ^2 , the $E[\text{sample mean}]$ is still μ but the variance of the sample mean shrinks by a factor of n . Stated in a different manner, this means that the spread of the sample mean reduces as we take the mean from more and more observations. This directly translates into the fact that our confidence on the estimate of the sample mean increases with more observations.

22.2 Sample Variance

The sample variance is defined as

$$S^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}$$

where \bar{X} is the sample mean. Similar to the sample mean, this is also a random variable.

The mean of sample variance can be calculated as follows

$$\begin{aligned} E[S^2] &= E\left[\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}\right] \\ &= \frac{1}{n-1} \sum_{i=1}^n E[(X_i - \bar{X})^2] \\ &= \frac{1}{n-1} \sum_{i=1}^n E[X_i^2 - 2X_i\bar{X} + \bar{X}^2] \\ &= \frac{1}{n-1} \left(\sum_{i=1}^n E[X_i^2] - E[2\bar{X} \sum_{i=1}^n X_i] + \sum_{i=1}^n E[\bar{X}^2] \right) \\ &= \frac{1}{n-1} \left(\sum_{i=1}^n E[X_i^2] - E[2n\bar{X}^2] + nE[\bar{X}^2] \right) \end{aligned}$$

Using $E[X^2] = \text{Var}(X) + E[X]^2$,

$$\begin{aligned} E[S^2] &= \frac{1}{n-1} \left(\sum_{i=1}^n E[\text{Var}(X_i) + E[X_i]^2] - nE[\bar{X}^2] \right) \\ &= \frac{1}{n-1} (n\sigma^2 + n\mu^2 - n(\frac{\sigma^2}{n} + \mu^2)) \end{aligned}$$

$$\boxed{E[S^2] = \sigma^2}$$

i.e., the mean of the sample variance is same as the variance of the distribution (population variance).

22.3 Distributions for a Normal Population

Consider X_1, X_2, \dots, X_n be independently derived from a normal population with mean μ and variance σ^2

i.e., $X_i \sim \mathcal{N}(\mu, \sigma^2) \forall i = 1, 2, \dots, n$

Based on the derivations above,

$$\begin{aligned} E[\bar{X}] &= \mu \\ \text{Var}(\bar{X}) &= \frac{\sigma^2}{n} \end{aligned}$$

And since the sum of normal random variables is also normal,

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$$

which is similar to the central limit theorem.

From the derivation above for the sample variance,

$$E[S^2] = \sigma^2$$

Now let's calculate the distribution of S^2

$$\begin{aligned}
S^2 &= \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1} \\
(n-1)S^2 &= \sum_{i=1}^n (X_i - \bar{X})^2 \\
&= \sum_{i=1}^n ((X_i - \mu) - (\bar{X} - \mu))^2 \\
&= \sum_{i=1}^n ((X_i - \mu)^2 + (\bar{X} - \mu)^2 - 2(X_i - \mu)(\bar{X} - \mu)) \\
&= \sum_{i=1}^n (X_i - \mu)^2 + n(\bar{X} - \mu)^2 - 2(\bar{X} - \mu) \sum_{i=1}^n (X_i - \mu) \\
&= \sum_{i=1}^n (X_i - \mu)^2 + n(\bar{X} - \mu)^2 - 2n(\bar{X} - \mu)^2 \\
&= \sum_{i=1}^n (X_i - \mu)^2 - n(\bar{X} - \mu)^2 \\
\frac{(n-1)S^2}{\sigma^2} &= \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma}\right)^2 - \left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}\right)^2 \text{ to make standard normals} \\
\text{or, } \frac{(n-1)S^2}{\sigma^2} + \left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}\right)^2 &= \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma}\right)^2
\end{aligned}$$

The right hand side is a chi-square variable with n degrees of freedom and the second part of the right hand side is a chi-square variable with 1 degree of freedom. We know that sum of independent chi-square variables is also a chi-square variable with degrees of freedom equal to the sum of individual degrees of freedom. Hence, it follows that

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$$

and also the fact that **for a normal population, the sample mean and sample variance are independent variables with normal and chi-square distributions respectively**. This independence is a unique property for a normal distribution and is useful in parameter estimation and hypothesis testing.

Another interesting observation from the above derivations is

$$\boxed{\sqrt{n} \frac{\bar{X} - \mu}{S} \sim t_{n-1}}$$

whereas $\sqrt{n} \frac{\bar{X} - \mu}{\sigma} \sim \mathcal{N}(0, 1)$

Note that the denominator in the first equation is sample variance. The derivation is

$$\begin{aligned} \frac{Z}{\sqrt{\chi_n^2/n}} &\sim t_n \text{ definition} \\ \text{or, } \frac{\frac{\bar{X}-\mu}{\sigma/\sqrt{n}}}{\sqrt{\frac{(n-1)S^2}{\sigma^2} \frac{1}{n-1}}} &\sim t_{n-1} \\ \text{or, } \sqrt{n} \frac{\bar{X}-\mu}{S} &\sim t_{n-1} \end{aligned}$$

23 Parameter Estimation

In probability theory, we usually have the distribution known to us and we try to use this information to obtain theoretical results applicable on population of this distribution. However, in statistics, we are given the samples drawn from the population, and we are interested in estimating the parameters of this population. For instance, it can be known that the samples are drawn from a normal distribution and the objective is to use the samples to obtain the mean and variance of the normal distribution. It is possible to partially know the parameters in some cases, for example mean is unknown but the variance is known. Following are some methods to obtain the *estimates* of these parameters.

23.1 Maximum Likelihood Estimator

Maximum Likelihood Estimator or MLE is based on the idea to **find that value of θ that maximizes the probability of observing the given set of samples of the population.** Alternately, let x_i for $i = 1, 2, \dots, n$ be n samples drawn from a population whose distribution is parametrized by θ (can be a vector as well). Then we define the likelihood function as

$$\text{likelihood} = f(x_1, x_2, \dots, x_n | \theta)$$

i.e., the joint probability (or density) of occurrence of all the samples under the given distribution for some value of θ . We aim to maximize this likelihood to get the estimate of θ . It is often the case that taking logarithm of both sides allows for an easy way of estimation. Note that both likelihood and log likelihood are maximized by the same value of estimator.

23.1.1 MLE for Bernoulli Variable

Suppose we observe n independent samples from a Bernoulli process and the aim is to find the MLE for the probability of success.

Let p denote the probability of success. Then,

$$\begin{aligned} P(X_i = x) &= p^x (1-p)^{1-x} \text{ where } x \text{ is } 0 \text{ or } 1 \\ \text{or, } P(X_i = x_i) &= p^{x_i} (1-p)^{1-x_i} \end{aligned}$$

Since all the samples are independent, the joint probability or likelihood is simply the product of all the probabilities

$$\begin{aligned} f(x_1, x_2, \dots, x_n | p) &= p^{x_1} (1-p)^{1-x_1} \dots p^{x_n} (1-p)^{1-x_n} \\ &= p^{\sum_{i=1}^n x_i} (1-p)^{n-\sum_{i=1}^n x_i} \\ \log(f(x_1, x_2, \dots, x_n | p)) &= \sum_{i=1}^n \log(p) + (n - \sum_{i=1}^n x_i) \log(1-p) \end{aligned}$$

Taking the derivative with respect to p to maximize,

$$\frac{d}{dp} \log(f(x_1, x_2, \dots, x_n | p)) = \sum_{i=1}^{x_i} \frac{1}{p} - (n - \sum_{i=1}^{x_i}) \frac{1}{1-p} = 0$$

$$\text{or, } \boxed{\hat{p} = \frac{\sum_{i=1}^n x_i}{n}}$$

which is the proportion of successful trials in the sample.

23.1.2 MLE for Poisson Variable

Suppose that we observe n random samples obtained from a poisson process. Recall

$$P(X = k) = \frac{e^{-\lambda} \lambda^k}{k!}$$

Hence, the joint distribution can be written as

$$f(x_1, x_2, \dots, x_n | p) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{x_i}}{x_i!}$$

$$f(x_1, x_2, \dots, x_n | p) = \frac{e^{-n\lambda} \lambda^{\sum_{i=1}^n x_i}}{\prod_{i=1}^n x_i!}$$

$$\log(f(x_1, x_2, \dots, x_n | p)) = -n\lambda + \left(\sum_{i=1}^n x_i\right) \log(\lambda) - \sum_{i=1}^n \log(x_i!)$$

$$\frac{d}{dp} (f(x_1, x_2, \dots, x_n | p)) = -n + \left(\sum_{i=1}^n x_i\right) \frac{1}{\lambda}$$

$$= 0$$

$$\text{or, } \boxed{\hat{\lambda} = \frac{\sum_{i=1}^n x_i}{n}}$$

i.e., the average rate is simply the average of all the observed arrivals.

23.1.3 MLE for Normal Variable

Suppose we observe n samples from a normal population whose mean is μ and variance is σ^2 . We will aim to obtain MLE estimates for both the mean and variance. Recall

$$P(X = x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right)$$

Hence, the likelihood will be

$$\begin{aligned}
f(x_1, x_2, \dots, x_n | \mu, \sigma^2) &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x_i - \mu)^2}{2\sigma^2}\right) \\
\log(f(x_1, x_2, \dots, x_n | \mu, \sigma^2)) &= -\frac{n}{2} \log(2\pi) - n \log(\sigma) - \frac{1}{2\sigma^2} \left(\sum_{i=1}^n (x_i - \mu)^2\right) \\
\frac{d}{d\mu}(\log(f(x_1, x_2, \dots, x_n | \mu, \sigma^2))) &= -\frac{1}{\sigma^2} \left(\sum_{i=1}^n (x_i - \mu)\right) \\
&= 0 \\
\text{or, } \boxed{\hat{\mu} &= \frac{\sum_{i=1}^n x_i}{n}} \\
\frac{d}{d\sigma}(\log(f(x_1, x_2, \dots, x_n | \mu, \sigma^2))) &= -\frac{n}{\sigma} - \frac{1}{2\sigma^2} \left(\sum_{i=1}^n (x_i - \mu)^2\right) \\
&= 0 \\
\text{or, } \boxed{\hat{\sigma}^2 &= \frac{\sum_{i=1}^n (x_i - \mu)^2}{n}} \\
&= \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n} \\
\text{where } \bar{x} &= \frac{\sum_{i=1}^n x_i}{n}
\end{aligned}$$

Note that the estimator for variance is different from the sample variance

$$\begin{aligned}
\text{MLE } \sigma^2 &= \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n} \\
S^2 &= \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n-1}
\end{aligned}$$

23.1.4 MLE for Uniform Random Variable

Consider observing n samples from a uniform distribution with the parameter θ . Then,

$$f(x_1, x_2, \dots, x_n | \theta) = \frac{1}{\theta^n}$$

which is clearly maximized when θ is minimum. But since θ has to be at least as large as the maximum observed value,

$$\boxed{\hat{\theta} = \max(x_1, x_2, \dots, x_n)}$$

23.2 Interval Estimates

The MLE estimates calculated above are estimates and do not reflect the true value. We expect the true value of the parameter to be close to the estimate, but not exactly equal to it. Hence, it makes sense to give an interval instead of a single estimate to reflect our confidence in the estimated value of the parameter.

Note that the below confidence intervals imply that α percent of times, the constructed interval will contain the true mean μ , when the same calculation is repeated with multiple samples. The calculations of intervals do not imply that μ is contained in the interval with α confidence. We calculate an interval that falls on μ rather than telling the interval that μ falls in.

23.2.1 Confidence interval for Mean of Normal Distribution when Variance is Known

Consider the problem of estimation of the mean of a normal distribution with known variance σ^2 . Since we know that the MLE for mean is just the sample mean, and the sample mean follows a normal distribution,

$$P(-1.96 < \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} < 1.96) = 0.95 \text{ using standard normal tables}$$

or, $P(\bar{X} - 1.96 \frac{\sigma}{\sqrt{n}} < \mu < \bar{X} + 1.96 \frac{\sigma}{\sqrt{n}}) = 0.95$

Thus, we are 95% confident that the true value of the mean lies in the range

$$(\bar{X} - 1.96 \frac{\sigma}{\sqrt{n}}, \bar{X} + 1.96 \frac{\sigma}{\sqrt{n}})$$

This trick to calculate the confidence interval can be generalized for any value of confidence. Recall that

$$P(Z > z_\alpha) = \alpha$$

$$P(Z < -z_\alpha) = \alpha$$

Hence, for a given confidence level α , the two sided confidence interval is

$$P(-z_{\alpha/2} < Z < z_{\alpha/2}) = 1 - \alpha$$

$$P(-z_{\alpha/2} < \sqrt{n} \frac{\bar{X} - \mu}{\sigma} < z_{\alpha/2}) = 1 - \alpha$$

$$P(\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}) = 1 - \alpha$$

two sided $1 - \alpha$ confidence interval = $(\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}})$

where \bar{x} is the observed value of \bar{X} .




Figure 4: Visualization of the double sided confidence interval on standard normal.

In a very similar manner, we can calculate the one sided confidence interval. Here, we are only

interested in the lower or upper bound of the said interval. The other side is inf or $-\inf$.

$$\begin{aligned}
P(Z > z_\alpha) &= \alpha \\
P(\sqrt{n} \frac{\bar{X} - \mu}{\sigma} > z_\alpha) &= \alpha \\
P(\mu < \bar{x} + z_\alpha \frac{\sigma}{\sqrt{n}}) &= \alpha
\end{aligned}$$

Lower $1 - \alpha$ confidence interval $= (-\inf, \bar{x} + z_\alpha \frac{\sigma}{\sqrt{n}})$

$$\begin{aligned}
P(Z < -z_\alpha) &= \alpha \\
P(\sqrt{n} \frac{\bar{X} - \mu}{\sigma} < -z_\alpha) &= \alpha \\
P(\bar{x} - z_\alpha \frac{\sigma}{\sqrt{n}} < \mu) &= \alpha
\end{aligned}$$

Upper $1 - \alpha$ confidence interval $= (\bar{x} - z_\alpha \frac{\sigma}{\sqrt{n}}, \inf)$

The interpretation of the right sided confidence interval is that we are $1 - \alpha$ confident that the value of the mean is more than the lower end of the interval. In a similar way, the left side interval gives the upper bound on the value of mean with the desired confidence.

23.2.2 Confidence interval for Mean of Normal Distribution when Variance is Unknown

The derivation of confidence intervals in this case is similar to the above, with the only difference of using a t-distribution. Recall

$$\sqrt{n} \frac{\bar{X} - \mu}{S} \sim t_{n-1}$$

where S is the sample variance. Following similar derivation to the known variance case,

$$\text{two sided } 1 - \alpha \text{ confidence interval} = (\bar{x} - t_{\alpha/2, n-1} \frac{s}{\sqrt{n}}, \bar{x} + t_{\alpha/2, n-1} \frac{s}{\sqrt{n}})$$

$$\text{Lower } 1 - \alpha \text{ confidence interval} = (-\inf, \bar{x} + t_{\alpha, n-1} \frac{s}{\sqrt{n}})$$

$$\text{Upper } 1 - \alpha \text{ confidence interval} = (\bar{x} - t_{\alpha, n-1} \frac{s}{\sqrt{n}}, \inf)$$

where s is the observed value of the sample variance S . However, notice that the intervals calculated will usually be larger than those when the variance is known because t-distribution is heavier tailed than a standard normal and thus has higher variance.

23.2.3 Confidence interval for Variance of Normal Distribution when Mean is Unknown

Recall that

$$(n-1) \frac{S^2}{\sigma^2} \sim \chi_{n-1}^2$$




Figure 5: Visualization of the double sided confidence interval on standard normal.

By noting that σ^2 is always positive, we have the following

$$\begin{aligned} \text{two sided } 1 - \alpha \text{ confidence interval} &= \left(\frac{(n-1)s^2}{\chi_{\alpha/2, n-1}^2}, \frac{(n-1)s^2}{\chi_{1-\alpha/2, n-1}^2} \right) \\ \text{Lower } 1 - \alpha \text{ confidence interval} &= \left(0, \frac{(n-1)s^2}{\chi_{1-\alpha, n-1}^2} \right) \\ \text{Upper } 1 - \alpha \text{ confidence interval} &= \left(\frac{(n-1)s^2}{\chi_{\alpha, n-1}^2}, \text{inf} \right) \end{aligned}$$

23.2.4 Estimating Difference in Means of Two Normal Populations

Suppose the following two independent sets of random variables

$$\begin{aligned} X_1, X_2, \dots, X_n &\sim \mathcal{N}(\mu_1, \sigma_1^2) \\ Y_1, Y_2, \dots, Y_m &\sim \mathcal{N}(\mu_2, \sigma_2^2) \end{aligned}$$

Then, we are interested in the distribution of $\mu_1 - \mu_2$. It is intuitive to see that the MLE estimator of this quantity is nothing but $\bar{X} - \bar{Y}$. Also, since \bar{X} and \bar{Y} are both normally distributed,

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_1 - \mu_2, \frac{\sigma_1^2}{n} + \frac{\sigma_2^2}{m}\right)$$

Consequently, using the confidence intervals derived for the case of a mean of a single normal distribution, we have the following intervals when the standard deviations are known

$$\begin{aligned} \text{two sided } 1 - \alpha \text{ confidence interval} &= (\bar{x} - \bar{y} - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n} + \frac{\sigma_2^2}{m}}, \bar{x} - \bar{y} + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n} + \frac{\sigma_2^2}{m}}) \\ \text{Lower } 1 - \alpha \text{ confidence interval} &= (-\text{inf}, \bar{x} - \bar{y} + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n} + \frac{\sigma_2^2}{m}}) \\ \text{Upper } 1 - \alpha \text{ confidence interval} &= (\bar{x} - \bar{y} - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n} + \frac{\sigma_2^2}{m}}, \text{inf}) \end{aligned}$$

where \bar{x} and \bar{y} are estimates of \bar{X} and \bar{Y} respectively.

A more challenging scenario arises when the variances are not known. In that case, it is only

logical to try to estimate the intervals using sample variances (themselves random variables)

$$S_1^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}$$

$$S_2^2 = \frac{\sum_{i=1}^m (Y_i - \bar{Y})^2}{m-1}$$

However, the distribution useful for deriving the confidence intervals

$$\frac{\bar{X} - \bar{Y} - (\mu_1 - \mu_2)}{\sqrt{\frac{S_1^2}{n} + \frac{S_2^2}{m}}}$$

is complicated and depends on the unknown variances. The variable is friendly if we assume that the two unknown variances are both same.

Assuming $\sigma_1 = \sigma_2 = \sigma$, it follows

$$(n-1) \frac{S_1^2}{\sigma^2} \sim \chi_{n-1}^2$$

$$(m-1) \frac{S_2^2}{\sigma^2} \sim \chi_{m-1}^2$$

$$(n-1) \frac{S_1^2}{\sigma^2} + (m-1) \frac{S_2^2}{\sigma^2} \sim \chi_{n+m-2}^2$$

since S_1^2 and S_2^2 are independent chi-square random variables and from section 16 that the sum of such variables is also chi-square.

We already know

$$\frac{\bar{X} - \bar{Y} - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma^2}{n} + \frac{\sigma^2}{m}}} \sim \mathcal{N}(0, 1)$$

and we also know that the ratio of a standard normal and the square root of a chi-square divided by it's degrees of freedom is a t-distribution

$$\frac{\bar{X} - \bar{Y} - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma^2}{n} + \frac{\sigma^2}{m}}} \div \sqrt{\frac{(n-1) \frac{S_1^2}{\sigma^2} + (m-1) \frac{S_2^2}{\sigma^2}}{n+m-2}} \sim t_{n+m-2}$$

Let

$$S_p = \frac{(n-1)S_1^2 + (m-1)S_2^2}{n+m-2}$$

Then,

$$P(-t_{\alpha/2, n+m-2} \leq \frac{\bar{X} - \bar{Y} - (\mu_1 - \mu_2)}{S_p \sqrt{\frac{1}{n} + \frac{1}{m}}} \leq t_{\alpha/2, n+m-2}) = 1 - \alpha$$

Two sided $1 - \alpha$ confidence interval $= (\bar{x} - \bar{y} - t_{\alpha/2, n+m-2} s_p \sqrt{\frac{1}{n} + \frac{1}{m}}, \bar{x} - \bar{y} + t_{\alpha/2, n+m-2} s_p \sqrt{\frac{1}{n} + \frac{1}{m}})$

where s_p is the sample estimate for S_p . Lower confidence interval is derived in a similar fashion to the previous derivations, but the upper confidence interval is the lower confidence interval of $\mu_2 - \mu_1$.

23.2.5 Confidence Interval for Mean of Bernoulli Random Variable

Suppose we obtain a sample of n independent Bernoulli random variables, where the probability of success is p . Let X denote the no of successes. Using the CLT for large n ,

$$\frac{X - np}{\sqrt{np(1-p)}} \sim \mathcal{N}(0, 1)$$

It is not tractable to calculate the confidence intervals from this formulation. Let $\hat{p} = X/n$ denote the MLE of the mean p . Substituting in the denominator of above,

$$P(-z_{\alpha/2} \leq \frac{X - np}{\sqrt{n\hat{p}(1-\hat{p})}} \leq z_{\alpha/2}) \approx 1 - \alpha$$

$$\text{Two sided } 1 - \alpha \text{ confidence interval} \approx (\hat{p} - z_{\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + z_{\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}})$$

and one sided confidence intervals can be obtained in similar manner to previous derivations.

24 Linear Regression

We are given pairs of data $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ (all independent) where we assume that x and y are governed by the linear relation

$$y \approx \theta_0 + \theta_1 x$$

The aim is to determine the model which is parametric consisting of two parameters θ_0 and θ_1 . We find it using the least squares estimate, i.e., minimizing

$$\underset{\theta_0, \theta_1}{\text{minimize}} \sum_{i=1}^n (y_i - \theta_0 - \theta_1 x_i)^2$$

The true model also includes noise and is given by

$$Y_i = \theta_0 + \theta_1 X_i + W_i$$

where we assume the noise $W_i \sim \mathcal{N}(0, \sigma^2)$ and is independently and identically distributed. Observing some X and Y is same as observing the noise.

$$P(X = x, Y = y) = P(W = y - \theta_0 - \theta_1 x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(y - \theta_0 - \theta_1 x)^2}{2}\right)$$

$$\begin{aligned} P(X_1 = x_1, Y_1 = y_1, \dots, X_n = x_n, Y_n = y_n) &= \prod_{i=1}^n P(X_i = x_i, Y_i = y_i) \\ &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(y_i - \theta_0 - \theta_1 x_i)^2}{2}\right) \end{aligned}$$

Maximizing the above product is maximizing the likelihood of the occurrence of the data under the model parameters θ_0 and θ_1 . Since taking log will not change the maxima, we usually maximize the log likelihood

$$\underset{\theta_0, \theta_1}{\text{maximize}} \prod_{i=1}^n \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(y_i - \theta_0 - \theta_1 x_i)^2}{2}\right) = \underset{\theta_0, \theta_1}{\text{minimize}} \sum_{i=1}^n (y_i - \theta_0 - \theta_1 x_i)^2$$

We can take derivatives with respect to the parameters of the above function to get the estimate for the parameters as

$$\begin{aligned}\bar{x} &= \frac{1}{n} \sum_{i=1}^n x_i, \bar{y} &= \frac{1}{n} \sum_{i=1}^n y_i \\ \hat{\theta}_1 &= \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{E[(X - \bar{X})(Y - \bar{Y})]}{E[(X - \bar{X})^2]} = \frac{Cov(X, Y)}{Var(X)} \\ \hat{\theta}_0 &= \bar{y} - \hat{\theta}_1 \bar{x}\end{aligned}$$

The above formulae can also be derived if the additives are a function of X . Since the linear relationship will still be respected and the loglikelihood can be maximized to get the estimates of the parameters.

25 Exercises

25.1 Problems

1. Independence in Complements

Given $A \perp B$, show $A \perp B^c$ and $A^c \perp B^c$. [Solution](#)

2. Conditional Independence

A, B , and C are independent with $P(C) > 0$. Show that $A \perp B|C$. [Solution](#)

3. Geometry of Meeting

R and J have to meet at a given place and each will arrive at the given place independent of each other with a delay of 0 to 1hr uniformly distributed. The pairs of delays are all equally likely. The first to arrive waits for 15 minutes and leaves. What is the probability of meeting ? [Solution](#)

4. Expectation of Function

Let X and Y be random variables with $Y = g(X)$. Show $E[Y] = \sum_x g(x)p_X(x)$. [Solution](#)

5. Cumulative Distribution Function

A random variable X is a combination of a continuous and discrete distribution as follows

$$f_X(x) = \begin{cases} 0.5 & a \leq x \leq b \\ 0.5 & x = 0.5 \\ 0 & \text{otherwise} \end{cases}$$

Find the Cumulative Distribution of X . [Solution](#)

6. Number of tosses till first head

When tossing a fair coin, what is the $E[\# \text{ tosses till the first H}]$. [Solution](#)

7. Iterated Expectation Proof

For discrete variables, show $E[X] = E[E[X|Y]]$. [Solution](#)

8. Iterated Expectation for three variables

For three random variables X, Y and Z , show $E[Z|X] = E[E[Z|X, Y]|X]$. [Solution](#)

9. Iterated Expectation practice

A class has two sections denoted by the random variable Y . Let X denote the quiz score of a student. Given that section 1 has 10 students, section 2 has 20 students, $E[X|Y = 1] = 90, E[X|Y = 2] = 60, Var(X|Y = 1) = 10, Var(X|Y = 2) = 20$, find $E[X]$ and $Var(X)$. [Solution](#)

10. **Hat Problem**

n people throw their hats in a box and then pick a hat at random. What is the expected number of people who pick their own hat ? [Solution](#)

11. **Breaking a stick**

A stick of length l is broken first at X uniformly chosen between $[0, l]$, and then at Y , uniformly chosen between $[0, X]$. Find the expected length of the shorter part. [Solution](#)

12. **Convolution of Exponentials**

Suppose $X \sim \exp(\lambda)$ and $Y \sim \exp(\mu)$, find the probability distribution $p_{X+Y}(x)$.

13. **Triangles from a Stick**

We have a stick of length 1. We randomly choose two points on the stick and break the stick at those points. Calculate the probability that the three pieces form a triangle. [Solution](#)

14. **PMF of $g(X)$**

Let X be uniform in $[0, 2]$, then find the PMF of $Y = X^3$. [Solution](#)

15. **Waiting for Taxi**

A taxi stand and bus stop near Al's home are at the same location. Al goes there and if a taxi is waiting $P = \frac{2}{3}$, he boards it. Otherwise, he waits for a taxi or bus to come, whichever is first. Taxi takes anywhere between 0 to 10 mins (uniform) while a bus arrives in exactly 5 mins. He boards whichever is first. Find CDF and $E[\text{wait time}]$. [Solution](#)

16. **Bayes Theorem**

Let Q be a continuous random variable with PDF

$$f_Q(q) = \begin{cases} 6q(1-q) & 0 \leq q \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

where Q represents $P(\text{success})$ for a Bernoulli X , i.e., $P(X = 1|Q = q) = q$. Find $f_{Q|X}(q|x) \forall x \in [0, 1]$ and q . [Solution](#)

17. **A Normal Transformation**

Let $X \sim \mathcal{N}(0, 1)$ and $Y = g(X)$. Find $p_Y(y)$.

$$g(t) = \begin{cases} -t & t \leq 0 \\ \sqrt{t} & t > 0 \end{cases}$$

[Solution](#)

18. **Binomial Shooter**

A shooter takes 10 hits in a shooting range and each shot has $p = 0.2$ of hitting target independent of each other. Let X = number of hits. Find

- (a) PMF of X
- (b) $P(\text{no hits})$
- (c) $P(\text{scoring more than misses})$
- (d) $E[X]$ and $\text{Var}(X)$
- (e) Suppose the entry is \$3 and each shot fetches \$2. Let Y = profit. Find $E[Y]$ and $\text{Var}(Y)$.
- (f) Suppose entry is free and total reward is square of number of hits. Let Z be profit. Find $E[Z]$.

Solution

19. Mosquito and Tick

Every second, a mosquito lands with $P = 0.5$. Once it lands, it bites with $P = 0.2$. Let X be the time between successive mosquito bites. Find $E[X]$ and $Var(X)$.

Now suppose a tick comes into play independent of mosquito. It lands with $P = 0.1$ and once landed, bites with $P = 0.7$. Let Y be the time between successive bug bites. Find $E[Y]$ and $Var(Y)$. [Solution](#)

20. HH or TT

Given a coin with $P(H) = p$, find the $E[\text{number of tosses till } HH \text{ or } TT]$. [Solution](#)

21. A Three Coin Game

Let 3 fair coins be tossed at every turn. Given all coins and turns are independent, calculate the following (assuming success is defined as all three coins landing the same side up))

- (a) PMF of K , no of trials upto but not including the 2^{nd} success
- (b) E and Var of M , the $E[\text{number of tails}]$ before first success.

Solution

22. Linear Expectations

Bob conducts trials in a similar manner to Problem 21, but with four coins. He repeatedly removes a coin at success until just a single coin remains. Calculate the Expected number of tosses till the finish of experiment. [Solution](#)

23. Papers Drawn with Replacement

Suppose there are n papers in a drawer. We take one paper, sign it, and then put it back into the drawer. We take one more paper out and if it is not signed, we sign it and put it back in the drawer. If the paper is already signed, we simply put it back in the drawer. We repeat this process until all the papers are signed. Find the $E[\text{papers drawn till all papers are signed}]$. What is the value of this quantity as $n \rightarrow \text{large}$. [Solution](#)

24. A Three Variable Inequality

Let X, Y, Z be three exponentially distributed random variables with parameters λ, μ , and ν respectively. Find $P(X < Y < Z)$. [Solution](#)

25. Poisson Emails

You get emails according to a Poisson process at the rate of 5 messages/hour. You check email every 30 minutes. Find

- $P(\text{no new message})$
- $P(\text{one new message})$

Solution

26. Poisson Fishing

We go fishing where we catch fishes at the rate of $0.6/\text{hour}$. We fish for two hours. If we do not catch a fish in the first two hours, we fish until the first catch. Find the following

- $P(\text{fish for } > 2 \text{ hours})$
- $P(\text{fish for } > 2 \text{ but } < 5 \text{ hours})$
- $P(\text{catch at least two fish})$
- $E[\text{fish}]$

- $E[\text{Total fishing time}]$
- $E[\text{future fishing time—fished for two hours}]$

[Solution](#)

27. Poisson Lightbulbs

We have three identical but independent lightbulbs whose lifetimes are modelled by a Poisson process with parameter λ . Given that we start all the three bulbs together, find the $E[\text{time until last bulb dies out}]$. [Solution](#)

28. Two Poisson Lightbulbs


Beginning at $t = 0$, we begin using bulbs one at a time until failure. Any broken bulb is immediately replaced. Each new bulb is selected independently and equally likely from type A (exponential life with $\lambda = 1$) or type B (exponential life with $\lambda = 3$). Lifetimes of all bulbs are independent.

- Find $E[\text{time until first failure}]$.
- $P(\text{no bulb failure before time } t)$.
- Given that there are no failures until time t , determine the conditional probability that the first bulb used is of type A.
- Find the probability that the total illumination by two type B bulbs $>$ one type A.
- Suppose the process terminates after 12 bulbs fail. Determine the expected value and variance of the total illumination provided by type B bulbs while the process is in operation.
- Given there are no failures until time t , find the expected value of time until first failure.

[Solution](#)


29. Steady State Markov Process

Find the steady state probabilities of the following Markov Process




[Solution](#)

30. Absorption Probabilities



Calculate the absorption probabilities for state 4 and expected time to absorption from all states. (for absorption time, assume $p_{35} = 0$ and $p_{32} = 0.5$) [Solution](#)

31. Selecting Courses with Markov Process



Consider the above markov process for changing courses. The probability being in some course tomorrow given a course today is mentioned along the edges. Suppose we start with course 6-1 (Note that course 6 is the combination of courses 6-1, 6-2 and 6-3). Calculate the following

- $P(\text{eventually leaving course 6})$.
- $P(\text{eventually landing in course 15})$.
- $E[\text{number of days till leaving course 6}]$.
- At every switch for 6-2 to 6-1 or 6-3 to 6-1, we buy an ice cream (but a maximum of two). Calculate the $E[\text{number of ice creams before leaving course 6}]$.
- Suppose we end up in 15. What is the $E[\text{number of steps to reach 15}]$.

- (f) Suppose we don't want to take course 15. Accordingly, when in 6-1, we stay there with probability $1/2$ while other three options have equal probabilities. If we are in 6-2, probability of going to 6-1 and 6-3 are in the same ratio as before. Calculate the $E[\text{number of days until we enter course 9}]$.
- (g) Assuming $P(X_{n+1} = 15|X_n = 9) = P(X_{n+1} = 9|X_n = 15) = P(X_{n+1} = 15|X_n = 15) = P(X_{n+1} = 9|X_n = 9) = 1/2$, what is $P(X_n = 15)$ and $P(X_n = 9)$ far into the future.
- (h) Suppose $P(X_{n+1} = 6-1|X_n = 9) = 1/8$, $P(X_{n+1} = 9|X_n = 9) = P(X_{n+1} = 15|X_n = 15) = 7/8$. What is the $E[\text{number of days till return to 6-1}]$.

[Solution](#)

32. Estimating Binomial with CLT, $1/2$ correction

Given a Bernoulli Process with $n = 36$ and $p = 0.5$, find $P(S_n \leq 21)$. [Solution](#)

33. Sample Variance for Normal Distribution

The time it takes a central processing unit to process a certain type of job is normally distributed with mean 20 seconds and standard deviation 3 seconds. If a sample of 15 such jobs is observed, what is the probability that the sample variance will exceed 12 ?

[Solution](#)

34. MLE Estimate

Suppose we observe n independent and identically distributed samples x_1, x_2, \dots, x_n from an exponential distribution. Estimate the parameter of the exponential. [Solution](#)

35. LMS Estimate

Given the prior $f_{\Theta|(\theta)}$, uniform in $[4, 10]$, and $f_{X|\Theta}(x|\theta)$ is uniform in $[\theta - 1, \theta + 1]$, estimate the posterior of θ . [Solution](#)

36. Probability Convergence

Let X be uniformly distributed between $[-1, 1]$. Let X_1, X_2, \dots, X_n be independently and identically distributed with the same distribution as X . Find whether the following sequences are convergent in probability and also find the limit.

- (a) X_i
 (b) $Y_i = X_i/i$
 (c) $Z_i = (X_i)^i$

[Solution](#)

25.2 Solutions

1. Question

- (a)

$$\begin{aligned}
 P(A \cap B) &= P(A)P(B) \\
 P(A) &= P((A \cap B) \cup (A \cap B^c)) \\
 &= P(A \cap B) + P(A \cap B^c) && \text{since disjoint} \\
 P(A \cap B^c) &= P(A) - P(A \cap B) \\
 &= P(A)(1 - P(B)) = P(A)P(B^c)
 \end{aligned}$$

(b)

$$\begin{aligned}(A \cup B)^c &= A^c \cap B^c \\ P(A^c \cap B^c) &= 1 - P(A \cup B) \\ &= 1 - P(A) - P(B) + P(A \cap B) \\ &= (1 - P(A))(1 - P(B)) \\ &= P(A^c)P(B^c)\end{aligned}$$

2. Question

$$P(A \cap B|C) = \frac{P(A \cap B \cap C)}{P(C)} = P(A)P(B) = P(A|C)P(B|C) \quad \text{Due to independence}$$

3. Question

Suppose R arrives at x hours. J has to arrive between x hrs to x hrs + 15 mins. Similarly if J arrives at y hours, R has to arrive between y hours to y hours + 15 mins. These are regions enclosed by the regions $x \leq 1, y \leq 1, y \leq x + \frac{1}{4}$ and $y \geq x - \frac{1}{4}$. The probability is then $1 - P(\text{not meeting}) = 1 - 2(\frac{1}{2} \frac{3}{4} \frac{3}{4}) = \frac{7}{16}$.

4. Question

$$\begin{aligned}E[Y] &= \sum_y y p_Y(y) = \sum_y \sum_{x:g(x)=y} p_X(x) = \sum_y \sum_{x:g(x)=y} y p_X(x) \\ &= \sum_y \sum_{x:g(x)=y} g(x) p_X(x) = \sum_x g(x) p_X(x)\end{aligned}$$

5. Question

Cumulative Distribution of X can be found by integration and is as follows

$$f_X(x) = \begin{cases} 0 & x < 0 \\ 0.5x & 0 \leq x < 0.5 \\ 0.75 & x = 0.5 \\ 0.75 + 0.5(x - 0.5) & 0.5 < x \leq 1 \\ 1 & 1 < x \end{cases}$$

6. Question

Let X be the # of tosses till first H . Then, $(X = 1) \cap (X > 1) = \phi$. Using *Total Expectation Theorem*

$$\begin{aligned}E[X] &= P(X = 1)E[X|X = 1] + P(X > 1)E[X|X > 1] \\ &= 0.5 * 1 + 0.5E[X] \\ \Rightarrow E[X] &= 2\end{aligned}$$

$P(X = 1) = 0.5$ because then we get the head in the first toss itself. Since $P(X = 1) + P(X > 1) = 1$, we have $P(X > 1) = 0.5$. $E[X] = E[X|X > 1]$ because the tosses are *independent* and thus memoryless.

7. [Question](#)

Note that $E[X|Y]$ is a function of y .

$$\begin{aligned}
 E[E[X|Y]] &= \sum_y E[X|Y]p_Y(y) \\
 &= \sum_y \sum_x xp_{X|Y}p_Y \\
 &= \sum_y \sum_x xp_{X,Y}(x,y) \\
 &= \sum_x x \sum_y p_{X,Y}(x,y) \\
 &= \sum_x xp_X(x) \\
 &= E[X]
 \end{aligned}$$

8. [Question](#)

Note that $E[Z|X, Y]$ will be a function of both X and Y .

$$\begin{aligned}
 E[Z|X, Y] &= \sum_z zp_{Z|X,Y}(z|x, y) \\
 E[E[Z|X, Y]|X] &= \sum_y E[Z|X, Y]p_{X,Y|X}(x, y|x) \\
 &= \sum_y \sum_z zp_{Z|X,Y}(z|x, y)p_{Y|X}(y|x) \\
 &= \sum_y \sum_z z \frac{p_{X,Y,Z}(x, y, z)}{p_X(x)} \\
 &= \sum_z z \sum_y \frac{p_{X,Y,Z}(x, y, z)}{p_X(x)} \\
 &= \sum_z z \frac{p_{X,Z}(x, z)}{p_X(x)} \\
 &= \sum_z zp_{Z|X}(z|x) \\
 &= E[Z|X]
 \end{aligned}$$

9. [Question](#)

We use the formulae from iterated expectation to calculate these.

$$\begin{aligned}
P_Y(y) &= \begin{cases} \frac{1}{3} & y = 1 \\ \frac{2}{3} & y = 2 \end{cases} \\
E[X] &= E[E[X|Y]] = \sum_y E[X|Y]P(Y) \\
&= 90 * \frac{1}{3} + 60 * \frac{2}{3} \\
Var(X) &= E[Var(X|Y)] + Var(E[X|Y]) \\
&= \sum_y Var(X|Y)P(Y) + ((90 - E[E[X|Y]])^2 \frac{1}{3} + (60 - E[E[X|Y]])^2 \frac{2}{3}) \\
&= \frac{650}{3}
\end{aligned}$$

10. Question

Let X denote the number of people who pick their own hat. We have been asked $E[X]$. Let X_i be a binary random variable denoting whether the i^{th} person picked their own hat, i.e.,

$$\begin{aligned}
X_i &= \begin{cases} 1 & \text{if } i^{th} \text{ person picks their own hat} \\ 0 & \text{otherwise} \end{cases} \\
P(X_i = 1) &= \frac{1}{n} \\
E[X_i] &= 1 * \frac{1}{n} + 0 * (1 - \frac{1}{n}) = \frac{1}{n}
\end{aligned}$$

Consequently

$$E[X] = E[\sum_{i=1}^n X_i] = \sum_{i=1}^n E[X_i] = 1$$

It is interesting to see the variance of X . Note that the formula for variance is $E[X^2] - E[X]^2$. Thus,

$$\begin{aligned}
X^2 &= (\sum_{i=1}^n X_i)^2 = \sum_{i=1}^n X_i^2 + \sum_{i=1}^n \sum_{j=1, j \neq i}^n X_i X_j \\
E[X^2] &= \sum_{i=1}^n E[X_i^2] + \sum_{i=1}^n \sum_{j=1, j \neq i}^n E[X_i X_j]
\end{aligned}$$

Note that X_i and X_j are not independent since after the first person has picked the hat, only $n - 1$ hats remain

$$\begin{aligned}
X_i X_j &= \begin{cases} 1 & \text{if } X_i = X_j = 1 \\ 0 & \text{otherwise} \end{cases} \\
P(X_i X_j = 1) &= P(X_i = 1)P(X_j = 1|X_i = 1) = \frac{1}{n} * \frac{1}{n-1} \\
E[X_i X_j] &= 1 * (\frac{1}{n} * \frac{1}{n-1}) + 0 * (1 - \frac{1}{n} * \frac{1}{n-1}) = \frac{1}{n(n-1)} \\
E[X_i^2] &= 1^2 \frac{1}{n} + 0^2 (1 - \frac{1}{n}) = \frac{1}{n}
\end{aligned}$$

Putting these values in the original equation for variance

$$\begin{aligned} E[X_2] &= n \frac{1}{n} + \frac{1}{n} \frac{1}{n-1} \left(\frac{n(n-1)}{2} * 2 \right) = 2 \\ Var(X) &= 2 - 1^2 = 1 \end{aligned}$$

11. [Question](#)

The following is the joint probability distribution of X and Y

$$f_{XY}(x, y) = f_X(x)f_{Y|X}(y|x) = \frac{1}{l} \frac{1}{x} = \frac{1}{xl} \quad \forall 0 \leq y \leq x \leq l$$

Using marginal probabilities, we can calculate $f_Y(y)$ and $E[Y]$ as

$$\begin{aligned} f_Y(y) &= \int f_{XY}(x, y) dx = \int_y^l \frac{1}{xl} dx = \frac{1}{l} \log \frac{l}{y} \quad \text{Note that for any } y, y \leq x \leq l \\ E[Y] &= \int y f_Y(y) dy = \int_0^l ly \frac{1}{l} \log \frac{l}{y} dy = \frac{l}{4} \end{aligned}$$

This problem can also be approached using iterated expectation

$$\begin{aligned} E[Y] &= E[E[Y|X]] = E[\text{uniform random variable between 0 and } x] \\ &= E\left[\frac{X}{2}\right] = \frac{1}{2}E[X] \\ &= \frac{l}{4} \end{aligned}$$

12. [Question](#)

Note that the required probability distribution is given by the following formula

$$p_{X+Y}(x) = \int_{-\inf}^{\inf} p_X(x-y) * p_Y(y) dy$$

However, note that the exponential distribution is not positive everywhere. For values < 0 , the probability density is 0. Hence, we break the integral into three parts as follows

$$p_{X+Y}(x) = \int_{-\inf}^0 p_X(x-y) * p_Y(y) dy + \int_0^x p_X(x-y) * p_Y(y) dy + \int_x^{\inf} p_X(x-y) * p_Y(y) dy$$

Carefully note that for y in range $(-\inf, 0]$, $p_Y(y) = 0$, and in the range $[x, \inf)$, $x-y < 0$, which implies $p_X(x-y) = 0$. Hence,

$$\begin{aligned} p_{X+Y}(x) &= \int_0^x p_X(x-y) * p_Y(y) dy \\ &= \lambda \mu \exp(-\lambda x) \int_0^x \exp((\lambda - \mu)y) dy \\ &= \frac{\lambda \mu}{\lambda - \mu} \exp(-\lambda x) (\exp((\lambda - \mu)x) - 1) \\ &= \frac{\lambda \mu}{\lambda - \mu} (\exp(\mu x) - \exp(-\lambda x)) \end{aligned}$$

13. [Question](#)

Assume that we break the stick at points X and Y . Assume $X < Y$. Then for the stick to form a triangle, the three lengths $X, Y - X$ and $1 - Y$ should satisfy the following three inequalities

$$\begin{aligned} X + (Y - X) &> 1 - Y \\ (Y - X) + (1 - Y) &> X \\ X + (1 - Y) &> Y - X \end{aligned}$$

which is nothing but the triangular region between the points $(0, 0.5)$, $(0.5, 0.5)$ and $(0.5, 1)$ and has the area of $1/8$. We should also consider the case $Y < X$ and by symmetry, the area is same. Now, X and Y comprise of the entire square region $X \leq 1$ and $Y \leq 1$. Hence the required probability is $2 * 1/8 = 1/4$.

14. [Question](#)

Always solve such questions using the cumulative distribution approach.

$$\begin{aligned} P(X \leq x) &= \begin{cases} 0 & x < 0 \\ \frac{1}{2}x & 0 \leq x \leq 2 \\ 1 & 2 < x \end{cases} \\ P(Y \leq y) &= P(X^3 \leq y) = P(X \leq y^{\frac{1}{3}}) \\ &= \begin{cases} 0 & y < 0 \\ \frac{1}{2}y^{\frac{1}{3}} & 0 \leq y^{\frac{1}{3}} \leq 2 \\ 1 & 2 < y^{\frac{1}{3}} \end{cases} \\ f_Y(y) &= \frac{dP(Y \leq y)}{dy}(y) \\ &= \begin{cases} 0 & y < 0 \\ \frac{1}{6}y^{-\frac{2}{3}} & 0 \leq y \leq 8 \\ 0 & 8 < y \end{cases} \end{aligned}$$

15. [Question](#)

Let X be the waiting time and $F_X(x)$ be the CDF. Then,

$$F_X(x) = \begin{cases} 0 & x < 0 \\ \frac{2}{3} & x = 0 \\ \frac{2}{3} + \frac{1}{30}x & 0 < x < 5 \\ 1 & 5 \leq x \end{cases}$$

The PDF is simply the derivate of the CDF. Thus, expectation is

$$E[X] = \frac{2}{3}(0) + \int_0^5 \frac{1}{30}x dx + \frac{1}{6}(5) = \frac{5}{4} \text{ mins}$$

16. [Question](#)

From Bayes' theorem

$$\begin{aligned} f_{Q|X}(q|x) &= \frac{f_{X|Q}(x|q)f_Q(q)}{f_X(x)} \\ &= \frac{f_{X|Q}(x|q)f_Q(q)}{\int_0^1 f_{X|Q}(x|q)f_Q(q) dq} \end{aligned}$$

We will need to solve separately for $x = 0$ and $x = 1$ as x is discrete.

$$f_{Q|X=0}(q|x=0) = \frac{(1-q) * 6q(1-q)}{\int_0^1 (1-q) * 6q(1-q) dq} = 12q(1-q)^2$$

$$f_{Q|X=1}(q|x=1) = \frac{q * 6q(1-q)}{\int_0^1 q * 6q(1-q) dq} = 12q^2(1-q)$$

17. [Question](#)

Questions of this type must only be approached through CDF. First find the CDF of Y and then it's PDF.

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(g(X) \leq y) \\ &= P(X \in [-y, 0] \cup X \in [0, y^2]) \\ &= (F_X(0) - F_X(-y)) + (F_X(y^2) - F_X(0)) \\ &= F_X(y^2) - F_X(-y) \\ p_Y(y) &= \frac{F_Y(y)}{dy} \\ &= \frac{dF_X(y^2)}{dx} \frac{d(y^2)}{dy} - \frac{dF_X(-y)}{dx} \frac{d(-y)}{dy} \\ &= 2yp_X(y^2) + p_X(-y) \\ &= 2y \frac{1}{\sqrt{2\pi}} e^{-\frac{y^4}{2}} + \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} \end{aligned}$$

18. [Question](#)

- (a) $P(X = k) = \binom{10}{k} 0.2^k 0.8^{10-k}$
- (b) $P(\text{no hits}) = 0.8^{10}$
- (c) $P(X \geq 6) = \sum_{k=6}^{10} \binom{10}{k} 0.2^k 0.8^{10-k}$
- (d) $E[X] = np = 2, \text{Var}(X) = np(1-p) = 1.6$ for Bernoulli distribution
- (e) $Y = 2X - 3, E[Y] = 2E[X] - 3 = 1, \text{Var}(Y) = 4\text{Var}(X) = 6.4$
- (f) $Z = X^2, E[Z] = E[X^2] = \text{Var}(X) + E[X]^2 = 5.6$

19. [Question](#)

For the mosquito, $P(\text{bite}) = P(\text{land})P(\text{bite}|\text{land}) = 0.1$. X is a geometric random variable. $E[X] = 1/p = 10$ and $\text{Var}(X) = \frac{1-p}{p^2} = 90$.

For the mosquito and tick combined, $P(\text{mosquito and tick}) = 0.1 + 0.1 * 0.7 - 0.1 * 0.1 * 0.7 = 0.163$. This is again a geometric random variable with $E[Y] = 1/0.163$ and $\text{Var}(Y) = (1 - 0.163)/(0.163^2)$.

20. [Question](#)

This quantity can be calculated using the law of total expectation

$$E[X] = E[X|A_1]P(A_1) + E[X|A_2]P(A_2) + \dots + E[X|A_n]P(A_n) \quad \text{where } A_i \text{ are disjoint}$$

Let H_1 denote heads at first toss, H_2 denote heads at the second toss, T_1 denote tails at first toss and T_2 denote tails at the second toss. Then,

$$\begin{aligned} E[X] &= E[X|H_1]P(H_1) + E[X|T_1]P(T_1) \\ E[X|H_1] &= E[X|H_1H_2]P(H_2|H_1) + E[X|H_1T_2]P(T_2|H_1) \\ &= 2p + (1 + E[X|T_1])(1 - p) \\ E[X|T_1] &= E[X|T_1T_2]P(T_2|T_1) + E[X|T_1H_2]P(H_2|T_1) \\ &= 2(1 - p) + (1 + E[X|H_1])p \end{aligned}$$

$E[X|H_1T_2] = 1 + E[X|T_1]$ because the tails after the first heads implies the first heads is now irrelevant and we have wasted one toss on the heads. The remaining process is same as starting from the first coin toss as tails.

Solving for the conditional expectations,

$$\begin{aligned} E[X|H_1] &= \frac{3 - 2p + p^2}{1 - p + p^2} \\ E[X|T_1] &= \frac{2 + p^2}{1 - p + p^2} \\ E[X] &= \frac{2 + p - p^2}{1 - p + p^2} \end{aligned}$$

21. Question

Define X as the following random variable

$$X = \begin{cases} 1, p = \frac{1}{4} & HHH \text{ or } TTT \\ 0, p = \frac{3}{4} & \text{otherwise} \end{cases}$$

- (a) K is simply a binomial distribution, where we want the 2^{nd} success to happen at the $K + 1$ th trial.

$$p_K(k) = \binom{k}{1} \frac{1}{4} \frac{3}{4}^{k-1} \quad \text{since the last trial is success}$$

- (b) M = number of tails before first success. Let the success be at $N + 1$. Defin Y as

$$\begin{aligned} Y &= \begin{cases} 1 & p = \frac{1}{2} \quad HHT, HTH, \text{ or } THH \\ 2 & p = \frac{1}{2} \quad HTT, THT, \text{ or } TTH \end{cases} \\ E[Y] &= 1 * \frac{1}{2} + 2 * \frac{1}{2} \\ Var(Y) &= (1 - \frac{3}{2})^2 * \frac{1}{2} + (2 - \frac{3}{2})^2 * \frac{1}{2} \\ E[N + 1] &= \frac{1}{p} = 4 \\ Var(N + 1) &= Var(N) = \frac{1 - p}{p^2} = \frac{1 - \frac{1}{4}}{\frac{1}{4}^2} \\ M &= Y_1 + Y_2 + \dots + Y_N \\ E[M] &= E[Y_1 + Y_2 + \dots + Y_N] \\ Var(M) &= Var(Y_1 + Y_2 + \dots + Y_N) \end{aligned}$$

Note that both Y and N are random variables here. Using the formulae for random number of random variables,

$$\begin{aligned} E[M] &= E[E[M|N]] = E[NE[Y]] = E[N]E[Y] = (4-1) * \frac{3}{2} = \frac{9}{2} \\ Var(M) &= Var(E[M|N]) + E[Var(M|N)] = Var(NE[Y]) + E[NVar(Y)] \\ &= E[Y]^2 Var(N) + E[N]Var(Y) = \frac{9}{4} * 12 + 3 * \frac{1}{4} = \frac{111}{4} \end{aligned}$$

22. Question

Let X be the number of tosses till the first coin is removed. This is a geometric random variable with $P(\text{success}) = \frac{1}{8}$. then $E[X] = 1/p = 8$. Now Y be the number of tosses till the second coin is removed (counting tosses after removal of first coin). Note that geometric random variables are memory less and what happened before the start of the "experiment" will not matter. Thus, $E[Y] = 1/(1/4) = 4$. Similarly, Z is the tosses till the last coin is removed and $E[Z] = 1/(1/2) = 2$. Note that the number of tosses till the end of experiment is simply $X + Y + Z$. $E[X + Y + Z] = E[X] + E[Y] + E[Z] = 14$.

23. Question

Note that the process till the end is a combination of multiple binomial process, such that any process lasts till the first success. Suppose we sign a paper and keep this in the drawer. Now the total signed papers in the drawer is k out of n and the $P(\text{success}) = \frac{n-k}{n}$ and $E[\text{draws till next unsigned paper}] = \frac{1}{p} = \frac{n}{n-k}$. Total draws

$$\begin{aligned} E &= \frac{n}{1} + \frac{n}{2} + \dots + \frac{n}{n} \\ &= n(1 + \frac{1}{2} + \dots + \frac{1}{n}) \\ \lim_{n \rightarrow \text{large}} E &= n \log(n) \end{aligned}$$

24. Question

A very straightforward way is to use a triple integral

$$P(X < Y < Z) = \int_0^{\text{inf}} \int_0^z \int_0^y \lambda e^{-\lambda x} \mu e^{-\mu y} \nu e^{-\nu z} dx dy dz = \frac{\lambda \mu}{(\lambda + \mu + \nu)(\mu + \nu)}$$

$P(X < Y < Z)$ can be broken down as $P(X < \min(Y, Z))P(Y < Z)$. Consider just $P(Y < Z)$

$$P(Y < Z) = \int_0^{\text{inf}} \int_0^z \mu e^{-\mu y} \nu e^{-\nu z} dy dz = \frac{\mu}{\mu + \nu}$$

Thus, when two exponential processes are considered, probability of arrival of 1st before 2nd is simply the percentage ratio of parameters. Thus,

$$\begin{aligned} P(X < \min(Y, Z)) &= \frac{\lambda}{\lambda + (\mu + \nu)} \quad Y \text{ and } Z \text{ can be combined as a single process} \\ P(Y < Z) &= \frac{\mu}{\mu + \nu} \\ P(X < Y < Z) &= P(X < \min(Y, Z))P(Y < Z) \\ &= \frac{\lambda \mu}{(\lambda + \mu + \nu)(\mu + \nu)} \end{aligned}$$

25. **Question** We can model the arrival process like a Poisson process. $\lambda = 5$ and $\tau = \frac{1}{2}$

$$P(\lambda, \tau, k) = \frac{(\lambda\tau)^k e^{-\lambda\tau}}{k!}$$

$$P(5, \frac{1}{2}, 0) = \frac{(5 * \frac{1}{2})^0 e^{-5 * \frac{1}{2}}}{0!}$$

$$P(5, \frac{1}{2}, 1) = \frac{(5 * \frac{1}{2})^1 e^{-5 * \frac{1}{2}}}{1!}$$

26. **Question**

- $P(\text{fish for } > 2 \text{ hours}) = P(k = 0, \tau = 2) = e^{-0.6 * 2}$
- $P(\text{fish for } > 2 \text{ but } < 5 \text{ hours}) = P(\text{first catch in } [2, 5] \text{ hours}) = P(k = 0, \tau = 2)(1 - P(k = 0, \tau = 3))$ which is no fish in $[0, 2]$ but at least 1 fish in the next 3 hours (which will be independent of first 2 hours)
- $P(\text{catch at least two fish}) = P(\text{at least 2 catches before 2 hours}) = 1 - P(k = 0, \tau = 2) - P(k = 1, \tau = 2)$
- $E[\text{fish}]$ has two possibilities, either single fish after 2 hours, or many fish before 2 hours. $E[\text{fish}] = E[\text{fish} | \tau \leq 2](1 - P(\tau > 2)) + E[\text{fish} | \tau > 2]P(\tau > 2) = (0.6 * 2) * (1 - P(k = 0, \tau = 2)) + 1 * P(k = 0, \tau = 2)$
- $E[\text{Total fishing time}] = 2 + P(k = 0, \tau = 2)\frac{1}{\lambda}$, since we fish for atleast 2 hours
- $E[\text{future fishing time—fished for two hours}]$ can be obtained using the memoryless property of Poisson process. The expected time till first arrival is independent of what has happened till now. Thus, $E[T_1] = \frac{1}{\lambda}$

27. **Question**

Start with the merged Poisson process which will denote the time till the first bulb will fail. For this process, $\lambda' = 3\lambda$. Hence, $E[\text{first bulb fails}] = \frac{1}{3\lambda}$. After the first bulb dies out, we are left with a process with $\lambda' = 3\lambda$. Due to memoryless property, $E[\text{second bulb fails}] = \frac{1}{2\lambda}$ and consequently $E[\text{last bulb fails}] = \frac{1}{\lambda}$. Note the above two times denote the time difference, i.e. the time taken for the bulb to die out after the last bulb died out. Thus, $E[\text{time until last bulb dies out}] = \frac{1}{3\lambda} + \frac{1}{2\lambda} + \frac{1}{\lambda}$

28. **Question**

(a)

$$E[\text{time till failure}] = E[\text{time till failure}|A]P(A) + E[\text{time till failure}|B]P(B)$$

$$= \frac{1}{\lambda_A} \frac{1}{2} + \frac{1}{\lambda_B} \frac{1}{2}$$

$$= \frac{1}{2} \left(1 + \frac{1}{3}\right) = \frac{2}{3}$$

- (b) Let C denote the event of no failure till time t . $P(C)$ for a given λ will be $\int_t^{\infty} \lambda e^{-\lambda t}$. Then,

$$P(C) = P(C|A)P(A) + P(C|B)P(B) \quad \text{Using total probability theorem}$$

$$= e^{-t} \left(\frac{1}{2}\right) + e^{-3t} \left(\frac{1}{2}\right)$$

$$= \frac{1}{2} (e^{-t} + e^{-3t})$$

(c) Let C denote the event of no failure till time t . Then,

$$\begin{aligned}
 P(A|C) &= \frac{P(C|A)P(A)}{P(C)} \\
 &= \frac{P(C|A)P(A)}{P(C|A)P(A) + P(C|B)P(B)} \\
 &= \frac{\frac{1}{2}e^{-t}}{\frac{1}{2}(e^{-t} + e^{-3t})} \\
 &= \frac{1}{1 + e^{-2t}}
 \end{aligned}$$

(d) Let T_{B1}, T_{B2} and T_A denote the life times of the first B bulb, second B bulb and the A bulb respectively. First consider the solution to $P(T_{B1} + T_{B2} = t)$

$$\begin{aligned}
 P(T_{B1} + T_{B2} = t) &= \int_0^t P(T_{B1} = t_1)P(T_{B2} = t - t_1)dt_1 \quad \text{Using independence} \\
 &= \int_0^t 3e^{-3t_1}3e^{-3(t-t_1)}dt_1 \\
 &= \int_0^t 9e^{-3t}dt_1 \\
 &= 9te^{-3t}
 \end{aligned}$$

Now, we can rewrite the required probability in a slightly different format

$$\begin{aligned}
 P(T_{B1} + T_{B2} > T_A) &= P(T_{B1} + T_{B2} = t)P(T_A \leq t) \\
 &= \int_0^{\inf} 9te^{-3t} \left(\int_0^t e^{-t_1} dt_1 \right) dt \\
 &= \int_0^{\inf} 9te^{-3t}(1 - e^{-t})dt \\
 &= \int_0^{\inf} 9te^{-3t} - 9te^{-4t} dt
 \end{aligned}$$

Using integration by parts, $\int uv' = uv - \int u'v$ and choosing $u = t, v = e^{-3t}/3$,

$$\begin{aligned}
 P(T_{B1} + T_{B2} > T_A) &= \left[9[te^{-3t}]_0^{\inf} - 3 \int_0^{\inf} e^{-3t} dt - 9[te^{-4t}]_0^{\inf} + \frac{9}{4} \int_0^{\inf} e^{-4t} dt \right] \\
 &= 0 + 1 - 0 - \frac{9}{16} = \frac{7}{16}
 \end{aligned}$$

(e) Let there be N bulbs of type B out of the 12 bulbs. Clearly N is a random variable and can be seen as the "successes" of choosing a given bulb as B. and the probability of choosing any i th bulb as B is $1/2$.

Let the life time of any bulb of type B be T . Then the total lifetime of all the type B bulbs will be NT , which is nothing but the sum of a random number of random variables.

$$\begin{aligned}
 E[NT] &= E[N]E[T] = np * \frac{1}{\lambda} = 12 * \frac{1}{2} * \frac{1}{3} = 2 \\
 Var(NT) &= E[Var(NT|N)] + Var(E[NT|N]) = E[N]Var(T) + E[T]^2Var(N) \\
 &= np * \frac{1}{\lambda^2} + \left(\frac{1}{\lambda}\right)^2 np(1 - p) = 1
 \end{aligned}$$

(f) Let D be the event that the lifetime is greater than t or $T > t$. Then,

$$\begin{aligned}
E[T|D] &= E[T|D, A]P(A|D) + E[T|D, B]P(B|D) \\
&= t + (E[T - t|D, A]P(A|D) + E[T - t|D, B]P(B|D)) \\
&= t + \left(\frac{1}{1}P(A|D) + \frac{1}{3}P(B|D)\right) && \text{Using memoryless property} \\
&= t + \left(\frac{1}{1+e^{-2t}} + \frac{1}{3}\left(1 - \frac{1}{1+e^{-2t}}\right)\right) && \text{Using part 28c} \\
&= t + \frac{1}{3} + \frac{2}{3} \frac{1}{1+e^{-2t}}
\end{aligned}$$

29. [Question](#)

Using balance equations, we have

$$\begin{aligned}
\pi_1 &= \pi_1 p_{11} + \pi_2 p_{21} \\
\pi_2 &= \pi_1 p_{12} + \pi_2 p_{22} \\
\pi_1 + \pi_2 &= 1
\end{aligned}$$

Solving, $\pi_1 = \frac{2}{7}$ and $\pi_2 = \frac{5}{7}$

30. [Question](#)

Let a_i denote the absorption probabilities into state 4 starting from i

$$\begin{aligned}
a_5 &= 0, a_4 = 1 \\
a_i &= \sum_j a_j p_{ij} \\
a_2 &= a_1 p_{21} + a_4 p_{24} \\
a_3 &= a_1 p_{31} + a_2 p_{32} + a_5 p_{35} \\
a_1 &= a_2 p_{12} + a_3 p_{13}
\end{aligned}$$

Solving, $a_1 = \frac{9}{14}$, $a_2 = \frac{5}{7}$ and $a_3 = \frac{15}{28}$

Let μ_i denote the expected time till absorption starting from i , then

$$\begin{aligned}
\mu_4 &= 0 \\
\mu_1 &= 1 + \mu_2 p_{12} + \mu_3 p_{13} \\
\mu_2 &= 1 + \mu_1 p_{21} + \mu_4 p_{24} \\
\mu_3 &= 1 + \mu_1 p_{31} + \mu_2 p_{32}
\end{aligned}$$

Solving, $\mu_1 = \frac{55}{4}$, $\mu_2 = 12$ and $\mu_3 = \frac{111}{8}$

31. [Question](#)

- (a) The probability of eventually leaving course 6 is 1 as states 15 and 9 are absorbing states.
- (b) Here we have to calculate the probability of absorption into state 15. Let a_i denote the probability of absorption into state 15 from state i . Then, $a_{15} = 1$ and $a_9 = 0$. Using equations from 20.4,

$$\begin{aligned}
a_{6-1} &= \frac{1}{2}a_{6-1} + \frac{1}{8}a_{6-2} + \frac{1}{8}a_{6-3} + \frac{1}{8}a_9 + \frac{1}{8}a_{15} \\
a_{6-2} &= \frac{1}{2}a_{15} + \frac{3}{8}a_{6-1} + \frac{1}{8}a_{6-3} \\
a_{6-3} &= \frac{1}{4}a_9 + \frac{3}{8}a_{6-1} + \frac{3}{8}a_{6-2}
\end{aligned}$$

Solving the 3 equations, 3 variable system, $a_{6-1} = 105/184$, $a_{6-2} = 143/184$ and $a_{6-3} = 93/184$.

- (c) Let μ_i denote the expected number of steps to get absorbed starting from state i . Then, $\mu_{15} = \mu_9 = 0$. Using equations from 20.4,

$$\begin{aligned}\mu_{6-1} &= 1 + \frac{1}{2}\mu_{6-1} + \frac{1}{8}\mu_{6-2} + \frac{1}{8}\mu_{6-3} + \frac{1}{8}\mu_9 + \frac{1}{8}\mu_{15} \\ \mu_{6-2} &= 1 + \frac{1}{2}\mu_{15} + \frac{3}{8}\mu_{6-1} + \frac{1}{8}\mu_{6-3} \\ \mu_{6-3} &= 1 + \frac{1}{4}\mu_9 + \frac{3}{8}\mu_{6-1} + \frac{3}{8}\mu_{6-2}\end{aligned}$$

Solving, $\mu_{6-1} = 81/23$, $\mu_{6-2} = 63/23$ and $\mu_{6-3} = 77/23$.

- (d) This question can be done in a manner similar to the equations described above but with a small adjustment. Note that, we can either have 0, 1, or 2 ice creams. Consider $v_i(j)$ as the probability of making j additional ice creams from 6-2 to 6-1 or 6-3 to 6-1 transitions, given the current state is i . Note $v_{15}(0) = v_9(0) = 1$. Then,

$$\begin{aligned}v_{6-1}(0) &= \frac{1}{2}v_{6-1}(0) + \frac{1}{8}v_{6-2}(0) + \frac{1}{8}v_{6-3}(0) + \frac{1}{8}v_9(0) + \frac{1}{8}v_{15}(0) \\ v_{6-2}(0) &= \frac{1}{2}v_{15}(0) + \frac{3}{8}v_{6-1}(0) + \frac{1}{8}v_{6-3}(0) \\ v_{6-3}(0) &= \frac{1}{4}v_9(0) + \frac{3}{8}v_{6-1}(0) + \frac{3}{8}v_{6-2}(0)\end{aligned}$$

Some of the transitions have been directly replaced with 0 as we are considering 0 ice creams and thus those transitions are not possible (6-2 to 6-1 for instance). Solving, $v_{6-1}(0) = 46/61$, $v_{6-2}(0) = 34/61$ and $v_{6-3}(0) = 28/61$.

The same way, we can construct equations for 1 additional steps where $v_{15}(1) = v_9 = 0$.

$$\begin{aligned}v_{6-1}(1) &= \frac{1}{2}v_{6-1}(1) + \frac{1}{8}v_{6-2}(1) + \frac{1}{8}v_{6-3}(1) + \frac{1}{8}v_9(1) + \frac{1}{8}v_{15}(1) \\ v_{6-2}(1) &= \frac{1}{2}v_{15}(1) + \frac{3}{8}v_{6-1}(0) + \frac{1}{8}v_{6-3}(1) \\ v_{6-3}(1) &= \frac{1}{4}v_9(1) + \frac{3}{8}v_{6-1}(0) + \frac{3}{8}v_{6-2}(1)\end{aligned}$$

In the second equation, after going from 6-2 to 6-1, we can only get 0 more ice creams. Hence, some of the values have been replaced with the $v_i(0)$ calculated above. Solving, $v_{6-1}(1) = 690/3721$, $v_{6-2}(1) = 1242/3721$ and $v_{6-3}(1) = 1518/3721$.

Note that since the total ice creams are 0, 1, or 2, we have $v_{6-1}(0) + v_{6-1}(1) + v_{6-1}(2) = 1$. $E[\text{ice creams}] = 0 * v_{6-1}(0) + 1 * v_{6-1}(1) + 2 * v_{6-1}(2) = 1140/3721$

- (e) We need to recalculate the the transition probabilities since we are conditioning on

the event A that we land up in state 15.

$$\begin{aligned}
P_{ij|A} &= P(X_{n+1} = j | X_i = i, A) \\
&= \frac{P(X_{n+1} = j, X_n = i, A)}{P(X_n = i, A)} \\
&= \frac{P(A | X_{n+1} = j, X_n = i) P(X_{n+1} = j | X_n = i) P(X_n = i)}{P(A | X_n = i) P(X_n = i)} \\
&= \frac{P(A | X_{n+1} = j) P(X_{n+1} = j | X_n = i)}{P(A | X_n = i)} \\
&= \frac{a_j}{a_i} P_{ij}
\end{aligned}$$

where a_i is the probability of absorption into state 15 starting from state i . Since markov process is only dependent on the last state, absorption probabilities are not dependent on n .

We can write equations similar to 20.4 for calculating the expected number of steps with the adjusted transition probabilities

$$\begin{aligned}
\mu_{6-1} &= 1 + \frac{a_{6-1}}{a_{6-1}} \frac{1}{2} \mu_{6-1} + \frac{a_{6-2}}{a_{6-1}} \frac{1}{8} \mu_{6-2} + \frac{a_{6-3}}{a_{6-1}} \frac{1}{8} \mu_{6-3} + \frac{a_{15}}{a_{6-1}} \frac{1}{8} \mu_{15} + \frac{a_9}{a_{6-1}} \frac{1}{8} \mu_9 \\
\mu_{6-2} &= 1 + \frac{a_{6-1}}{a_{6-2}} \frac{3}{8} \mu_{6-1} + \frac{a_{6-3}}{a_{6-2}} \frac{1}{8} \mu_{6-3} + \frac{a_{15}}{a_{6-2}} \frac{1}{2} \mu_{15} \\
\mu_{6-3} &= 1 + \frac{a_{6-1}}{a_{6-3}} \frac{3}{8} \mu_{6-1} + \frac{a_{6-2}}{a_{6-3}} \frac{3}{8} \mu_{6-2} + \frac{a_9}{a_{6-3}} \frac{1}{4} \mu_9
\end{aligned}$$

where $\mu_{15} = \mu_9 = 0$, $a_{15} = 1$, and $a_9 = 0$. The absorption probabilities can be taken from the part 31b. Solving, $\mu_{6-1} = 1763/483$.

- (f) The changed probabilities become $P(X_{n+1} = 15 | X_n = 6-1) = P(X_{n+1} = 6-2 | X_n = 6-1) = P(X_{n+1} = 6-3 | X_n = 6-1) = 1/6$, $P(X_{n+1} = 6-1 | X_n = 6-2) = 3/4$ and $P(X_{n+1} = 6-3 | X_n = 6-2) = 1/4$. We then use equations from 20.4 to calculate the expected values

$$\begin{aligned}
\mu_{6-1} &= 1 + \frac{1}{2} \mu_{6-1} + \frac{1}{6} \mu_{6-2} + \frac{1}{6} \mu_{6-3} + \frac{1}{6} \mu_9 \\
\mu_{6-2} &= 1 + \frac{3}{4} \mu_{6-1} + \frac{1}{4} \mu_{6-3} \\
\mu_{6-3} &= 1 + \frac{1}{4} \mu_9 + \frac{3}{8} \mu_{6-1} + \frac{3}{8} \mu_{6-2}
\end{aligned}$$

where $\mu_{15} = 0$. Solving, $\mu_{6-1} = 86/13$, $\mu_{6-2} = 98/13$ and $\mu_{6-3} = 82/13$.

- (g) If we look carefully at the new probabilities, states 15 and 9 become recurrent. Far into the future, we are sure to land up in those states, and will be in either one of those. By symmetry, the two should be same. $\pi_{15} = \pi_9 = 1/2$.
- (h) We assume that 6-1 is an absorbing state, and accordingly calculate the probabilities. Note that there will not be an equation for 6-1 since we are then already in the final

state.

$$\begin{aligned}\mu_{6-2} &= 1 + \frac{1}{8}\mu_{6-3} + \frac{1}{2}\mu_{15} \\ \mu_{6-3} &= 1 + \frac{3}{8}\mu_{6-2} + \frac{1}{4}\mu_9 \\ \mu_9 &= 1 + \frac{7}{8}\mu_9 \\ \mu_{15} &= 1 + \frac{7}{8}\mu_{15}\end{aligned}$$

Solving, $\mu_{6-2} = 344/61$, $\mu_{6-3} = 312/61$ and $\mu_9 = \mu_{15} = 8$. Plugging these into the following equation (which corresponds to taking one step out of 6-1),

$$\mu_{6-1} = 1 + \frac{1}{2}\mu_{6-1} + \frac{1}{8}\mu_{15} + \frac{1}{8}\mu_{6-2} + \frac{1}{8}\mu_{6-3} + \frac{1}{8}\mu_{15} = \frac{265}{61}$$

32. **Question** The exact answer will be

$$\sum_{k=0}^{21} \binom{36}{k} \left(\frac{1}{2}\right)^{36} = 0.8785$$

But the same can be estimated using the CLT as follows

$$\begin{aligned}\mu &= np = 18 \\ \sigma^2 &= np(1-p) = 9 \\ P(S_n \leq 21) &\approx P\left(\frac{S_n - 18}{3} \leq \frac{21 - 18}{3}\right) \approx 0.843\end{aligned}$$

Our estimate is in the rough range of the answer but not quite close. We can do better using the $\frac{1}{2}$ correction

$$\begin{aligned}P(S_n \leq 21) &= P(S_n < 22) \text{ since } S_n \text{ is an integer} \\ \text{Consider } P(S_n \leq 21.5) &\text{ as a compromise between the two} \\ P(S_n \leq 21.5) &= P\left(\frac{S_n - 18}{3} \leq \frac{21.5 - 18}{3}\right) \approx 0.879\end{aligned}$$

In a similar manner, $P(S_n = 19) = P(18.5 \leq S_n \leq 19.5)$ using $\frac{1}{2}$ correction.

33. **Question** We know that for a normal population,

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$$

In the given problem, $n = 15$ and $\sigma^2 = 9$. Hence,

$$\begin{aligned}P(S^2 > 12) &= P\left(\frac{14S^2}{9} > \frac{14}{9}12\right) \\ &= P\left(\chi_{14}^2 > \frac{56}{3}\right) = 0.178 \text{ from standard tables or chi-square calculators}\end{aligned}$$

34. Question

Since the observations are independent, the likelihood of all the observations under some θ is given by

$$p_{X|\Theta}(x|\theta) = \prod_{i=1}^n \theta \exp(-\theta x_i)$$


$$\log(p_{X|\Theta}(x|\theta)) = n \log(\theta) - \theta \left(\sum_{i=1}^n x_i \right)$$

Taking the derivative and maximizing with respect to θ , $\hat{\theta}_{MLE} = \frac{n}{\sum_{i=1}^n x_i}$

35. Question

We need to evaluate $f_{\Theta|X}(\theta|x)$ in order to get $E[\Theta|X]$.

$f_{X,\Theta}(x, \theta) = f_X(x)f_{\Theta|X}(\theta|x)$ which is a parallelogram on the $\theta - x$ plane at the points (3, 4), (5, 4), (9, 10) and (11, 10). Then $E[\Theta|X]$ can be obtained by drawing vertical lines on the planes and calculating the $E[\theta]$ over that line. It is a line which bends at two points.



36. Question

(a) No, since X_i is also uniform in $[-1, 1]$

(b) Yes, $E[Y_i] = 0$ by symmetry. For $\epsilon > 0$,

$$\begin{aligned} \lim_{i \rightarrow \infty} P(|Y_i - \mu_i| > \epsilon) &= \lim_{i \rightarrow \infty} P\left(\left|\frac{X_i}{i} - 0\right| > \epsilon\right) \\ &= \lim_{i \rightarrow \infty} P\left(\frac{X_i}{i} > \epsilon \text{ and } \frac{X_i}{i} < -\epsilon\right) \\ &= \lim_{i \rightarrow \infty} [P(X_i > i\epsilon) + P(X_i < -i\epsilon)] = 0 \end{aligned}$$

(c) Yes, $E[Y_i] = 0$ by symmetry. For $\epsilon > 0$,

$$\begin{aligned} \lim_{i \rightarrow \infty} P(|Z_i - 0| > \epsilon) &= \lim_{i \rightarrow \infty} P((X_i)^i > \epsilon \text{ or } (X_i)^i < -\epsilon) \\ &= \lim_{i \rightarrow \infty} \left[\frac{1}{2}(1 - \epsilon^{1/i}) + \frac{1}{2}(1 - \epsilon^{1/i}) \right] \\ &= \lim_{i \rightarrow \infty} (1 - \epsilon^{1/i}) = 0 \end{aligned}$$