

Additions

Kevin O'Brien

November 9, 2014

Contents

0.0.1	Influence Analysis for LME Models	1
0.0.2	Influence Statistics for LME models	1
0.0.3	Likelihood and estimation	2
0.1	Likelihood ratio tests	4
0.1.1	Test for inter-method bias	6
0.1.2	Testing Procedures	7

Influence Analysis for LME Models

The linear mixed effects model is a useful methodology for fitting a wide range of models. However, linear mixed effects models are known to be sensitive to outliers. ? advises that identification of outliers is necessary before conclusions may be drawn from the fitted model.

Standard statistical packages concentrate on calculating and testing parameter estimates without considering the diagnostics of the model. The assessment of the effects of perturbations in data, on the outcome of the analysis, is known as statistical influence analysis. Influence analysis examines the robustness of the model. Influence analysis methodologies have been used extensively in classical linear models, and provided the basis for methodologies for use with LME models. Computationally inexpensive diagnostics tools have been developed to examine the issue of influence (?). Studentized residuals, error contrast matrices and the inverse of the response variance covariance matrix are regular components of these tools.

Influence Statistics for LME models

Influence statistics can be coarsely grouped by the aspect of estimation that is their primary target:

- overall measures compare changes in objective functions: (restricted) likelihood distance (Cook and Weisberg 1982, Ch. 5.2)
- influence on parameter estimates: Cook's (Cook 1977, 1979), MDFFITS (Belsley, Kuh, and Welsch 1980, p. 32)
- influence on precision of estimates: CovRatio and CovTrace

- influence on fitted and predicted values: PRESS residual, PRESS statistic (Allen 1974), DFFITS (Belsley, Kuh, and Welsch 1980, p. 15)
- outlier properties: internally and externally studentized residuals, leverage
- LMEs
- Likelihood and log likelihood functions
- Likelihood ratio test
- more on score functions etc
- MLEs
- Algorithms

The estimate for the fixed effects are referred to as the best linear unbiased estimates (BLUE). Henderson's estimate for the random effects is known as the best linear unbiased predictor (BLUP).

Likelihood and estimation

Likelihood is the hypothetical probability that an event that has already occurred would yield a specific outcome. Likelihood differs from probability in that probability refers to future occurrences, while likelihood refers to past known outcomes.

The likelihood function is a fundamental concept in statistical inference. It indicates how likely a particular population is to produce an observed sample. The set of values that maximize the likelihood function are considered to be optimal, and are used as the estimates of the parameters.

Maximum likelihood (ML) estimation is a method of obtaining parameter estimates by optimizing the likelihood function. The likelihood function is constructed as a function of the parameters in the specified model.

Restricted maximum likelihood (REML) is an alternative methods of computing parameter estimated. REML is often preferred to ML because it produces unbiased estimates of covariance parameters by taking into account the loss of degrees of freedom that results from estimating the fixed effects in β .

REML estimation reduces the bias in the variance component, and also handles high correlations more effectively, and is less sensitive to outliers than ML. The problem with REML for model building is that the "likelihoods" obtained for different fixed effects are not comparable. Hence it is not valid to compare models with different fixed effects using a likelihood ratio test or AIC when REML is used to estimate the model. Therefore models derived using ML must be used instead.

Assuming a statistical model $f_\theta(y)$ parameterized by a fixed and unknown set of parameters θ , the likelihood $L(\theta)$ is the probability of the observed data y considered as a function of θ (?).

The log likelihood $l(\theta)$

0.1 Likelihood ratio tests

Likelihood ratio tests are a class of tests based on the comparison of the values of the likelihood functions of two candidate models. LRTs can be used to test hypotheses about covariance parameters or fixed effects parameters in the context of LMEs.

The test statistic for the LRT is the difference of the log-likelihood functions, multiplied by -2 . The probability distribution of the test statistic is approximated by the χ^2 distribution with $(\nu_1 - \nu_2)$ degrees of freedom, where ν_1 and ν_2 are the degrees of freedom of models 1 and 2 respectively.

The score function $S(\theta)$ is the derivative of the log likelihood with respect to θ ,

$$S(\theta) = \frac{\partial}{\partial \theta} l(\theta),$$

and the maximum likelihood estimate is the solution to the score equation

$$S(\theta) = 0.$$

The Fisher information $I(\theta)$, which is defined as

$$I(\theta) = -\frac{\partial^2}{\partial \theta^2} l(\theta),$$

give rise to the observed Fisher information ($I(\hat{\theta})$) and the expected Fisher information ($\mathcal{I}(\theta)$).

? used ML estimation to estimate the true correlation between the variables when the measurements are linked over time. The methodology relies on the assumption that the two variables with repeated measures follow a multivariate normal distribution. The methodology currently does not extend to any more than two cases. The MLE of the correlation takes into account the dependency among repeated measures.

The true correlation ρ_{xy} is repeated measurements can be considered as having two components: between subject and within-subject correlation. The usefulness of estimating repeated measure correlation coefficients is the calculation of between-method and within-method variabilities are produced as by-products.

0.1.1 Test for inter-method bias

Bias is determinable by examination of the 't-table'. Estimate for both methods are given, and the bias is simply the difference between the two. Because the *R* implementation does not account for an intercept term, a p -value is not given. Should a p -value be required specifically for the bias, and simple restructuring of the model is required wherein an intercept term is included. Output from a second implementation will yield a p -value.

0.1.2 Testing Procedures

Roy's methodology requires the construction of four candidate models. The first candidate model is compared to each of the three other models successively. It is the alternative model in each of the three tests, with the other three models acting as the respective null models.

The probability distribution of the test statistic can be approximated by a chi-square distribution with $(\nu_1 - \nu_2)$ degrees of freedom, where ν_1 and ν_2 are the degrees of freedom of models 1 and 2 respectively.

Likelihood ratio tests are very simple to implement in **R**, simply use the 'anova()' commands. Sample output will be given for each variability test. The likelihood ratio test is the procedure used to compare the fit of two models. For each candidate model, the '-2 log likelihood' ($M2LL$) is computed. The test statistic for each of the three hypothesis tests is the difference of the $M2LL$ for each pair of models. If the p -value in each of the respective tests exceed as significance level chosen by the analyst, then the null model must be rejected.

$$-2 \ln \Lambda_d = [M2LL \text{ under } H_0 \text{ model}] - [M2LL \text{ under } H_A \text{ model}] \quad (1)$$

These test statistics follow a chi-square distribution with the degrees of freedom computed as the difference of the LRT degrees of freedom.

$$\nu = [\text{LRT df under } H_0 \text{ model}] - [\text{LRT df under } H_A \text{ model}] \quad (2)$$


```

> anova(MCS1,MCS2)
>
>

```

	Model	df	AIC	BIC	logLik	Test	L.Ratio	p-value
MCS1	1	8	4077.5	4111.3	-2030.7			
MCS2	2	7	4075.6	4105.3	-2030.8	1 vs 2	0.15291	0.6958