

Chapter 1

Limits of Agreement

1.1 Computing LoAs from LME models

One important feature of replicate observations is that they should be independent of each other. In essence, this is achieved by ensuring that the observer makes each measurement independent of knowledge of the previous value(s). This may be difficult to achieve in practice.

1.2 New

Computing limits of agreement features prominently in many method comparison studies, further to Bland and Altman (1986, 1999). Bland and Altman (1999) addresses the issue of computing LoAs in the presence of replicate measurements, suggesting several computationally simple approaches. When repeated measures data are available, it is desirable to use all the data to compare the two methods. However, the original Bland-Altman method was developed for two sets of measurements done on one occasion (i.e. independent data), and so this approach is not suitable for replicate measures data. However, as a naive analysis, it may be used to explore the data because of the simplicity of the method. Carstensen et al. (2008) computes the limits of agreement to the case with replicate measurements by using LME models.

Roy (2009) formulates a very powerful method of assessing whether two methods of measurement, with replicate measurements, also using LME models. Roy's approach is based on the construction of variance-covariance matrices. Importantly, Roy's approach does not address the issue of limits of agreement (though another related analysis, the coefficient of repeatability, is mentioned).

This paper seeks to use Roy's approach to estimate the limits of agreement. These estimates will be compared to estimates computed under Carstensen's formulation.

In computing limits of agreement, it is first necessary to have an estimate for the standard deviations of the differences. When the agreement of two methods is analyzed using LME models, a clear method of how to compute the standard deviation is required. As the estimate for inter-method bias and the quantile would be the same for both methodologies, the focus is solely on the standard deviation.

1.2.1 Roy's method

Roy proposes a novel method using the LME model with Kronecker product covariance structure in a doubly multivariate set-up to assess the agreement between a new method and an established method with unbalanced data and with unequal replications for different subjects (Roy, 2009).

Using Roy's method, four candidate models are constructed, each differing by constraints applied to the variance covariance matrices. In addition to computing the inter-method bias, three significance tests are carried out on the respective formulations to make a judgement on whether or not two methods are in agreement.

1.2.2 Carstensen's Model

MORE

1.2.3 Carstensen's LOAs

Carstensen presents a model where the variation between items for method m is captured by σ_m and the within item variation by τ_m .

Further to his model, Carstensen computes the limits of agreement as

$$\hat{\alpha}_1 - \hat{\alpha}_2 \pm \sqrt{2\hat{\tau}^2 + \hat{\sigma}_1^2 + \hat{\sigma}_2^2}$$

1.2.4 Roy's LOAs

The limits of agreement computed by Roy's method are derived from the variance covariance matrix for overall variability. This matrix is the sum of the between subject VC matrix and the within-subject VC matrix.

The standard deviation of the differences of methods x and y is computed using values from the overall VC matrix.

$$\text{var}(x - y) = \text{var}(x) + \text{var}(y) - 2\text{cov}(x, y)$$

The respective estimates computed by both methods are tabulated as follows. Evidently there is close correspondence between both sets of estimates.

Carstensen et al. (2008) formulates an LME model, both in the absence and the presence of an interaction term. ? uses both to demonstrate the importance of using an interaction term. Failure to take the replication structure into account results in over-estimation of the limits of agreement. For the Carstensen estimates below, an interaction term was included when computed.

Roy (2006) uses the “Blood” data set, which featured in Bland and Altman (1999).

Carstensen et al. (2008) describes the sampling method when discussing of a motivating example

Diabetes patients attending an outpatient clinic in Denmark have their HbA_{1c} levels routinely measured at every visit. Venous and Capillary blood samples were obtained from all patients appearing at the clinic over two days. Samples were measured on four consecutive days on each machines, hence there are five analysis days.

Carstensen et al. (2008) notes that every machine was calibrated every day to the manufacturers guidelines. Measurements are classified by method, individual and replicate. In this case the replicates are clearly not exchangeable, neither within patients nor simulataneously for all patients.

1.3 Hamlett

Hamlett re-analyses the data of Lam et al. (1999) to generalize their model to cover other settings not covered by the Lam method.

In many cases, repeated observation are collected from each subject in sequence and/or longitudinally.

$$y_i = \alpha + \mu_i + \epsilon$$

1.3.1 T

he classical model is based on measurements y_{mi} by method $m = 1, 2$ on item $i = 1, 2 \dots$

$$y_{mi} = \alpha_m + \mu_i + e_{mi}$$

$$e_{mi} \sim \mathcal{N}(0, \sigma_m^2)$$

Even though the separate variances can not be identified, their sum can be estimated by the empirical variance of the differences.

Like wise the separate α can not be estimated, only their difference can be estimated as \bar{D}

Bibliography

- Bland, J. and D. Altman (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *The Lancet* *i*, 307–310.
- Bland, J. and D. Altman (1999). Measuring agreement in method comparison studies. *Statistical Methods in Medical Research* *8*(2), 135–160.
- Carstensen, B., J. Simpson, and L. C. Gurrin (2008). Statistical models for assessing agreement in method comparison studies with replicate measurements. *The International Journal of Biostatistics* *4*(1).
- Lam, M., K. Webb, and D. O’Donnell (1999). Correlation between two variables in repeated measurements. *American Statistical Association, Proceedings of the Biometric Session*, 213–218.
- Roy, A. (2006). Estimating correlation coefficient between two variables with repeated observations using mixed effects models. *Biometric Journal* *2*, 286–301.
- Roy, A. (2009). An application of linear mixed effects model to assess the agreement between two methods with replicated observations. *Journal of Biopharmaceutical Statistics* *19*, 150–173.