

1 Model Validation Framework

In statistical modelling, the process of model validation is a critical step of model fitting process, but also a step that is too often overlooked. A very simple procedure is to examine commonly-used metrics, such as the R^2 value. However, using a small handful of simple measures and methods is insufficient to properly assess the quality of a fitted model. To do so properly, a full and comprehensive analysis that tests all of the assumptions, as far as possible, must be carried out.

A statistical model, whether of the fixed-effects or mixed-effects variety, represents how you think your data were generated. Following model specification and estimation, it is of interest to explore the model-data agreement by raising questions such as

- Does the model-data agreement support the model assumptions?
- Should model components be refined, and if so, which components? For example, should regressors be added or removed, and is the covariation of the observations modeled properly?
- Are the results sensitive to model and/or data? Are individual data points or groups of cases particularly influential on the analysis?

? describes the model validation framework as comprised of the following tasks

- overall measures of goodness-of-fit
- the informal, graphical examination of estimates of model errors to assess the quality of distributional assumptions: residual analysis
- the quantitative assessment of the inter-relationship of model components; for example, collinearity diagnostics
- the qualitative and quantitative assessment of influence of cases on the analysis, i.e. influence analysis.

The sensitivity of a model is studied through measures that express its stability under perturbations. You are not interested in a model that is either overly stable or overly sensitive. Changes in the data or model components should produce commensurate changes in the model output. The difficulty is to determine when the changes are substantive enough to warrant further investigation, possibly leading to a reformulation of the model or changes in the data (such as dropping outliers). This paper is primarily concerned with stability of linear mixed models to perturbations of the data; that is, with influence analysis.

Statistical software environments, such as the R Programming language, provides a suite of tests and graphical procedures for appraising a fitted linear model, with several of these procedures analysing the model residuals.

2 Introduction

In classical linear models model diagnostics have become a required part of any statistical analysis, and the methods are commonly available in statistical packages and standard textbooks on applied regression. However it has been noted by several papers that model diagnostics do not often accompany LME model analyses. Model diagnostic techniques determine whether or not the distributional assumptions are satisfied, and to assess the influence of unusual observations.

3 Case-Deletion Diagnostics

Influence diagnostics are formal techniques that assess the influence of observations on parameter estimates for β and θ . A common technique is to refit the model with an observation or group of observations omitted.

Case-deletion diagnostics provide a useful tool for identifying influential observations and outliers. Since the pioneering work of Cook in 1977, deletion measures have been applied to many statistical models for identifying influential observations.

Standard statistical packages concentrate on calculating and testing parameter estimates without considering the diagnostics of the model.

The assessment of the effects of perturbations in data, on the outcome of the analysis, is known as statistical influence analysis. Influence analysis examines the robustness of the model.

4 Introduction

In classical linear models model diagnostics have become a required part of any statistical analysis, and the methods are commonly available in statistical packages and standard textbooks on applied regression. However it has been noted by several papers that model diagnostics do not often accompany LME model analyses.

4.1 Introduction

The linear mixed effects model is a useful methodology for fitting a wide range of models. However, linear mixed effects models are known to be sensitive to outliers. Christensen et al. (1992) advises that identification of outliers is necessary before conclusions may be drawn from the fitted model.

Standard statistical packages concentrate on calculating and testing parameter estimates without considering the diagnostics of the model.

The assessment of the effects of perturbations in data, on the outcome of the analysis, is known as statistical influence analysis. Influence analysis examines the robustness of the model.

Influence analysis methodologies have been used extensively in classical linear models, and provided the basis for methodologies for use with LME models.

Influence analysis examines the robustness of the model. Influence analysis methodologies have been used extensively in classical linear models, and provided the basis for methodologies for use with LME models.

Computationally inexpensive diagnostics tools have been developed to examine the issue of influence (Zewotir and Galpin, 2005). Studentized residuals, error contrast matrices and the inverse of the response variance covariance matrix are regular components of these tools.

Influence arises at two stages of the LME model. Firstly when V is estimated by \hat{V} , and subsequent estimations of the fixed and random regression coefficients β and u , given \hat{V} .

5 Checking model assumptions

In classical linear regression, it is important to carry out model diagnostic techniques to determine whether or not the distributional assumptions are satisfied. Model diagnostics are also used to determine the influence of unusual observations.

Schabenberger (2004) describes the examination of model-data agreement as comprising several elements; residual analysis, goodness of fit, collinearity diagnostics and influence analysis.

6 Influence analysis

Likelihood based estimation methods, such as ML and REML, are sensitive to unusual observations. Influence diagnostics are formal techniques that assess the influence of observations on parameter estimates for β and θ . A common technique is to refit the model with an observation or group of observations omitted.

West et al. (2007) examines a group of methods that examine various aspects of influence diagnostics for LME models. For overall influence, the most common approaches are the ‘likelihood distance’ and the ‘restricted likelihood distance’.

6.1 Model Data Agreement

Schabenberger (2004) describes the examination of model-data agreement as comprising several elements; residual analysis, goodness of fit, collinearity diagnostics and influence analysis.

6.2 Cook’s 1986 paper on Local Influence

Cook 1986 introduced methods for local influence assessment. These methods provide a powerful tool for examining perturbations in the assumption of a model, particularly

the effects of local perturbations of parameters of observations.

The local-influence approach to influence assessment is quite different from the case deletion approach, comparisons are of interest.

7 What is Influence

Broadly defined, “influence” is understood as the ability of a single or multiple data points, through their presence or absence in the data, to alter important aspects of the analysis, yield qualitatively different inferences, or violate assumptions of the statistical model. The goal of influence analysis is not primarily to mark data points for deletion so that a better model fit can be achieved for the reduced data, although this might be a result of influence analysis (Schabenberger, 2004).

Influence is defined as the ‘ability of a single or multiple data points, through their presence or absence

The goal of influence analysis is not primarily to mark data points for deletion so that a better model fit can be achieved for the reduced data, although this might be a result of influence analysis. The goal is rather to determine which cases are influential and the manner in which they are important to the analysis. Outliers, for example, may be the most noteworthy data points in an analysis. They can point to a model breakdown and lead to development of a better model.

Influence is understood to be the ability of a single or multiple data points, through their presences or absence in the data, to alter important aspects of the analysis, yield qualitatively different inferences, or violate assumptions of the statistical model (Schabenberger, 2004).

Outliers are the most noteworthy data points in an analysis, and an objective of influence analysis is how influential they are, and the manner in which they are influential.

7.1 Influence

The influence of an observation can be thought of in terms of how much the predicted scores for other observations would differ if the observation in question were not included.

Cook’s D is a good measure of the influence of an observation and is proportional to the sum of the squared differences between predictions made with all observations in the analysis and predictions made leaving out the observation in question. If the predictions are the same with or without the observation in question, then the observation has no influence on the regression model. If the predictions differ greatly when the observation is not included in the analysis, then the observation is influential.

7.2 Quantifying Influence

The basic procedure for quantifying influence is simple as follows:

- Fit the model to the data and obtain estimates of all parameters.
- Remove one or more data points from the analysis and compute updated estimates of model parameters.
- Based on full- and reduced-data estimates, contrast quantities of interest to determine how the absence of the observations changes the analysis.

Cook (1986) introduces powerful tools for local-influence assessment and examining perturbations in the assumptions of a model. In particular the effect of local perturbations of parameters or observations are examined.

7.3 Influence Analysis for LME Models

7.4 Influence Diagnostics: Basic Idea and Statistics

The general idea of quantifying the influence of one or more observations relies on computing parameter estimates based on all data points, removing the cases in question from the data, refitting the model, and computing statistics based on the change between full-data and reduced-data estimation. Influence statistics can be coarsely grouped by the aspect of estimation that is their primary target:

- overall measures compare changes in objective functions: (restricted) likelihood distance (Cook and Weisberg 1982, Ch. 5.2)
- influence on parameter estimates: Cook's (Cook 1977, 1979), MDFFITS (Belsley, Kuh, and Welsch 1980, p. 32)
- influence on precision of estimates: CovRatio and CovTrace
- influence on fitted and predicted values: PRESS residual, PRESS statistic (Allen 1974), DFFITS (Belsley, Kuh, and Welsch 1980, p. 15)
- outlier properties: internally and externally studentized residuals, leverage

Residual analysis is a widely used model validation technique. A residual is simply the difference between an observed value and the corresponding fitted value, as predicted by the model. The rationale is that, if the model is properly fitted to the model, then the residuals would approximate the random errors that one should expect. that is to say, if the residuals behave randomly, with no discernible trend, the model has fitted the data well. If some sort of non-random trend is evident in the model, then the model can be considered to be poorly fitted.

For LME models the matter of residual is more complex. ? describes two types of residuals, marginal residuals and conditional residuals. A marginal residual is the difference between the observed data and the estimated marginal mean. A conditional residual is the difference between the observed data and the predicted value of the observation. In a model without random effects, both sets of residuals coincide. We shall revert to this matter in due course.

Further to the analysis of residuals, ? recommends the examination of the following questions.

- Does the model-data agreement support the model assumptions?
- Should model components be refined, and if so, which components? For example, should regressors be added or removed, and is the covariation of the observations modeled properly?
- Are the results sensitive to model and/or data? Are individual data points or groups of cases particularly influential on the analysis?

7.5 Extension of Diagnostic Methods to LME models

When similar notions of statistical influence are applied to mixed models, things are more complicated. Removing data points affects fixed effects and covariance parameter estimates. Update formulas for *leave-one-out* estimates typically fail to account for changes in covariance parameters.

? noted the case deletion diagnostics techniques had not been applied to linear mixed effects models and seeks to develop methodologies in that respect. ? develops these techniques in the context of REML.

Demidenko (2004) extends several regression diagnostic techniques commonly used in linear regression, such as leverage, infinitesimal influence, case deletion diagnostics, Cook's distance, and local influence to the linear mixed-effects model. In each case, the proposed new measure has a direct interpretation in terms of the effects on a parameter of interest, and reduces to the familiar linear regression measure when there are no random effects.

The new measures that are proposed by Demidenko (2004) are explicitly defined functions and do not require re-estimation of the model, especially for cluster deletion diagnostics. The basis for both the cluster deletion diagnostics and Cook's distance is a generalization of Miller's simple update formula for case deletion for linear models. Furthermore Demidenko (2004) shows how Pregibon's infinitesimal case deletion diagnostics is adapted to the linear mixed-effects model.

Demidenko (2004) proposes two likelihood-based diagnostics for identifying individuals that can influence the choice between two competing models.

7.6 Extension of techniques to LME Models

Model diagnostic techniques, well established for classical models, have since been adapted for use with linear mixed effects models. Diagnostic techniques for LME models are inevitably more difficult to implement, due to the increased complexity.

Beckman et al. (1987) applied the local influence method of Cook (1986) to the analysis of the linear mixed model.

While the concept of influence analysis is straightforward, implementation in mixed models is more complex. Update formulae for fixed effects models are available only when the covariance parameters are assumed to be known.

If the global measure suggests that the points in U are influential, the nature of that influence should be determined. In particular, the points in U can affect the following

- the estimates of fixed effects,
- the estimates of the precision of the fixed effects,
- the estimates of the covariance parameters,
- the estimates of the precision of the covariance parameters,
- fitted and predicted values.

Diagnostic Methods for OLS models

Influence diagnostics are formal techniques allowing for the identification of observations that exert substantial influence on the estimates of fixed effects and variance covariance parameters.

The idea of influence diagnostics for a given observation is to quantify the effect of omission of this observation from the data on the results of the model fit. To this aim, the concept of likelihood displacement is used.

8 Analysis of Influence

Likelihood based estimation techniques, such as ML and REML, are sensitive to unusual observations. Influence diagnostics are formal techniques that assess the influence of observations on parameter estimates for β and θ . A common technique is to refit the model with an observation or group of observations omitted. West et al. (2007) examines a groups of methods that examine various aspects of the influence diagnostics for LME models. For overall influence, the most common approaches are the ‘likelihood distance’ and the ‘restricted likelihood distance’.

8.1 Further Assumptions of Linear Models

As with fitted models, the assumption of normality of residuals and homogeneity of variance is applicable to LMEs also.

Homoscedascity is the technical term to describe the variance of the residuals being constant across the range of predicted values. Heteroscedascity is the converse scenario : the variance differs along the range of values.

In recent years, mixed models have become invaluable tools in the analysis of experimental and observational data. In these models, more than one term can be subject to random variation. Mixed model technology enables you to analyze complex experimental data with hierarchical random processes, temporal, longitudinal, and spatial data, to name just a few important applications.

8.2 Summary of Schabenberger's Paper

On occasion, quantification is not possible. Assume, for example, that a data point is removed and the new estimate of the G matrix is not positive definite. This may occur if a variance component estimate now falls on the boundary of the parameter space. Thus, it may not be possible to compute certain influence statistics comparing the full-data and reduced-data parameter estimates. However, knowing that a new singularity was encountered is important qualitative information about the data points influence on the analysis.

We use the subscript (U) to denote quantities obtained without the observations in the set U. For example, (U) denotes the fixed-effects *leave-U-out* estimates. Note that the set U can contain multiple observations.

If the global measure suggests that the points in U are influential, you should next determine the nature of that influence. In particular, the points can affect

- the estimates of fixed effects
- the estimates of the precision of the fixed effects
- the estimates of the covariance parameters
- the estimates of the precision of the covariance parameters
- fitted and predicted values

It is important to further decompose the initial finding to determine whether data points are actually troublesome. Simply because they are influential somehow, should not trigger their removal from the analysis or a change in the model. For example, if points primarily affect the precision of the covariance parameters without exerting much influence on the fixed effects, then their presence in the data may not distort hypothesis tests or confidence intervals about β .

8.3 Summary of Paper

Standard residual and influence diagnostics for linear models can be extended to LME models. The dependence of the fixed effects solutions on the covariance parameters has important ramifications on the perturbation analysis. Calculating the studentized residuals-And influence statistics whereas each software procedure can calculate both conditional and marginal raw residuals, only SAs Proc Mixed is currently the only program that provide studentized residuals Which ave preferred for model diagnostics. The conditional Raw residuals ave not well suited to detecting outliers as are the studentized conditional residuals (Schabenberger, 2004).

LME are flexible tools for the analysis of clustered and repeated measurement data. LME extend the capabilities of standard linear models by allowing unbalanced and missing data, as long as the missing data are MAR. Structured covariance matrices for both the random effects G and the residuals R. missing at Random.

A conditional residual is the difference between the observed value and the predicted value of a dependent variable- Influence diagnostics are formal techniques that allow the identification observation that heavily influence estimates of parameters. To alleviate the problems with the interpretation of conditional residuals that may have unequal variances, we consider scaling. Residuals obtained in this manner ave called studentized residuals.

- Standard residual and inuence diagnostics for linear models can be extended to linear mixed models. The dependence of xed-effects solutions on the covariance parameter estimates has important ramications in perturbation analysis.
- To gauge the full impact of a set of observations on the analysis, covariance parameters need to be updated, which requires retting of the model.
- The conditional (subject-specific) and marginal (population-averaged) formulations in the linear mixed model enable you to consider conditional residuals that use the estimated BLUPs of the random effects, and marginal residuals which are deviations from the overall mean.
- Residuals using the BLUPs are useful to diagnose whether the random effects components in the model are specied correctly, marginal residuals are useful to diagnose the xed-effects components.
- Both types of residuals are available in SAS 9.1 as an experimental option of the MODEL statement in the MIXED procedure.
- It is important to note that influence analyses are performed under the assumption that the chosen model is correct. Changing the model structure can alter the conclusions. Many other variance models have been t to the data presented in the repeated measures example. You need to see the conclusions about which model component is affected in light of the model being fit.

9 Schabenberger: Summary and Conclusions

- Standard residual and influence diagnostics for linear models can be extended to linear mixed models. The dependence of fixed-effects solutions on the covariance parameter estimates has important ramifications in perturbation analysis.
- To gauge the full impact of a set of observations on the analysis, covariance parameters need to be updated, which requires refitting of the model.
- The experimental INFLUENCE option of the MODEL statement in the MIXED procedure (SAS 9.1) enables you to perform iterative and noniterative influence analysis for individual observations and sets of observations.
- The conditional (subject-specific) and marginal (population-averaged) formulations in the linear mixed model enable you to consider conditional residuals that use the estimated BLUPs of the random effects, and marginal residuals which are deviations from the overall mean.
- Residuals using the BLUPs are useful to diagnose whether the random effects components in the model are specified correctly, marginal residuals are useful to diagnose the fixed-effects components.
- Both types of residuals are available in SAS 9.1 as an experimental option of the MODEL statement in the MIXED procedure.
- It is important to note that influence analyses are performed under the assumption that the chosen model is correct. Changing the model structure can alter the conclusions. Many other variance models have been fit to the data presented in the repeated measures example. You need to see the conclusions about which model component is affected in light of the model being fit.

10 Effects on fitted and predicted values

$$\hat{e}_{i(U)} = y_i - x\hat{\beta}_{(U)} \quad (1)$$

11 Covariance Parameters

The unknown variance elements are referred to as the covariance parameters and collected in the vector θ .

12 Influence in LME Models

Model diagnostic techniques, well established for classical models, have since been adapted for use with linear mixed effects models. Diagnostic techniques for LME models are inevitably more difficult to implement, due to the increased complexity.

12.1 Key Definitions

Residual The difference between the predicted value (based on the regression equation) and the actual, observed value.

Outlier In linear regression, an outlier is an observation with large residual. In other words, it is an observation whose dependent-variable value is unusual given its value on the predictor variables. An outlier may indicate a sample peculiarity or may indicate a data entry error or other problem.

Leverage An observation with an extreme value on a predictor variable is a point with high leverage. Leverage is a measure of how far an independent variable deviates from its mean. High leverage points can have a great amount of effect on the estimate of regression coefficients.

Influence An observation is said to be influential if removing the observation substantially changes the estimate of the regression coefficients. Influence can be thought of as the product of leverage and outlieriness.

Cook's distance A measure that combines the information of leverage and residual of the observation.

13 Schabenberger

schab examines the use and implementation of influence measures in LME models.

Influence is understood to be the ability of a single or multiple data points, through their presences or absence in the data, to alter important aspects of the analysis, yield qualitatively different inferences, or violate assumptions of the statistical model (*schabenberger*).

Outliers are the most noteworthy data points in an analysis, and an objective of influence analysis is how influential they are, and the manner in which they are influential.

schab describes a simple procedure for quantifying influence. Firstly a model should be fitted to the data, and estimates of the parameters should be obtained. The second step is that either single or multiple data points, specifically outliers, should be omitted from the analysis, with the original parameter estimates being updated.

This is known as ‘*leave one out leave k out*’ analysis. The final step of the procedure is comparing the sets of estimates computed from the entire and reduced data sets to determine whether the absence of observations changed the analysis.

schabenberger notes that it is not always possible to derive influence statistics necessary for comparing full- and reduced-data parameter estimates.

In recent years, mixed models have become invaluable tools in the analysis of experimental and observational data. In these models, more than one term can be subject to random variation. Mixed model technology enables you to analyze complex experimental data with hierarchical random processes, temporal, longitudinal, and spatial data, to name just a few important applications.

schab remarks that the concept of critiquing the model-data agreement applies in mixed models in the same way as in linear fixed-effects models. In fact, because of the more complex model structure, you can argue that model and data diagnostics are even more important. For example, you are not only concerned with capturing the important variables in the model. You are also concerned with “distributing them correctly between the fixed and random components of the model. The mixed model structure presents unique and interesting challenges that prompt us to reexamine the traditional ideas of influence and residual analysis.

13.1 Overall Influence

An overall influence statistic measures the change in the objective function being minimized. For example, in OLS regression, the residual sums of squares serves that purpose. In linear mixed models fit by maximum likelihood (ML) or restricted maximum likelihood (REML), an overall influence measure is the likelihood distance [Cook and Weisberg].

14 Iterative and non-iterative influence analysis

Schabenberger (2004) highlights some of the issue regarding implementing mixed model diagnostics.

A measure of total influence requires updates of all model parameters.

however, this doesnt increase the procedures execution time by the same degree.

14.1 Iterative Influence Analysis

For linear models, the implementation of influence analysis is straightforward. However, for LME models, the process is more complex. Update formulas for the fixed effects are available only when the covariance parameters are assumed to be known. A measure of total influence requires updates of all model parameters. This can only be achieved in general is by omitting observations, then refitting the model.

Schabenberger (2004) describes the choice between iterative influence analysis and non-iterative influence analysis.

14.2 Iterative vs Non-Iterative Influence Analysis

While the basic idea of influence analysis is straightforward, the implementation in mixed models can be tricky. For example, update formulas for the fixed effects are available only when the covariance parameters are assumed to be known. At most the profiled residual variance can be updated without refitting the model.

A measure of total influence requires updates of all model parameters, and the only way that this can be achieved in general is by removing the observations in question and refitting the model.

Because this **bruteforce** method involves iterative reestimation of the covariance parameters, it is termed *iterative influence analysis*. Reliance on closed-form update formulas for the fixed effects without updating the (un-profiled) covariance parameters is termed a noniterative influence analysis.

An iterative analysis seems like a costly, computationally intensive enterprise. If you compute iterative influence diagnostics for all n observations, then a total of $n + 1$ mixed models are fit iteratively. This does not imply, of course, that the procedures execution time increases n -fold. Keep in mind that

- iterative reestimation always starts at the converged full-data estimates. If a data point is not influential, then its removal will have little effect on the objective function and parameter estimates. Within one or two iterations, the process should arrive at the reduced-data estimates.
- if complete reestimation does require many iterations, then this is important information in itself. The likelihood surface has probably changed drastically, and the reduced-data estimates are moving away

from the full-data estimates.

References

- Beckman, R., C. Nachtsheim, and R. Cook (1987). Diagnostics for mixed-model analysis of variance. *Technometrics* 29(4), 413–426.
- Christensen, R., L. M. Pearson, and W. Johnson (1992). Case-deletion diagnostics for mixed models. *Technometrics* 34(1), 38–45.
- Cook, R. (1986). Assessment of local influence. *Journal of the Royal Statistical Society. Series B (Methodological)* 48(2), 133–169.

- Demidenko, E. (2004). *Mixed Models: Theory And Application*. Dartmouth College: Wiley Interscience.
- Schabenberger, O. (2004). Mixed model influence diagnostics. 18929.
- West, B., K. Welch, and A. Galecki (2007). *Linear Mixed Models: a Practical Guide Using Statistical Software*. Chapman and Hall CRC.
- Zewotir, T. and J. S. Galpin (2005). Influence diagnostics for linear mixed models. *Journal of Data Science* 3(2), 153–177.