

# Contents

<b>1</b>	<b>Method Comparison Studies</b>	<b>4</b>
1.1	Introduction . . . . .	4
1.2	Repeatability . . . . .	7
1.2.1	Repeatability and Gold Standards . . . . .	8
<b>2</b>	<b>Review of Current Methodologies</b>	<b>10</b>
2.1	Bland-Altman Approach . . . . .	10
2.1.1	Bland-Altman plots . . . . .	13
2.1.2	Bland-Altman plots for the Grubbs data . . . . .	14
2.1.3	Prevalence of the Bland-Altman plot . . . . .	17
2.1.4	Adverse features . . . . .	18
2.2	Limits of Agreement . . . . .	23
2.2.1	Inferences on Bland-Altman estimates . . . . .	25
2.2.2	Formal definition of limits of agreement . . . . .	25
2.2.3	Alternative Agreement Indices . . . . .	26
2.3	Variations and Alternative Graphical Methods . . . . .	30
2.3.1	Variants of the Bland-Altman Plot . . . . .	31
2.3.2	Replicate Measurements . . . . .	37
2.3.3	Prevalence of the Bland-Altman plot . . . . .	38
2.4	The Technology Acceptance Model . . . . .	38
2.5	Formal Models and Tests . . . . .	40

2.5.1	Morgan Pitman Testing . . . . .	42
2.5.2	Paired sample $t$ test . . . . .	42
2.6	Blackwood Bradley Model . . . . .	43
2.6.1	Bland-Altman correlation test . . . . .	44
2.6.2	Identifiability . . . . .	45
2.7	Regression Methods for Method Comparison . . . . .	46
2.8	Regression Methods (duplication) . . . . .	47
2.8.1	Deming Regression . . . . .	49
2.9	Other Types of Studies . . . . .	51
<b>3</b>	<b>Linear Mixed effects Models</b>	<b>54</b>
3.1	Linear Mixed effects Models . . . . .	54
3.1.1	Estimation . . . . .	56
3.2	Repeated measurements in LME models . . . . .	61
3.2.1	Formulation of the Response Vector . . . . .	61
3.2.2	Decomposition of the response covariance matrix . . . . .	62
3.2.3	Correlation terms . . . . .	63
3.3	Using LME for method comparison . . . . .	65
3.3.1	Roy's Approach . . . . .	65
3.3.2	Likelihood Ratio Tests . . . . .	66
3.3.3	Testing Procedures . . . . .	67
3.3.4	Correlation . . . . .	68
3.3.5	Variability test 1 . . . . .	69
3.3.6	Variability test 2 . . . . .	69
3.3.7	Variability test 3 . . . . .	70
3.3.8	Demonstration of Roy's testing . . . . .	70
3.4	Limits of agreement in LME models . . . . .	73
3.4.1	Linked replicates . . . . .	74
3.5	Implementation in R . . . . .	77

<b>4</b>	<b>Extending Current Methodologies</b>	<b>81</b>
4.1	Extension of Roy's methodology . . . . .	81
4.2	Conclusion . . . . .	82
4.3	Outline of Thesis . . . . .	83
	Bibliography . . . . .	83

# Chapter 1

## Method Comparison Studies

### 1.1 Introduction

The problem of assessing the agreement between two or more methods of measurement is ubiquitous in scientific research, and is commonly referred to as a ‘method comparison study’. Published examples of method comparison studies can be found in disciplines as diverse as pharmacology (Ludbrook, 1997), anaesthesia (Myles, 2007), and cardiac imaging methods (Krummenauer et al., 2000).

To illustrate the characteristics of a typical method comparison study consider the data in Table I (Grubbs, 1973). In each of twelve experimental trials, a single round of ammunition was fired from a 155mm gun and its velocity was measured simultaneously (and independently) by three chronographs devices, identified here by the labels ‘Fotobalk’, ‘Counter’ and ‘Terma’.

Round	Fotobalk [F]	Counter [C]	Terma [T]
1	793.8	794.6	793.2
2	793.1	793.9	793.3
3	792.4	793.2	792.6
4	794.0	794.0	793.8
5	791.4	792.2	791.6
6	792.4	793.1	791.6
7	791.7	792.4	791.6
8	792.3	792.8	792.4
9	789.6	790.2	788.5
10	794.4	795.0	794.7
11	790.9	791.6	791.3
12	793.5	793.8	793.5

Table 1.1.1: Velocity measurement from the three chronographs (Grubbs 1973).

An important aspect of these data is that all three methods of measurement are assumed to have an attended measurement error, and the velocities reported in Table 1.1 can not be assumed to be ‘true values’ in any absolute sense.

A method of measurement should ideally be both accurate and precise. Barnhart et al. (2007) describes agreement as being a broader term that contains both of those qualities. An accurate measurement method will give results close to the unknown ‘true value’. The precision of a method is indicated by how tightly measurements obtained under identical conditions are distributed around their mean measurement value. A precise and accurate method will yield results consistently close to the true value. Of course a method may be accurate, but not precise, if the average of its measurements is close to the true value, but those measurements are highly dispersed. Conversely a method that is not accurate may be quite precise, as it consistently indicates the same level of inaccuracy. The tendency of a method of measurement to consistently give

results above or below the true value is a source of systematic bias. The smaller the systematic bias, the greater the accuracy of the method.

In the context of the agreement of two methods, there is also a tendency of one measurement method to consistently give results above or below the other method. Lack of agreement is a consequence of the existence of ‘inter-method bias’. For two methods to be considered in good agreement, the inter-method bias should be in the region of zero. A simple estimate of the inter-method bias is given by the differences between pairs of measurements, for example, Table 1.1.2 is a good example of possible inter-method bias; the ‘Fotobalk’ consistently recording smaller velocities than the ‘Counter’ method. A cursory inspection of the table will indicate a systematic tendency for the Counter method to result in higher measurements than the Fotobalk method.

The absence of inter-method bias is, by itself, not sufficient to establish that two measurement methods agree. The two methods must also have equivalent levels of precision. Should one method yield results considerably more variable than those of the other, they can not be considered to be in agreement. Hence, method comparison studies are required to take account of both inter-method bias and difference in precision of measurements.

Round	Fotobalk (F)	Counter (C)	Difference (F-C)
1	793.8	794.6	-0.8
2	793.1	793.9	-0.8
3	792.4	793.2	-0.8
4	794.0	794.0	0.0
5	791.4	792.2	-0.8
6	792.4	793.1	-0.7
7	791.7	792.4	-0.7
8	792.3	792.8	-0.5
9	789.6	790.2	-0.6
10	794.4	795.0	-0.6
11	790.9	791.6	-0.7
12	793.5	793.8	-0.3

Table 1.1.2: Difference between Fotobalk and Counter measurements.

## 1.2 Repeatability

Repeatability is the ability of a measurement method to give consistent results for a particular subject, i.e. a measurement will agree with prior and subsequent measurements of the same subject. Barnhart et al. (2007) emphasizes the importance of repeatability as part of an overall method comparison study, a view endorsed by Carstensen et al. (2008). Before there can be good agreement between two methods, a method must have good agreement with itself. If one method has poor repeatability in the sense of considerable variability, then agreement between two methods is bound to be poor (Roy, 2009b). Barnhart et al. (2007) remarks that it is important to report repeatability when assessing measurement, because it measures the purest form of random error not influenced by other factors, while further remarking ‘*curiously replicate measure-*

ments are rarely made in method comparison studies, so that an important aspect of comparability is often overlooked. Bland and Altman (1999) strongly recommends the simultaneous estimation of repeatability and agreement by collecting replicated data. However Roy (2009b) notes the lack of convenience in such calculations. Repeatability is defined by the IUPAC (2009) as ‘*the closeness of agreement between independent results obtained with the same method on identical test material, under the same conditions (same operator, same apparatus, same laboratory and after short intervals of time)*’ and is determined by taking multiple measurements on a series of subjects.

A measurement is said to be repeatable when this variation is smaller than some pre-specified limit. In these situations, there is often a predetermined ”critical difference”, and for differences in monitored values that are smaller than this critical difference, the possibility of pre-test variability as a sole cause of the difference may be considered in addition to, for examples, changes in diseases or treatments.

The British Standards Institute (1979) defines a coefficient of repeatability as *the value below which the difference between two single test results may be expected to lie within a specified probability*. In the absence of other indications, the probability is 95%.

### 1.2.1 Repeatability and Gold Standards

Currently the phrase ‘gold standard’ describes the most accurate method of measurement available. No other criteria are set out. Further to Dunn (2002), various gold standards have a varying levels of repeatability. Dunn cites the example of the sphygmomanometer, which is prone to measurement error. Consequently it can be said that a measurement method can be the ‘gold standard’, yet have poor repeatability.

Dunn (2002) recognizes this problem. Hence, if the most accurate method is considered to have poor repeatability, it is referred to as a ‘bronze standard’. Again, no formal definition of a ‘bronze standard’ exists.

The coefficient of repeatability may provide the basis of formulation a formal defi-



nition of a ‘gold standard’. For example, by determining the ratio of  $CR$  to the sample mean  $\bar{X}$ . Advisably the sample size should be specified in advance. A gold standard may be defined as the method with the lowest value of  $\lambda = CR/\bar{X}$  with  $\lambda < 0.1\%$ . Similarly, a silver standard may be defined as the method with the lowest value of  $\lambda$  with  $0.1\% \leq \lambda < 1\%$ . Such thresholds are solely for expository purposes.

# Chapter 2

## Review of Current Methodologies

### 2.1 Bland-Altman Approach

The issue of whether two measurement methods comparable to the extent that they can be used interchangeably with sufficient accuracy is encountered frequently in scientific research. Historically, comparison of two methods of measurement was carried out by use of paired sample  $t$ -test, correlation coefficients or simple linear regression. However, simple linear regression is unsuitable for method comparison studies due to the assumption that one variable is measured without error. In comparing two methods, both methods are assumed to have attendant random error.

Altman and Bland (1983) highlighted the inadequacies of these approaches for comparing two methods of measurement, and proposed methodologies with this specific application in mind. Although the authors also acknowledge the opportunity to apply other, more complex, approaches, but argue that simpler approaches is preferable, especially when the results must be ‘explained to non-statisticians’.

Notwithstanding previous remarks about linear regression, the first step recommended, which the authors argue should be mandatory, is the construction of a scatter plot of the data. Scatterplots can facilitate an initial judgement and helping to identify potential outliers, with the addition of the line of equality. In the case of good agree-

ment, the observations would be distributed closely along this line. However, they are not useful for a thorough examination of the data. O'Brien et al. (1990) notes that data points will tend to cluster around the line of equality, obscuring interpretation.

A scatter plot of the Grubbs data is shown in Figure 1.1. Visual inspection confirms the previous conclusion that inter-method bias is present, i.e. the Fotobalk device has a tendency to record a lower velocity.




Figure 2.1.1: Scatter plot for Fotobalk and Counter methods.

Dewitte et al. (2002) notes that scatter plots were very seldom presented in the *Annals of Clinical Biochemistry*. This apparently results from the fact that the 'Instructions for Authors' dissuade the use of regression analysis, which conventionally is

accompanied by a scatter plot.

### 2.1.1 Bland-Altman plots

In light of shortcomings associated with scatterplots, Altman and Bland (1983) recommend a further analysis of the data. Firstly case-wise differences of measurements of two methods  $d_i = y_{1i} - y_{2i}$ , for  $i = 1, 2, \dots, n$ , on the same subject should be calculated, and then the average of those measurements, ( $a_i = (y_{1i} + y_{2i})/2$  for  $i = 1, 2, \dots, n$ ).

Altman and Bland (1983) proposed that  $a_i$  should be plotted against  $d_i$ , a plot now widely known as the Bland-Altman plot, and motivated this plot as follows:

“From this type of plot it is much easier to assess the magnitude of disagreement (both error and bias), spot outliers, and see whether there is any trend, for example an increase in (difference) for high values. This way of plotting the data is a very powerful way of displaying the results of a method comparison study.”

The case wise-averages capture several aspects of the data, such as expressing the range over which the values were taken, and assessing whether the assumptions of constant variance holds. Case-wise averages also allow the case-wise differences to be presented on a two-dimensional plot, with better data visualization qualities than a one dimensional plot. Bland and Altman (1986) cautions that it would be the difference against either measurement value instead of their average, as the difference relates to both value. This approach has proved very popular, and the Bland-Altman plots is widely regarded as powerful graphical tool for making a visual assessment of the data.

The magnitude of the inter-method bias between the two methods is simply the average of the differences  $\bar{d}$ . This inter-method bias is represented with a line on the Bland-Altman plot. As the objective of the Bland-Altman plot is to advise on the agreement of two methods, the individual case-wise differences are also particularly relevant. The variances around this bias is estimated by the standard deviation of these differences  $S_d$ .

### 2.1.2 Bland-Altman plots for the Grubbs data

In the case of the Grubbs data the inter-method bias is  $-0.61$  metres per second, and is indicated by the dashed line on Figure 1.2. By inspection of the plot, it is also possible to compare the precision of each method. Noticeably the differences tend to increase as the averages increase.

The Bland-Altman plot for comparing the ‘Fotobalk’ and ‘Counter’ methods, which shall henceforth be referred to as the ‘F vs C’ comparison, is depicted in Figure 1.2, using data from Table 1.3. The presence and magnitude of the inter-method bias is indicated by the dashed line.

Round	Fotobalk [F]	Counter [C]	Differences [F-C]	Averages [(F+C)/2]
1	793.8	794.6	-0.8	794.2
2	793.1	793.9	-0.8	793.5
3	792.4	793.2	-0.8	792.8
4	794.0	794.0	0.0	794.0
5	791.4	792.2	-0.8	791.8
6	792.4	793.1	-0.7	792.8
7	791.7	792.4	-0.7	792.0
8	792.3	792.8	-0.5	792.5
9	789.6	790.2	-0.6	789.9
10	794.4	795.0	-0.6	794.7
11	790.9	791.6	-0.7	791.2
12	793.5	793.8	-0.3	793.6

Table 2.1.1: Fotobalk and Counter methods: differences and averages.

Round	Fotobalk [F]	Terma [T]	Differences [F-T]	Averages [(F+T)/2]
1	793.8	793.2	0.6	793.5
2	793.1	793.3	-0.2	793.2
3	792.4	792.6	-0.2	792.5
4	794.0	793.8	0.2	793.9
5	791.4	791.6	-0.2	791.5
6	792.4	791.6	0.8	792.0
7	791.7	791.6	0.1	791.6
8	792.3	792.4	-0.1	792.3
9	789.6	788.5	1.1	789.0
10	794.4	794.7	-0.3	794.5
11	790.9	791.3	-0.4	791.1
12	793.5	793.5	0.0	793.5

Table 2.1.2: Fotobalk and Terma methods: differences and averages.




Figure 2.1.2: Bland-Altman plot For Fotobalk and Counter methods.

In Figure 1.3 Bland-Altman plots for the ‘F vs C’ and ‘F vs T’ comparisons are shown, where ‘F vs T’ refers to the comparison of the ‘Fotobalk’ and ‘Terma’ methods. Usage of the Bland-Altman plot can be demonstrate in the contrast between these comparisons. By inspection, there exists a larger inter-method bias in the ‘F vs C’ comparison than in the ‘F vs T’ comparison. Conversely there appears to be less precision in ‘F vs T’ comparison, as indicated by the greater dispersion of covariates.






Figure 2.1.3: Bland-Altman plots for Grubbs’ F vs C and F vs T comparisons.

### 2.1.3 Prevalence of the Bland-Altman plot

Bland and Altman (1986), which further develops the Bland-Altman approach, was found to be the sixth most cited paper of all time by Ryan and Woodall (2005). Dewitte et al. (2002) describes the rate at which prevalence of the Bland-Altman plot has developed in scientific literature, by examining all articles in the journal ‘Clinical Chemistry’ between 1995 and 2001. This study concluded that use of the BlandAltman plot increased over the years, from 8% in 1995 to 14% in 1996, and 3136% in 2002.

The Bland-Altman Plot has since become expected, and often obligatory, approach for presenting method comparison studies in many scientific journals (Hollis, 1996). Furthermore O’Brien et al. (1990) recommend its use in papers pertaining to method comparison studies for the journal of the British Hypertension Society.

### 2.1.4 Adverse features

Estimates for inter-method bias and variance of differences are only meaningful if there is uniform inter-bias and variability throughout the range of measurements. Fulfilment of these assumptions can be checked by visual inspection of the plot. The prototype Bland-Altman plots depicted in Figures 1.4, 1.5 and 1.6 are derived from simulated data, for the purpose of demonstrating how the plot would inform an analyst of features that would adversely affect use of the recommended approach.

Figure 1.4 demonstrates how the Bland-Altman plot would indicate increasing variance of differences over the measurement range. Fitted regression lines, for both the upper and lower half of the plot, has been added to indicate the trend. Figure 1.5 is an example of cases where the inter-method bias changes over the measurement range. This is known as proportional bias, and is defined by Ludbrook (1997) as meaning that ‘one method gives values that are higher (or lower) than those from the other by an amount that is proportional to the level of the measured variable’. In both Figures 1.4 and 1.5, the assumptions necessary for further analysis using the limits of agreement are violated.

Application of regression techniques to the Bland-Altman plot, and subsequent formal testing for the constant variability of differences is informative. The data set may be divided into two subsets, containing the observations wherein the difference values are less than and greater than the inter-method bias respectively. For both of these fits, hypothesis tests for the respective slopes can be performed. While both tests could be considered separately, multiple comparison procedures, such as the Benjamini-Hochberg (Benjamini and Hochberg, 1995) test, are advisable.




Figure 2.1.4: Bland-Altman plot demonstrating the increase of variance over the range.

The Bland-Altman plot also can be used to identify outliers. An outlier is an observation that is conspicuously different from the rest of the data that it arouses suspicion that it occurs due to a mechanism, or conditions, different to that of the rest of the observations. Bland and Altman (1999) do not recommend excluding outliers from analyses, but remark that recalculation of the inter-method bias estimate, and further calculations based upon that estimate, are useful for assessing the influence of outliers. The authors remark that ‘we usually find that this method of analysis is not too sensitive to one or two large outlying differences’. Figure 1.6 demonstrates how the Bland-Altman plot can be used to visually inspect the presence of potential outliers.

As a complement to the Bland-Altman plot, Bartko (1994) proposes the use of a bivariate confidence ellipse, constructed for a predetermined level. Altman (1978) provides the relevant calculations for the ellipse. This ellipse is intended as a visual guidelines for the scatter plot, for detecting outliers and to assess the within- and between-subject variances.




Figure 2.1.5: Bland-Altman plot indicating the presence of proportional bias.




Figure 2.1.6: Bland-Altman plot indicating the presence of potential outliers.

The minor axis relates to the between subject variability, whereas the major axis relates to the error mean square, with the ellipse depicting the size of both relative to each other. Consequently Bartko's ellipse provides a visual aid to determining the relationship between variances. If  $\text{var}(a)$  is greater than  $\text{var}(d)$ , the orientation of the ellipse is horizontal. Conversely if  $\text{var}(a)$  is less than  $\text{var}(d)$ , the orientation of the ellipse is vertical.

The Bland-Altman plot for the Grubbs data, complemented by Bartko's ellipse, is depicted in Figure 1.7. The fourth observation is shown to be outside the bounds of the ellipse, indicating that it is a potential outlier.




Figure 2.1.7: Bartko's Ellipse for Grubbs' data.

The limitations of using bivariate approaches to outlier detection in the Bland-Altman plot can be demonstrated using Bartko's ellipse. A covariate is added to the 'F vs C' comparison that has a difference value equal to the inter-method bias, and an average value that markedly deviates from the rest of the average values in the comparison, i.e. 786. Table 1.8 depicts a 95% confidence ellipse for this manipulated data set. By inspection of the confidence interval, we would conclude that this extra

covariate is an outlier, in spite of the fact that this observation is very close to the inter-method bias as determined by this approach.




Figure 2.1.8: Bartko's Ellipse for Grubbs' data, with an extra covariate.

Importantly, outlier classification must be informed by the logic of the mechanism that produces the data. In the Bland-Altman plot, the horizontal displacement (i.e. the average) of any observation is supported by two separate measurements. Any observation should not be considered an outlier on the basis of a noticeable horizontal displacement from the main cluster, as in the case with the extra covariate. Conversely, the fourth observation, from the original data set, should be considered an outlier, as it has a noticeable vertical displacement from the rest of the observations.

In classifying whether a observation from a univariate data set is an outlier, many formal tests are available, such as the Grubbs test for outliers. In assessing whether a covariate in a Bland-Altman plot is an outlier, this test is useful when applied to the case-wise difference values treated as a univariate data set. The null hypothesis of the Grubbs test procedure is the absence of any outliers in the data set. Conversely, the alternative hypotheses is that there is at least one outlier present.

The test statistic for the Grubbs test ( $G$ ) is the largest absolute deviation from the sample mean divided by the standard deviation of the differences,

$$G = \max_{i=1,\dots,n} \frac{|d_i - \bar{d}|}{S_d}.$$

For the ‘F vs C’ comparison it is the fourth observation gives rise to the test statistic,  $G = 3.64$ . The critical value is calculated using Student’s  $t$  distribution and the sample size,

$$U = \frac{n-1}{\sqrt{n}} \sqrt{\frac{t_{\alpha/(2n),n-2}^2}{n-2+t_{\alpha/(2n),n-2}^2}}.$$

For this test  $U = 0.75$ . The conclusion of this test is that the fourth observation in the ‘F vs C’ comparison is an outlier, with  $p$ -value = 0.003, in accordance with the previous result of Bartko’s ellipse.

## 2.2 Limits of Agreement

A third element of the Bland-Altman approach, an interval known as ‘limits of agreement’ is introduced in Bland and Altman (1986) (sometimes referred to in literature as 95% limits of agreement). Limits of agreement are used to assess whether the two methods of measurement can be used interchangeably. Bland and Altman (1986) refer to this as the ‘equivalence’ of two measurement methods. The specific question to which limits of agreement are intended as the answer to must be established clearly. Bland and Altman (1995) comment that the limits of agreement show ‘how far apart measurements by the two methods were likely to be for most individuals’, a definition echoed in their 1999 paper:

“We can then say that nearly all pairs of measurements by the two methods will be closer together than these extreme values, which we call 95% limits of agreement. These values define the range within which most differences between measurements by the two methods will lie.”

The limits of agreement (LoA) are computed by the following formula:

$$LoA = \bar{d} \pm 1.96s_d$$

with  $\bar{d}$  as the estimate of the inter method bias,  $s_d$  as the standard deviation of the differences and 1.96 (sometimes rounded to 2) is the 95% quantile for the standard normal distribution. The limits of agreement methodology assumes a constant level of bias throughout the range of measurements. Importantly the authors recommend prior determination of what would constitute acceptable agreement, and that sample sizes should be predetermined to give an accurate conclusion. However Mantha et al. (2000) highlight inadequacies in the correct application of limits of agreement, resulting in contradictory estimates of limits of agreement in various papers.

For the Grubbs ‘F vs C’ comparison, these limits of agreement are calculated as -0.132 for the upper bound, and -1.08 for the lower bound. Figure 1.9 shows the resultant Bland-Altman plot, with the limits of agreement shown in dashed lines.




Figure 2.2.9: Bland-Altman plot with limits of agreement



### 2.2.1 Inferences on Bland-Altman estimates

Bland and Altman (1999) advises on how to calculate confidence intervals for the inter-method bias and limits of agreement. For the inter-method bias, the confidence interval is simply that of a mean:  $\bar{d} \pm t_{(\alpha/2, n-1)} S_d / \sqrt{n}$ . The confidence intervals and standard error for the limits of agreement follow from the variance of the limits of agreement, which is shown to be

$$\text{Var}(LoA) = \left( \frac{1}{n} + \frac{1.96^2}{2(n-1)} \right) s_d^2.$$

If  $n$  is sufficiently large this can be following approximation can be used

$$\text{Var}(LoA) \approx 1.71^2 \frac{s_d^2}{n}.$$

Consequently the standard errors of both limits can be approximated as 1.71 times the standard error of the differences.

A 95% confidence interval can be determined, by means of the  $t$  distribution with  $n - 1$  degrees of freedom. However, Bland and Altman (1999) comment that such calculations may be ‘somewhat optimistic’ on account of the associated assumptions not being realized.

### 2.2.2 Formal definition of limits of agreement

Bland and Altman (1999) note the similarity of limits of agreement to confidence intervals, but are clear that they are not the same thing. Interestingly, they describe the limits as ‘being like a reference interval’.

Limits of agreement have very similar construction to Shewhart control limits. The Shewhart chart is a well known graphical methodology used in statistical process control. Consequently there is potential for misinterpreting the limits of agreement as they were Shewhart control limits.

Carstensen et al. (2008) regards the limits of agreement as a prediction interval for the difference between future measurements with the two methods on a new individual,

but states that it does not fit the formal definition of a prediction interval, since the definition does not consider the errors in estimation of the parameters. Prediction intervals, which are often used in regression analysis, are estimates of an interval in which future observations will fall, with a certain probability, given what has already been observed. Carstensen et al. (2008) offers an alternative formulation, a 95% prediction interval for the difference

$$\bar{d} \pm t_{(0.025, n-1)} s_d \sqrt{1 + \frac{1}{n}}$$

where  $n$  is the number of subjects. Carstensen is careful to consider the effect of the sample size on the interval width, adding that only for 61 or more subjects is the quantile less than 2.

Luiz et al. (2003) offers an alternative description of limits of agreement, this time as tolerance limits. A tolerance interval for a measured quantity is the interval in which a specified fraction of the population's values lie, with a specified level of confidence. Barnhart et al. (2007) describes them as a probability interval, and offers a clear description of how they should be used; 'if the absolute limit is less than an acceptable difference  $d_0$ , then the agreement between the two methods is deemed satisfactory'.

The prevalence of contradictory definitions of what limits of agreement strictly are will inevitably attenuate the poor standard of reporting using limits of agreement, as mentioned by Mantha et al. (2000).

### 2.2.3 Alternative Agreement Indices

As an alternative to limits of agreement, Lin et al. (2002) proposes the use of the mean square deviation in assessing agreement. The mean square deviation is defined as the expectation of the squared differences of two readings. The MSD is usually used for the case of two measurement methods  $X$  and  $Y$ , each making one measurement for the same subject, and is given by

$$MSD_{xy} = E[(x - y)^2] = (\mu_x - \mu_y)^2 + (\sigma_x - \sigma_y)^2 + 2\sigma_x\sigma_y(1 - \rho_{xy}).$$

Barnhart et al. (2007) advises the use of a predetermined upper limit for the MSD value,  $MSD_{ul}$ , to define satisfactory agreement. However, a satisfactory upper limit may not be easily determinable, thus creating a drawback to this methodology.

Alternative indices, proposed by Barnhart et al. (2007), are the square root of the MSD and the expected absolute difference (EAD).

$$EAD = E(|x - y|) = \frac{\sum |x_i - y_i|}{n}$$

Both of these indices can be interpreted intuitively, since their units are the same as that of the original measurements. Also they can be compared to the maximum acceptable absolute difference between two methods of measurement  $d_0$ .

The EAD can be used to supplement the inter-method bias in an initial comparison study, as the EAD is informative as a measure of dispersion, is easy to calculate and requires no distributional assumptions. A consequence of using absolute differences is that high variances would result in a higher EAD value.

	X	Y	U	V
1	101.83	102.52	98.05	99.53
2	101.68	102.69	99.17	96.53
3	97.89	99.01	100.31	97.55
4	98.15	99.57	100.35	96.03
5	99.94	100.85	99.51	99.00
6	98.85	98.86	98.50	100.76
7	99.86	97.85	100.66	99.37
8	101.57	100.21	99.66	108.87
9	100.12	99.85	99.70	105.16
10	99.49	98.77	101.55	94.31

Differences	2.5% limit	97.5% limit	SD(diff)
-0.08078844	-2.39471014	2.23313327	1.15696085

	X	Y	$X - Y$	$ X - Y $
1	98.05	99.53	-1.49	1.49
2	99.17	96.53	2.64	2.64
3	100.31	97.55	2.75	2.75
4	100.35	96.03	4.32	4.32
5	99.51	99.00	0.51	0.51
6	98.50	100.76	-2.26	2.26
7	100.66	99.37	1.29	1.29
8	99.66	108.87	-9.21	9.21
9	99.70	105.16	-5.45	5.45
10	101.55	94.31	7.24	7.24

Table 2.2.3: Example data set

To illustrate the use of EAD, consider table 2.2.3. The inter-method bias is 0.03, which is quite close to zero, and conducive to agreement between methods. However, an identity plot would indicate very poor agreement, as the points are noticeably distant from the line of equality.

The limits of agreement are  $[-9.61, 9.68]$ , a wide interval for this data. As with the identity plot, this would indicate lack of agreement. The EAD is 3.71.

The Bland-Altman plot remains a useful part of the analysis. In 2.2.11, it is clear there is a systematic decrease in differences across the range of measurements.

Barnhart et al. (2007) remarks that a comparison of EAD and MSD, using simulation studies, would be interesting, while further adding that ‘*It will be of interest to investigate the benefits of these possible new unscaled agreement indices*’. For the Grubbs’ ‘F vs C’ and ‘F vs T’ comparisons, the inter-method bias, difference variances, limits of agreement and EADs are shown in Table 1.5. The corresponding Bland-Altman plots for ‘F vs C’ and ‘F vs T’ comparisons were depicted previously on Figure 1.3. While the inter-method bias for the ‘F vs T’ comparison is smaller, the EAD penalizes the comparison for having a greater variance of differences. Hence the




Figure 2.2.10: Identity Plot for example data




Figure 2.2.11: Expected Absolute Difference

EAD values for both comparisons are much closer.

	F vs C	F vs T
Inter-method bias	-0.61	0.12
Difference variance	0.06	0.22
Limits of agreement	(-1.08, -0.13)	(-0.81,1.04)
EAD	0.61	0.35

Table 2.2.4: Agreement indices for Grubbs’ data comparisons.

Further to Lin (2000) and Lin et al. (2002), individual agreement between two measurement methods may be assessed using the the coverage probability (CP) criteria or the total deviation index (TDI). If  $d_0$  is predetermined as the maximum acceptable absolute difference between two methods of measurement, the probability that the absolute difference of two measures being less than  $d_0$  can be computed. This is known as the coverage probability (CP).

$$CP = P(|x_i - y_i| \leq d_0) \quad (2.1)$$

If  $\pi_0$  is set as the predetermined coverage probability, the boundary under which the proportion of absolute differences is  $\pi_0$  may be determined. This boundary is known as the ‘total deviation index’ (TDI). Hence the TDI is the  $100\pi_0$  percentile of the absolute difference of paired observations.

## 2.3 Variations and Alternative Graphical Methods

In this section, we will look at some variations and enhancements of the Bland-Altman plot, as well as some alternative graphical techniques. Strictly speaking, the Identity Plot is advised by Bland and Altman as a prior analysis to the Bland-Alman plot, and therefore is neither a variant nor an alternative approach. However it is worth mentioning, as it is a simple, powerful and elegant technique that is often overlooked

in method comparison studies. The identity plot is a simple scatter-plot approach of measurements for both methods on either axis, with the line of equality (the  $X = Y$  line, i.e. the 45 degree line through the origin). This plot can give the analyst a cursory examination of how well the measurement methods agree. In the case of good agreement, the covariates of the plot accord closely with the line of equality.

### **2.3.1 Variants of the Bland-Altman Plot**

In light of some potential pitfalls associated with the conventional difference plot, a series of alternative formulations for the Bland-Altman approach have been proposed.

Referring to the assumption that bias and variability are constant across the range of measurements, Bland and Altman (1999) address the case where there is an increase in variability as the magnitude increases. They remark that it is possible to ignore the issue altogether, but the limits of agreement would be wider apart than necessary when just lower magnitude measurements are considered. Conversely the limits would be too narrow should only higher magnitude measurements be used. To address the issue, they propose the logarithmic transformation of the data. The plot is then formulated as the difference of paired log values against their mean. Bland and Altman acknowledge that this is not easy to interpret, and may not be suitable in all cases.

### **Bland and Altman's Percentage and Ratio Plots**

Bland and Altman (1999) offer two variations of the Bland-Altman plot intended to overcome situations where the conventional plot is inappropriate. The first variation is a plot of casewise differences as percentage of averages, and is appropriate when the variability of the differences increase as the magnitude increases.

The second variation is a plot of casewise ratios as percentage of averages. This will remove the need for logarithmic transformation. This approach is useful when there is an increase in variability of the differences as the magnitude of the measurement increases. Eksborg (1981) proposed such a ratio plot, independently of Bland and

Altman. Dewitte et al. (2002) commented on the reception of this article by saying *‘Strange to say, this report has been overlooked’*.



## Bartko's Ellipse

As an enhancement on the Bland Altman Plot, Bartko (1994) has expounded a confidence ellipse for the covariates. Bartko (1994) proposes a bivariate confidence ellipse as a boundary for dispersion. The stated purpose is to ‘amplify dispersion’, which presumably is for the purposes of outlier detection. The orientation of the the ellipse is key to interpreting the results. The minor axis is related to the between-item variability whereas the major axis is related to the mean squared error (referred to here as Error Mean Square). The ellipse illustrates the size of both relative to each other.

Consequently Bartko's ellipse provides a visual aid to determining the relationship between variances. Furthermore, the ellipse provides a visual aid to determining the relationship between the variance of the means  $Var(a_i)$  and the variance of the differences  $Var(d_i)$ . If  $var(a)$  is greater than  $var(d)$ , the orientation of the ellipse is horizontal. Conversely if  $var(a)$  is less than  $var(d)$ , the orientation of the ellipse is vertical. The more horizontal the ellipse, the greater the degree of agreement between the two methods being tested.

Bartko states that the ellipse can, inter alia, be used to detect the presence of outliers (furthermore Bartko (1994) proposes formal testing procedures, that shall be discussed in due course). The Bland-Altman plot for the Grubbs data, complemented by Bartko's ellipse, is depicted in Figure 2.3.12. The fourth observation is shown to be outside the bounds of the ellipse, indicating that it is a potential outlier.

The limitations of using bivariate approaches to outlier detection in the Bland-Altman plot can demonstrated using Bartko's ellipse. A covariate is added to the ‘F vs C’ comparison that has a difference value equal to the inter-method bias, and an average value that markedly deviates from the rest of the average values in the comparison, i.e. 786. Table 1.8 depicts a 95% confidence ellipse for this manipulated data set. By inspection of the confidence interval, a conclusion would be reached that this extra covariate is an outlier, in spite of the fact that this observation is wholly consistent with the conclusion of the Bland-Altman plot.




Figure 2.3.12: Bartko's Ellipse For Grubbs' Data.

Importantly, outlier classification must be informed by the logic of the data's formulation. In the Bland-Altman plot, the horizontal displacement of any observation is supported by two independent measurements. Any observation should not be considered an outlier on the basis of a noticeable horizontal displacement from the main cluster, as in the case with the extra covariate. Conversely, the fourth observation, from the original data set, should be considered an outlier, as it has a noticeable vertical displacement from the rest of the observations.




Figure 2.3.13: Bartko's Ellipse For Grubbs' Data, with an extra covariate.

In the Bland-Altman plot, the horizontal displacement of any point on the plot is supported by two independent measurements. Any point should not be considered an outlier on the basis of a noticeable horizontal displacement from the main cluster, as in the case with the extra co-variate. Conversely, the fourth point, from the original data set, should be considered an outlier, as it has a noticeable vertical displacement from the rest of the observations.

## Survival-Agreement Plot

A graphical technique for method comparison studies, that is entirely different to the Bland-Altman plot, was proposed by Luiz et al. (2003). This approach, known as the survival-agreement plot, is used to determine the degree of agreement using the Kaplan-Meier method, a well known graphical technique in the area of Survival Analysis. Furthermore Luiz et al. (2003) propose that commonly used survival analysis techniques should complement this method, *providing a new analytical insight for agreement*. Two survival-agreement plots are used to detect the bias between to measurements of the same variable. The presence of inter-method bias is tested with the log-rank test, and its magnitude with Cox regression.

The degree of agreement (or disagreement) of a measure is expressed as a function of several limits of tolerance, using the Kaplan-Meier method, where the failures occur exactly at absolute values of the differences between the two methods of measurement.

According to Luiz et al, the survival-agreement plot is a step function of a typical survival analysis without censored data, where the Y axis represents the proportion of discordant cases. This is equivalent to a step function where the X axis represents the absolute observed differences and the Y axis is the proportion of the cases with at least the observed difference ( $x_i$ ).

## Mountain Plot

Krouwer and Monti have proposed a folded empirical cumulative distribution plot, otherwise known as a Mountain plot.

They argue that it is suitable for detecting large, infrequent errors. This is a non-parametric method that can be used as a complement with the Bland Altman plot. Mountain plots are created by computing a percentile for each ranked difference between a new method and a reference method. (Folded plots are so called because of the following transformation is performed for all percentiles above 50: percentile = 100 - percentile.) These percentiles are then plotted against the differences between

the two methods.

Krouwer and Monti argue that the mountain plot offers some following advantages. It is easier to find the central 95% of the data, even when the data are not normally distributed. Also, comparison on different distributions can be performed with ease.

### 2.3.2 Replicate Measurements

Thus far, the formulation for comparison of two measurement methods is based on one measurement by each method per subject. Should there be two or more measurements by each method, these measurements are known as ‘replicate measurements’. Carstensen et al. (2008) recommends the use of replicate measurements, but acknowledges the additional computational complexity.

Bland and Altman (1986) address this situation via two different approaches. The premise of the first approach is that replicate measurements can be treated as independent measurements. The second approach is based upon using the mean of the each group of replicates as one single representative value.

Although either approach may be used to estimate the inter-method bias, removal of the effects of replicate measurements error leads to the underestimation of the standard deviation of the differences. Bland and Altman (1986) propose a correction for this.

Carstensen et al. (2008) take issue with the limits of agreement based on mean values of replicate measurements, since these must be interpreted as prediction limits for the difference between means of repeated measurements by both methods, rather than the difference of individual measurements. Carstensen et al. (2008) demonstrates how the limits of agreement calculated using the mean of replicates are ‘much too narrow as prediction limits for differences between future single measurements’. This paper also comments that, while treating the replicate measurements as independent will cause a downward bias on the limits of agreement calculation, this method is preferable to the ‘mean of replicates’ approach.

### 2.3.3 Prevalence of the Bland-Altman plot

Bland and Altman (1986), which further develops the Bland-Altman methodology, was found to be the sixth most cited paper of all time by the Ryan and Woodall (2005). Dewitte et al. (2002) describes the rate at which prevalence of the Bland-Altman plot has developed in scientific literature. Dewitte et al. (2002) reviewed the use of Bland-Altman plots by examining all articles in the journal ‘Clinical Chemistry’ between 1995 and 2001. This study concluded that use of the Bland-Altman plot increased over the years, from 8% in 1995 to 14% in 1996, and 31-36% in 2002.

The Bland-Altman Plot has since become expected, and often obligatory, approach for presenting method comparison studies in many scientific journals (Hollis, 1996). Furthermore O’Brien et al. (1990) recommend its use in papers pertaining to method comparison studies for the journal of the British Hypertension Society.

## 2.4 The Technology Acceptance Model

Davis (1989) proposes the TAM model, which suggests an hypothesis as to why users may adopt particular technologies, and not others. According to this theory, when users are presented with a new technology, two important factors will influence their decision about how and when they will adopt it.

**Perceived usefulness (PU)** - This was defined by Fred Davis as "the degree to which a person believes that using a particular system would enhance his or her job performance".

**Perceived ease-of-use (PEOU)** - Davis defined this as "the degree to which a person believes that using a particular system would be free from effort"

Davis’s explanations of these term can be rephrased for application to statistical analysis. Perceived Use could refer to the degree to which an user would deem a particular statistical method would properly establish the results of an analysis. In the case of method comparison studies, proper indication of agreement, or lack thereof.

Perceived ease-of-use requires only applying the context of a statistical problem. A very modest statistical skill set is the only prerequisite for constructing a Bland-Altman plot, and computing limits of agreement. The main building blocks are simple descriptive statistics and a knowledge of the normal distribution. These are topics that feature in almost every undergraduate statistics courses.

In short, the user perceives the Bland-Altman methodology to be an easy-to-implement technique, that will properly address the question of agreement.

Conversely the Survival plot is a derivative of the Kaplan-Meier Curve, a non-parametric graphical technique that features in Survival Analysis. This subject area is a well known domain of statistics, but would be encountered on curriculums of specialist courses.

The Mountain Plot is formally called the empirical folder cumulative distribution plot. While not particularly hard to render, the procedure is not straight-forward for the casual user. Currently there is only one software implementation , *medcalc.be* toolkit.

## 2.5 Formal Models and Tests

While the Bland-Altman plot is useful for inspection of data, Kinsella (1986) notes the lack of formal testing offered by this methodology. Furthermore, Kinsella (1986) formulates a model for single measurement observations as a linear mixed effects model, i.e. a model that additively combines fixed effects and random effects:

$$Y_{ij} = \mu + \beta_j + u_i + \epsilon_{ij} \quad i = 1, \dots, n \quad j = 1, 2$$

The true value of the measurement is represented by  $\mu$  while the fixed effect due to method  $j$  is  $\beta_j$ . For simplicity these terms can be combined into single terms;  $\mu_1 = \mu + \beta_1$  and  $\mu_2 = \mu + \beta_2$ . The inter-method bias is the difference of the two fixed effect terms,  $\beta_1 - \beta_2$ . Each individual is assumed to give rise to a random error, represented by  $u_i$ . This random effects term is assumed to have mean zero and be normally distributed with variance  $\sigma^2$ . There is assumed to be an attendant error for each measurement on each individual, denoted  $\epsilon_{ij}$ . This is also assumed to have mean zero. The variance of measurement error for both methods are not assumed to be identical for both methods variance, hence it is denoted  $\sigma_j^2$ . The set of observations  $(x_i, y_i)$  by methods  $X$  and  $Y$  are assumed to follow a bivariate normal distribution with expected values  $E(x_i) = \mu_i$  and  $E(y_i) = \tau_i$  respectively. The variance covariance of the observations  $\Sigma$  is given by

$$\Sigma = \begin{bmatrix} \sigma^2 + \sigma_1^2 & \sigma^2 \\ \sigma^2 & \sigma^2 + \sigma_2^2 \end{bmatrix}$$

Kinsella (1986) demonstrates the estimation of the variance terms and relative precisions relevant to a method comparison study, with attendant confidence intervals for both. The measurement model introduced by Grubbs (1948, 1973) provides a formal procedure for estimating the variances  $\sigma^2$ ,  $\sigma_1^2$  and  $\sigma_2^2$ . Grubbs (1948) offers estimates, commonly known as Grubbs estimators, for the various variance components. These



estimates are maximum likelihood estimates, which shall be revisited in due course.

$$\begin{aligned}\hat{\sigma}^2 &= \sum \frac{(x_i - \bar{x})(y_i - \bar{y})}{n-1} = S_{xy} \\ \hat{\sigma}_1^2 &= \sum \frac{(x_i - \bar{x})^2}{n-1} = S^2_x - S_{xy} \\ \hat{\sigma}_2^2 &= \sum \frac{(y_i - \bar{y})^2}{n-1} = S^2_y - S_{xy}\end{aligned}$$

Thompson (1963) defines  $\Delta_j = \sigma^2/\sigma_j^2, j = 1, 2$ , to be a measure of the relative precision of the measurement methods, and demonstrates how to make statistical inferences about  $\Delta_j$ . Based on the following identities,

$$\begin{aligned}C_x &= (n-1)S_x^2, \\ C_{xy} &= (n-1)S_{xy}, \\ C_y &= (n-1)S_y^2, \\ |A| &= C_x \times C_y - (C_{xy})^2,\end{aligned}$$

the confidence interval limits of  $\Delta_1$  are

$$\frac{C_{xy} - t(\frac{|A|}{n-2})^{\frac{1}{2}}}{C_x - C_{xy} + t(\frac{|A|}{n-2})^{\frac{1}{2}}} < \Delta_1 < \frac{C_{xy} + t(\frac{|A|}{n-2})^{\frac{1}{2}}}{C_x - C_{xy} - t(\frac{|A|}{n-2})^{\frac{1}{2}}}$$

The value  $t$  is the  $100(1 - \alpha/2)\%$  upper quantile of Student's  $t$  distribution with  $n - 2$  degrees of freedom (Kinsella, 1986). The confidence limits for  $\Delta_2$  are found by substituting  $C_y$  for  $C_x$  in (1.2). Negative lower limits are replaced by the value 0.

The case-wise differences and means are calculated as  $d_i = x_i - y_i$  and  $a_i = (x_i + y_i)/2$  respectively. Both  $d_i$  and  $a_i$  are assumed to follow a bivariate normal distribution with  $E(d_i) = \mu_d = \mu_1 - \mu_2$  and  $E(a_i) = \mu_a = (\mu_1 + \mu_2)/2$ , and the variance matrix  $\Sigma_{(a,d)}$  is

$$\Sigma_{(a,d)} = \begin{bmatrix} \sigma_1^2 + \sigma_2^2 & \frac{1}{2}(\sigma_1^2 - \sigma_2^2) \\ \frac{1}{2}(\sigma_1^2 - \sigma_2^2) & \sigma^2 + \frac{1}{4}(\sigma_1^2 + \sigma_2^2) \end{bmatrix}. \quad (2.2)$$

### 2.5.1 Morgan Pitman Testing

An early contribution to formal testing in method comparison was made by both Morgan (1939) and Pitman (1939), in separate contributions. The basis of this approach is that the distribution of the original measurements is bivariate normal. Morgan and Pitman noted that the correlation coefficient depends upon the difference  $\sigma_1^2 - \sigma_2^2$ , being zero if and only if  $\sigma_1^2 = \sigma_2^2$ .

The classical Pitman-Morgan test is a hypothesis test for equality of the variance of two data sets;  $\sigma_1^2 = \sigma_2^2$ , based on the correlation value  $\rho_{a,d}$ , and is evaluated as follows;

$$\rho(a, d) = \frac{\sigma_1^2 - \sigma_2^2}{\sqrt{(\sigma_1^2 + \sigma_2^2)(4\sigma_S^2 + \sigma_1^2 + \sigma_2^2)}} \quad (2.3)$$

The correlation constant takes the value zero if, and only if, the two variances are equal. Therefore a test of the hypothesis  $H : \sigma_1^2 = \sigma_2^2$  is equivalent to a test of the hypothesis  $H : \rho(D, A) = 0$ . This corresponds to the well-known  $t$  test for a correlation coefficient with  $n - 2$  degrees of freedom. Bartko (1994) describes the Morgan-Pitman test as identical to the test of the slope equal to zero in the regression of  $Y_{i1}$  on  $Y_{i2}$ , a result that can be derived using straightforward algebra.

### 2.5.2 Paired sample $t$ test

Bartko (1994) discusses the use of the well known paired sample  $t$  test to test for inter-method bias;  $H : \mu_d = 0$ . The test statistic is distributed a  $t$  random variable with  $n - 1$  degrees of freedom and is calculated as follows,

$$t^* = \frac{\bar{d}}{\frac{s_d}{\sqrt{n}}} \quad (2.4)$$

where  $\bar{d}$  and  $s_d$  is the average of the differences of the  $n$  observations. Only if the two methods show comparable precision then the paired sample student  $t$ -test is appropriate for assessing the magnitude of the bias.

## 2.6 Blackwood Bradley Model

Bradley and Blackwood (1989) have developed a regression based procedure for assessing the agreement. This approach performs a simultaneous test for the equivalence of means and variances of the respective methods. Using simple linear regression of the differences of each pair against the sums, a line is fitted to the model, with estimates for intercept and slope ( $\hat{\beta}_0$  and  $\hat{\beta}_1$ ).

$$D = (X_1 - X_2) \quad (2.5)$$

$$M = (X_1 + X_2)/2 \quad (2.6)$$

The Bradley Blackwood Procedure fits D on M as follows:

$$D = \beta_0 + \beta_1 M \quad (2.7)$$

This technique offers a formal simultaneous hypothesis test for the mean and variance of two paired data sets. The null hypothesis of this test is that the mean ( $\mu$ ) and variance ( $\sigma^2$ ) of both data sets are equal if the slope and intercept estimates are equal to zero (i.e.  $\sigma_1^2 = \sigma_2^2$  and  $\mu_1 = \mu_2$  if and only if  $\beta_0 = \beta_1 = 0$  )

A test statistic is then calculated from the regression analysis of variance values (Bradley and Blackwood, 1989) and is distributed as ‘ $F$ ’ random variable. The degrees of freedom are  $\nu_1 = 2$  and  $\nu_2 = n - 2$  (where  $n$  is the number of pairs). The critical value is chosen for  $\alpha\%$  significance with those same degrees of freedom. Bartko (1994) amends this approach for use in method comparison studies, using the averages of the pairs, as opposed to the sums, and their differences. This approach can facilitate simultaneous usage of test with the Bland-Altman approach. Bartko’s test statistic take the form:

$$F.test = \frac{(\Sigma d^2) - SSReg}{2MSReg}$$

For the Grubbs data,  $\Sigma d^2 = 5.09$ ,  $SSReg = 0.60$  and  $MSreg = 0.06$  Therefore the test statistic is 37.42, with a critical value of 4.10. Hence the means and variance of

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Averages	1	0.04	0.04	0.74	0.4097
Residuals	10	0.60	0.06		

Table 2.6.5: Regression ANOVA of case-wise differences and averages for Grubbs Data

the Fotobalk and Counter chronometers are assumed to be simultaneously equal.

Importantly, this approach determines whether there is both inter-method bias and precision present, or alternatively if there is neither present. It has previously been demonstrated that there is a inter-method bias present, but as this procedure does not allow for separate testing, no conclusion can be drawn on the comparative precision of both methods.

### 2.6.1 Bland-Altman correlation test

The approach proposed by Altman and Bland (1983) is a formal test on the Pearson correlation coefficient of case-wise differences and means ( $\rho_{AD}$ ). According to the authors, this test is equivalent to the ‘Pitman Morgan Test’. For the Grubbs data, the correlation coefficient estimate ( $r_{AD}$ ) is 0.2625, with a 95% confidence interval of (-0.366, 0.726) estimated by Fishers ‘ $r$  to  $z$ ’ transformation (Cohen, Cohen, West, and Aiken, Cohen et al.). The null hypothesis ( $\rho_{AD} = 0$ ) fail to be rejected. Consequently the null hypothesis of equal variances of each method would also fail to be rejected. There has no been no further mention of this particular test in Bland and Altman (1986), although Bland and Altman (1999) refers to Spearman’s rank correlation coefficient. Bland and Altman (1999) state that they ‘do not see a place for methods of analysis based on hypothesis testing’. Bland and Altman (1999) also states that consider structural equation models to be inappropriate.

## 2.6.2 Identifiability

? highlights an important issue regarding using models such as structural equation modelling, which is the identifiability problem. This comes as a result of there being too many parameters to be estimated. Therefore assumptions about some parameters, or estimators used, must be made so that others can be estimated. For example, in the literature, the variance ratio  $\lambda = \frac{\sigma_1^2}{\sigma_2^2}$  must often be assumed to be equal to 1 (Linnet, 1998). ? considers approaches based on two methods with single measurements on each subject as inadequate for a serious study on the measurement characteristics of the methods. This is because there would not be enough data to allow for a meaningful analysis. There is, however, a counter-argument that in many practical settings it is very difficult to get replicate observations when, for example, the measurement method requires invasive medical procedure.

Bradley and Blackwood (1989) offer a formal simultaneous hypothesis test for the mean and variance of paired data sets. This approach is based upon regressing the differences of each pair on the sum of each pair, a line is fitted to the model, with estimates for intercept and slope ( $\hat{\beta}_0$  and  $\hat{\beta}_1$ ). The null hypothesis of this test is that the mean ( $\mu$ ) and variance ( $\sigma^2$ ) of both data sets are equal if the slope and intercept estimates are equal to zero (i.e  $\sigma_1^2 = \sigma_2^2$  and  $\mu_1 = \mu_2$  if and only if  $\beta_0 = \beta_1 = 0$  )

A test statistic is then calculated from the regression analysis of variance values (Bradley and Blackwood, 1989) and is distributed as ‘ $F$ ’ random variable. The degrees of freedom are  $\nu_1 = 2$  and  $\nu_2 = n - 2$  (where  $n$  is the number of pairs). Bartko (1994) amends this approach for use in method comparison studies, using the averages of the pairs, as opposed to the sums, and their differences. This approach can facilitate simultaneous usage of test with the Bland-Altman approach. Bartko’s test statistic take the form:

$$F.test = \frac{(\Sigma d^2) - SSReg}{2MSReg}$$

For the Grubbs data,  $\Sigma d^2 = 5.09$ ,  $SSReg = 0.60$  and  $MSreg = 0.06$ . Therefore

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Averages	1	0.04	0.04	0.74	0.4097
Residuals	10	0.60	0.06		

Table 2.6.6: Regression ANOVA of case-wise differences and averages for Grubbs Data

the test statistic is 3.742, with a critical value of 4.10. Hence the means and variance of the Fotobalk and Counter chronometers are assumed to be simultaneously equal.

Importantly, this methodology determines whether there is both inter-method bias and precision present, or alternatively if there is neither present. It has previously been demonstrated that there is a inter-method bias present, but as this procedure does not allow for separate testing, no conclusion can be drawn on the comparative precision of both methods.

## 2.7 Regression Methods for Method Comparison

Conventional regression models are estimated using the ordinary least squares (OLS) technique, and are referred to as ‘Model I regression’ (Cornbleet and Cochrane, 1979; Ludbrook, 1997). A key feature of Model I models is that the independent variable is assumed to be measured without error. However this assumption invalidates simple linear regression for use in method comparison studies, as both methods must be assumed to be measured with error (Altman and Bland, 1983; Ludbrook, 1997).

The use of regression models that assumes the presence of error in both variables  $X$  and  $Y$  have been proposed for use instead (Cornbleet and Cochrane, 1979; Ludbrook, 1997). These methodologies are collectively known as ‘Model II regression’. They differ in the method used to estimate the parameters of the regression.

Regression estimates depend on formulation of the model. A formulation with one method considered as the  $X$  variable will yield different estimates for a formulation where it is the  $Y$  variable. With Model I regression, the models fitted in both cases

will entirely different and inconsistent. However with Model II regression, they will be consistent and complementary.

Regression approaches are useful for a making a detailed examination of the biases across the range of measurements, allowing bias to be decomposed into fixed bias and proportional bias. Fixed bias describes the case where one method gives values that are consistently different to the other across the whole range. Proportional bias describes the difference in measurements getting progressively greater, or smaller, across the range of measurements. A measurement method may have either an attendant fixed bias or proportional bias, or both. (Ludbrook, 1997). Determination of these biases shall be discussed in due course.

## 2.8 Regression Methods (duplication)

Conventional regression models are estimated using the ordinary least squares (OLS) technique, and are referred to as ‘Model I regression’ (Cornbleet and Cochrane, 1979; Ludbrook, 1997). A key feature of Model I models is that the independent variable is assumed to be measured without error. As often pointed out in several papers (Altman and Bland, 1983; Ludbrook, 1997), this assumption invalidates simple linear regression for use in method comparison studies, as both methods must be assumed to be measured with error.

The use of regression models that assumes the presence of error in both variables  $X$  and  $Y$  have been proposed for use instead (Cornbleet and Cochrane, 1979; Ludbrook, 1997). These approaches are collectively known as ‘Model II regression’. They differ in the method used to estimate the parameters of the regression.

Regression estimates depend on formulation of the model. A formulation with one method considered as the  $X$  variable will yield different estimates for a formulation where it is the  $Y$  variable. With Model I regression, the models fitted in both cases will entirely different and inconsistent. However with Model II regression, they will be consistent and complementary.

Regression approaches are useful for making a detailed examination of the biases across the range of measurements, allowing bias to be decomposed into fixed bias and proportional bias. Fixed bias describes the case where one method gives values that are consistently different to the other across the whole range. Proportional bias describes the difference in measurements getting progressively greater, or smaller, across the range of measurements. A measurement method may have either an attendant fixed bias or proportional bias, or both. (?). Determination of these biases shall be discussed in due course.



### 2.8.1 Deming Regression

As stated previously, the fundamental flaw of simple linear regression is that it allows for measurement error in one variable only. This causes a downward biased slope estimate.

Deming regression is a regression fitting approach that assumes error in both variables. Deming regression is recommended by Cornbleet and Cochrane (1979) as the preferred Model II regression for use in method comparison studies. The sum of squared distances from measured sets of values to the regression line is minimized at an angles specified by the ratio  $\lambda$  of the residual variance of both variables. I When  $\lambda$  is one, the angle is 45 degrees. In ordinary linear regression, the distances are minimized in the vertical directions (Linnet, 1999). In cases involving only single measurements by each method,  $\lambda$  may be unknown and is therefore assumes a value of one. While this will produce biased estimates, they are less biased than ordinary linear regression.

The Bland Altman Plot is uninformative about the comparative influence of proportional bias and fixed bias. Model II approaches, such as Deming regression, can provide independent tests for both types of bias.

For a given  $\lambda$ , Kummel (1879) derived the following estimate that would later be used for the Deming regression slope parameter. The intercept estimate  $\alpha$  is simply estimated in the same way as in conventional linear regression, by using the identity  $\bar{Y} - \hat{\beta}\bar{X}$ ;

$$\hat{\beta} = \frac{S_{yy} - \lambda S_{xx} + [(S_{yy} - \lambda S_{xx})^2 + 4\lambda S_{xy}^2]^{1/2}}{2S_{xy}} \quad (2.8)$$

, with  $\lambda$  as the variance ratio. As stated previously  $\lambda$  is often unknown, and therefore must be assumed to equal one. Carroll and Ruppert (1996) states that Deming regression is acceptable only when the precision ratio ( $\lambda$ , in their paper as  $\eta$ ) is correctly specified, but in practice this is often not the case, with the  $\lambda$  being underestimated. Several candidate models, with varying variance ratios may be fitted, and estimates of the slope and intercept are produced. However no model selection information is available to determine the best fitting model.

As with conventional regression methodologies, Deming regression calculates an estimate for both the slope and intercept for the fitted line, and standard errors thereof. Therefore there is sufficient information to carry out hypothesis tests on both estimates, that are informative about presence of fixed and proportional bias.

A 95% confidence interval for the intercept estimate can be used to test the intercept, and hence fixed bias, is equal to zero. This hypothesis is accepted if the confidence interval for the estimate contains the value 0 in its range. Should this be, it can be concluded that fixed bias is not present. Conversely, if the hypothesis is rejected, then it is concluded that the intercept is non zero, and that fixed bias is present.

Testing for proportional bias is a very similar procedure. The 95% confidence interval for the slope estimate can be used to test the hypothesis that the slope is equal to 1. This hypothesis is accepted if the confidence interval for the estimate contains the value 1 in its range. If the hypothesis is rejected, then it is concluded that the slope is significant different from 1 and that a proportional bias exists.

For convenience, a new data set shall be introduced to demonstrate Deming regression. Measurements of transmitral volumetric flow (MF) by doppler echocardiography, and left ventricular stroke volume (SV) by cross sectional echocardiography in 21 patients with aortic valve disease are tabulated in Zhang et al. (1986). This data set features in the discussion of method comparison studies in Altman (1991, p.398) .

Carroll and Ruppert (1996) states that Deming's regression is acceptable only when the precision ratio ( $\lambda$ , in their paper as  $\eta$ ) is correctly specified, but in practice this is often not the case, with the  $\lambda$  being underestimated.

Patient	MF ( $cm^3$ )	SV ( $cm^3$ )	Patient	MF ( $cm^3$ )	SV ( $cm^3$ )	Patient	MF ( $cm^3$ )	SV ( $cm^3$ )
1	47	43	8	75	72	15	90	82
2	66	70	9	79	92	16	100	100
3	68	72	10	81	76	17	104	94
4	69	81	11	85	85	18	105	98
5	70	60	12	87	82	19	112	108
6	70	67	13	87	90	20	120	131
7	73	72	14	87	96	21	132	131

Table 2.8.7: Transmitral volumetric flow(MF) and left ventricular stroke volume (SV) in 21 patients. (Zhang et al 1986)

## 2.9 Other Types of Studies

Lewis et al. (1991) categorize method comparison studies into three different types. The key difference between the first two is whether or not a ‘gold standard’ method is used. In situations where one instrument or method is known to be ‘accurate and precise’, it is considered as the ‘gold standard’ (Lewis et al., 1991). A method that is not considered to be a gold standard is referred to as an ‘approximate method’. In calibration studies they are referred to a criterion methods and test methods respectively.

**1. Calibration problems.** The purpose is to establish a relationship between methods, one of which is an approximate method, the other a gold standard. The results of the approximate method can be mapped to a known probability distribution of the results of the gold standard (Lewis et al., 1991). (In such studies, the gold standard method and corresponding approximate method are generally referred to a criterion method and test method respectively.) Altman and Bland (1983) make clear that their methodology is not intended for calibration problems.

**2. Comparison problems.** When two approximate methods, that use the same




Figure 2.8.14: Deming Regression For Zhang's Data

units of measurement, are to be compared. This is the case which the Bland-Altman methodology is specifically intended for, and therefore it is the most relevant of the three.

**3. Conversion problems.** When two approximate methods, that use different units of measurement, are to be compared. This situation would arise when the measurement methods use 'different proxies', i.e different mechanisms of measurement. Lewis et al. (1991) deals specifically with this issue. In the context of this study, it is the least relevant of the three.

?, p.47 cautions that 'gold standards' should not be assumed to be error free. 'It is

of necessity a subjective decision when we come to decide that a particular method or instrument can be treated as if it was a gold standard'. The clinician gold standard, the sphygmomanometer, is used as an example thereof. The sphygmomanometer 'leaves considerable room for improvement' (?). Pizzi (1999) similarly addresses the issue of gold standards, 'well-established gold standard may itself be imprecise or even unreliable'.

The NIST F1 Caesium fountain atomic clock is considered to be the gold standard when measuring time, and is the primary time and frequency standard for the United States. The NIST F1 is accurate to within one second per 60 million years (NIST, 2009).

Measurements of the interior of the human body are, by definition, invasive medical procedures. The design of method must balance the need for accuracy of measurement with the well-being of the patient. This will inevitably lead to the measurement error as described by ?. The magnetic resonance angiogram, used to measure internal anatomy, is considered to the gold standard for measuring aortic dissection. Medical test based upon the angiogram is reported to have a false positive reporting rate of 5% and a false negative reporting rate of 8%. This is reported as sensitivity of 95% and a specificity of 92% (ACR, 2008).

In literature they are, perhaps more accurately, referred to as 'fuzzy gold standards' (Phelps and Hutson, 1995). Consequently when one of the methods is essentially a fuzzy gold standard, as opposed to a 'true' gold standard, the comparison of the criterion and test methods should be consider in the context of a comparison study, as well as of a calibration study.

# Chapter 3

## Linear Mixed effects Models

### 3.1 Linear Mixed effects Models

A linear mixed effects (LME) model is a statistical model containing both fixed effects and random effects (random effects are also known as variance components). LME models are a generalization of the classical linear model, which contain fixed effects only. When the levels of factors are considered to be sampled from a population, and each level is not of particular interest, they are considered random quantities with associated variances. The effects of the levels, as described, are known as random effects. Random effects are represented by unobservable normally distributed random variables. Conversely fixed effects are considered non-random and the levels of each factor are of specific interest.

Fisher (1918) introduced variance components models for use in genetical studies. Whereas an estimate for variance must take a non-negative value, an individual variance component, i.e. a component of the overall variance, may be negative.

The framework has developed since, including contributions from Tippett (1931), who extend the use of variance components into linear models, and Eisenhart (1947), who introduced the ‘mixed model’ terminology and formally distinguished between mixed and random effects models. Henderson (1950) devised a framework for deriving

estimates for both the fixed effects and the random effects, using a set of equations that would become known as ‘mixed model equations’ or ‘Henderson’s equations’. LME methodology is further enhanced by Henderson’s later works (Henderson, 1953; Henderson et al., 1959, 1963, 1973, 1984). The key features of Henderson’s work provide the basis for the estimation techniques.

Hartley and Rao (1967) demonstrated that unique estimates of the variance components could be obtained using maximum likelihood methods. However these estimates are known to be biased ‘downwards’ (i.e. underestimated) , because of the assumption that the fixed estimates are known, rather than being estimated from the data. Patterson and Thompson (1971) produced an alternative set of estimates, known as the restricted maximum likelihood (REML) estimates, that do not require the fixed effects to be known. Thusly there is a distinction the REML estimates and the original estimates, now commonly referred to as ML estimates.

Laird and Ware (1982) provides a form of notation for notation for LME models that has since become the standard form, or the basis for more complex formulations. Due to computation complexity, linear mixed effects models have not seen widespread use until many well known statistical software applications began facilitating them. SAS Institute added PROC MIXED to its software suite in 1992 (Singer, 1998). Pinheiro and Bates (1994) described how to compute LME models in the **S-plus** environment.

Using Laird-Ware form, the LME model is commonly described in matrix form,

$$y = X\beta + Zb + \epsilon \tag{3.1}$$

where  $y$  is a vector of  $N$  observable random variables,  $\beta$  is a vector of  $p$  fixed effects,  $X$  and  $Z$  are  $N \times p$  and  $N \times q$  known matrices, and  $b$  and  $\epsilon$  are vectors of  $q$  and  $N$ , respectively, random effects such that  $E(b) = 0$ ,  $E(\epsilon) = 0$  and

$$\text{var} \begin{pmatrix} b \\ \epsilon \end{pmatrix} = \begin{pmatrix} D & 0 \\ 0 & \Sigma \end{pmatrix}$$

where  $D$  and  $\Sigma$  are positive definite matrices parameterized by an unknown variance component parameter vector  $\theta$ . The variance-covariance matrix for the vector of

observations  $y$  is given by  $V = ZDZ' + \Sigma$ . This implies  $y \sim (X\beta, V) = (X\beta, ZDZ' + \Sigma)$ . It is worth noting that  $V$  is an  $n \times n$  matrix, as the dimensionality becomes relevant later on. The notation provided here is generic, and will be adapted to accord with complex formulations that will be encountered in due course.

### 3.1.1 Estimation

Estimation of LME models involve two complementary estimation issues'; estimating the vectors of the fixed and random effects estimates  $\hat{\beta}$  and  $\hat{b}$  and estimating the variance covariance matrices  $D$  and  $\Sigma$ . Inference about fixed effects have become known as 'estimates', while inferences about random effects have become known as 'predictions'. The most common approach to obtain estimators are Best Linear Unbiased Estimator (BLUE) and Best Linear Unbiased Predictor (BLUP). For an LME model given by (3.1), the BLUE of  $\hat{\beta}$  is given by

$$\hat{\beta} = (X'V^{-1}X)^{-1}X'V^{-1}y,$$

whereas the BLUP of  $\hat{b}$  is given by

$$\hat{b} = DZ'V^{-1}(y - X\hat{\beta}).$$

### Henderson's equations

Because of the dimensionality of  $V$  (i.e.  $n \times n$ ) computing the inverse of  $V$  can be difficult. As a way around this problem Henderson (1953); Henderson et al. (1959, 1963, 1973, 1984) offered a more simpler approach of jointly estimating  $\hat{\beta}$  and  $\hat{b}$ . Henderson (1950) made the (ad-hoc) distributional assumptions  $y|b \sim N(X\beta + Zb, \Sigma)$  and  $b \sim N(0, D)$ , and proceeded to maximize the joint density of  $y$  and  $b$

$$\left| \begin{array}{cc} D & 0 \\ 0 & \Sigma \end{array} \right|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} \begin{pmatrix} b \\ y - X\beta - Zb \end{pmatrix}' \begin{pmatrix} D & 0 \\ 0 & \Sigma \end{pmatrix}^{-1} \begin{pmatrix} b \\ y - X\beta - Zb \end{pmatrix} \right\}, \quad (3.2)$$

with respect to  $\beta$  and  $b$ , which ultimately requires minimizing the criterion

$$(y - X\beta - Zb)' \Sigma^{-1} (y - X\beta - Zb) + b' D^{-1} b. \quad (3.3)$$



This leads to the mixed model equations

$$\begin{pmatrix} X'\Sigma^{-1}X & X'\Sigma^{-1}Z \\ Z'\Sigma^{-1}X & X'\Sigma^{-1}X + D^{-1} \end{pmatrix} \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} X'\Sigma^{-1}y \\ Z'\Sigma^{-1}y \end{pmatrix}. \quad (3.4)$$

Using these equations, obtaining the estimates requires the inversion of a matrix of dimension  $p + q \times p + q$ , considerably smaller in size than  $V$ . ? shows that these mixed model equations do not depend on normality and that  $\hat{\beta}$  and  $\hat{b}$  are the BLUE and BLUP under general conditions, provided  $D$  and  $\Sigma$  are known.

Robinson (1991) points out that although Henderson (1950) initially referred to the estimates  $\hat{\beta}$  and  $\hat{b}$  from (3.4) as “joint maximum likelihood estimates”, Henderson (1973) later advised that these estimates should not be referred to as “maximum likelihood” as the function being maximized in (3.3) is a joint density rather than a likelihood function. Lee et al. (2006) remarks that it is clear that Henderson used joint estimation for computational purposes, without recognizing the theoretical implications.

### Estimation of the fixed parameters

The vector  $y$  has marginal density  $y \sim N(X\beta, V)$ , where  $V = \Sigma + ZDZ'$  is specified through the variance component parameters  $\theta$ . The log-likelihood of the fixed parameters  $(\beta, \theta)$  is

$$\ell(\beta, \theta | y) = -\frac{1}{2} \log |V| - \frac{1}{2} (y - X\beta)' V^{-1} (y - X\beta), \quad (3.5)$$

and for fixed  $\theta$  the estimate  $\hat{\beta}$  of  $\beta$  is obtained as the solution of

$$(X'V^{-1}X)\beta = X'V^{-1}y. \quad (3.6)$$

Substituting  $\hat{\beta}$  from (3.6) into  $\ell(\beta, \theta | y)$  from (3.5) returns the *profile* log-likelihood

$$\begin{aligned} \ell_P(\theta | y) &= \ell(\hat{\beta}, \theta | y) \\ &= -\frac{1}{2} \log |V| - \frac{1}{2} (y - X\hat{\beta})' V^{-1} (y - X\hat{\beta}) \end{aligned}$$

of the variance parameter  $\theta$ . Estimates of the parameters  $\theta$  specifying  $V$  can be found by maximizing  $\ell_P(\theta | y)$  over  $\theta$ . These are the ML estimates.

For REML estimation the *restricted* log-likelihood is defined as

$$\ell_R(\theta \mid y) = \ell_P(\theta \mid y) - \frac{1}{2} \log |X'VX|.$$

The REML approach does not base estimates on a maximum likelihood fit of all the information, but instead uses a likelihood function derived from a data set, transformed to remove the irrelevant influences (Dodge, 2003). Restricted maximum likelihood is often preferred to maximum likelihood because REML estimation reduces the bias in the variance component by taking into account the loss of degrees of freedom that results from estimating the fixed effects in  $\beta$ . Restricted maximum likelihood also handles high correlations more effectively, and is less sensitive to outliers than maximum likelihood. The problem with REML for model building is that the likelihoods obtained for different fixed effects are not comparable. Hence it is not valid to compare models with different fixed effects using a likelihood ratio test or AIC when REML is used to estimate the model. Therefore models derived using ML must be used instead.

### **Estimation of the random effects**

The established approach for estimating the random effects is to use the best linear predictor of  $b$  from  $y$ , which for a given  $\beta$  equals  $DZ'V^{-1}(y - X\beta)$ . In practice  $\beta$  is replaced by an estimator such as  $\hat{\beta}$  from (3.6) so that  $\hat{b} = DZ'V^{-1}(y - X\hat{\beta})$ . Pre-multiplying by the appropriate matrices it is straightforward to show that these estimates  $\hat{\beta}$  and  $\hat{b}$  satisfy the equations in (3.4).

### **Algorithms for likelihood function optimization**

Iterative numerical techniques are used to optimize the log-likelihood function and estimate the covariance parameters  $\theta$ . The procedure is subject to the constraint that  $R$  and  $D$  are both positive definite. The most common iterative algorithms for optimizing the likelihood function are the Newton-Raphson method, which is the preferred method, the expectation maximization (EM) algorithm and the Fisher scoring methods.

The EM algorithm, introduced by Dempster et al. (1977), is an iterative technique for maximizing complicated likelihood functions. The algorithm alternates between performing an expectation (E) step and the maximization (M) step. The ‘E’ step computes the expectation of the log-likelihood evaluated using the current estimate for the variables. In the ‘M’ step, parameters that maximize the expected log-likelihood, found on the previous ‘E’ step, are computed. These parameter estimates are then used to determine the distribution of the variables in the next ‘E’ step. The algorithm alternates between these two steps until convergence is reached.

The main drawback of the EM algorithm is its slow rate of convergence. Consequently the EM algorithm is rarely used entirely in LME estimation, instead providing an initial set of values that can be passed to other optimization techniques.

The Newton Raphson (NR) method is the most common, and recommended technique for ML and REML estimation. The NR algorithm minimizes an objective function defined as  $-2$  times the log likelihood for the covariance parameters  $\theta$ . At every iteration the NR algorithm requires the calculation of a vector of partial derivatives, known as the gradient, and the second derivative matrix with respect to the covariance parameters. This is known as the observed Hessian matrix. Due to the Hessian matrix, the NR algorithm is more time-consuming, but convergence is reached with fewer iterations compared to the EM algorithm. The Fisher scoring algorithm is a variant of the NR algorithm that is more numerically stable and likely to converge, but not recommended to obtain final estimates.

## The extended likelihood

The desire to have an entirely likelihood-based justification for estimates of random effects, in contrast to Henderson’s equation, has motivated Pawitan (2001, page 429) to define the *extended likelihood*. He remarks “In mixed effects modelling the extended likelihood has been called *h-likelihood* (for hierarchical likelihood) by Lee and Nelder (1996), while in smoothing literature it is known as the *penalized likelihood* (e.g. Green and Silverman 1994).” The extended likelihood can be written  $L(\beta, \theta, b|y) =$

$p(y|b; \beta, \theta)p(b; \theta)$  and adopting the same distributional assumptions used by Henderson (1950) yields the log-likelihood function

$$\begin{aligned} \ell_h(\beta, \theta, b|y) = & -\frac{1}{2} \{ \log |\Sigma| + (y - X\beta - Zb)' \Sigma^{-1} (y - X\beta - Zb) \\ & + \log |D| + b' D^{-1} b \}. \end{aligned}$$

Given  $\theta$ , differentiating with respect to  $\beta$  and  $b$  returns Henderson's equations in (3.4).

### **The LME model as a general linear model**

Henderson's equations in (3.4) can be rewritten  $(T'W^{-1}T)\delta = T'W^{-1}y_a$  using

$$\delta = \begin{pmatrix} \beta \\ b \end{pmatrix}, \quad y_a = \begin{pmatrix} y \\ \psi \end{pmatrix}, \quad T = \begin{pmatrix} X & Z \\ 0 & I \end{pmatrix}, \quad \text{and } W = \begin{pmatrix} \Sigma & 0 \\ 0 & D \end{pmatrix},$$

where Lee et al. (2006) describe  $\psi = 0$  as quasi-data with mean  $E(\psi) = b$ . Their formulation suggests that the joint estimation of the coefficients  $\beta$  and  $b$  of the linear mixed effects model can be derived via a classical augmented general linear model  $y_a = T\delta + \varepsilon$  where  $E(\varepsilon) = 0$  and  $\text{var}(\varepsilon) = W$ , with *both*  $\beta$  and  $b$  appearing as fixed parameters. The usefulness of this reformulation of an LME as a general linear model will be revisited.

## 3.2 Repeated measurements in LME models

In many statistical analyzes, the need to determine parameter estimates where multiple measurements are available on each of a set of variables often arises. Further to Lam et al. (1999), Hamlett et al. (2004) performs an analysis of the correlation of replicate measurements, for two variables of interest, using LME models.

Let  $y_{Aij}$  and  $y_{Bij}$  be the  $j$ th repeated observations of the variables of interest  $A$  and  $B$  taken on the  $i$ th subject. The number of repeated measurements for each variable may differ for each individual. Both variables are measured on each time points. Let  $n_i$  be the number of observations for each variable, hence  $2 \times n_i$  observations in total.

It is assumed that the pair  $y_{Aij}$  and  $y_{Bij}$  follow a bivariate normal distribution.

$$\begin{pmatrix} y_{Aij} \\ y_{Bij} \end{pmatrix} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \text{ where } \boldsymbol{\mu} = \begin{pmatrix} \mu_A \\ \mu_B \end{pmatrix}$$

The matrix  $\boldsymbol{\Sigma}$  represents the variance component matrix between response variables at a given time point  $j$ .

$$\boldsymbol{\Sigma} = \begin{pmatrix} \sigma_A^2 & \sigma_{AB} \\ \sigma_{AB} & \sigma_B^2 \end{pmatrix}$$

$\sigma_A^2$  is the variance of variable  $A$ ,  $\sigma_B^2$  is the variance of variable  $B$  and  $\sigma_{AB}$  is the covariance of the two variable. It is assumed that  $\boldsymbol{\Sigma}$  does not depend on a particular time point, and is the same over all time points.

### 3.2.1 Formulation of the Response Vector

Information of individual  $i$  is recorded in a response vector  $\mathbf{y}_i$ . The response vector is constructed by stacking the response of the 2 responses at the first instance, then the 2 responses at the second instance, and so on. Therefore the response vector is a  $2n_i \times 1$  column vector. The covariance matrix of  $\mathbf{y}_i$  is a  $2n_i \times 2n_i$  positive definite matrix  $\boldsymbol{\Omega}_i$ .

Consider the case where three measurements are taken by both methods  $A$  and  $B$ ,

$\mathbf{y}_i$  is a  $6 \times 1$  random vector describing the  $i$ th subject.

$$\mathbf{y}_i = (y_i^{A1}, y_i^{B1}, y_i^{A2}, y_i^{B2}, y_i^{A3}, y_i^{B3})'$$

The response vector  $\mathbf{y}_i$  can be formulated as an LME model according to Laird-Ware form.

$$\mathbf{y}_i = \mathbf{X}_i\boldsymbol{\beta} + \mathbf{Z}_i\mathbf{b}_i + \boldsymbol{\epsilon}_i$$

$$\mathbf{b}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{D})$$

$$\boldsymbol{\epsilon}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{R}_i)$$

Information on the fixed effects are contained in a three dimensional vector  $\boldsymbol{\beta} = (\beta_0, \beta_1, \beta_2)'$ . For computational purposes  $\beta_2$  is conventionally set to zero. Consequently  $\boldsymbol{\beta}$  is the solutions of the means of the two methods, i.e.  $E(\mathbf{y}_i) = \mathbf{X}_i\boldsymbol{\beta}$ . The variance covariance matrix  $\mathbf{D}$  is a general  $2 \times 2$  matrix, while  $\mathbf{R}_i$  is a  $2n_i \times 2n_i$  matrix.

### 3.2.2 Decomposition of the response covariance matrix

The variance covariance structure can be re-expressed in the following form,

$$\text{Cov}(\mathbf{y}_i) = \boldsymbol{\Omega}_i = \mathbf{Z}_i\mathbf{D}\mathbf{Z}_i' + \mathbf{R}_i.$$

$\mathbf{R}_i$  can be shown to be the Kronecker product of a correlation matrix  $\mathbf{V}$  and  $\boldsymbol{\Lambda}$ . The correlation matrix  $\mathbf{V}$  of the repeated measures on a given response variable is assumed to be the same for all response variables. Both Hamlett et al. (2004) and Lam et al. (1999) use the identity matrix, with dimensions  $n_i \times n_i$  as the formulation for  $\mathbf{V}$ . Roy (2009a) remarks that, with repeated measures, the response for each subject is correlated for each variable, and that such correlation must be taken into account in order to produce a valid inference on correlation estimates. Roy (2006) proposes various correlation structures may be assumed for repeated measure correlations, such as the compound symmetry and autoregressive structures, as alternative to the identity matrix.

However, for the purposes of method comparison studies, the necessary estimates are currently only determinable when the identity matrix is specified, and the results in Roy (2009a) indicate its use.

For the response vector described, Hamlett et al. (2004) presents a detailed covariance matrix. A brief summary shall be presented here only. The overall variance matrix is a  $6 \times 6$  matrix composed of two types of  $2 \times 2$  blocks. Each block represents one separate time of measurement.

$$\mathbf{\Omega}_i = \begin{pmatrix} \mathbf{\Sigma} & \mathbf{D} & \mathbf{D} \\ \mathbf{D} & \mathbf{\Sigma} & \mathbf{D} \\ \mathbf{D} & \mathbf{D} & \mathbf{\Sigma} \end{pmatrix}$$

The diagonal blocks are  $\Sigma$ , as described previously. The  $2 \times 2$  block diagonal matrix in  $\mathbf{\Omega}$  gives  $\Sigma$ .  $\Sigma$  is the sum of the between-subject variability  $\mathbf{D}$  and the within subject variability  $\mathbf{\Lambda}$ .

$\mathbf{\Omega}_i$  can be expressed as

$$\mathbf{\Omega}_i = \mathbf{Z}_i \mathbf{D} \mathbf{Z}_i' + (\mathbf{I}_{n_i} \otimes \mathbf{\Lambda}).$$

The notation  $\text{dim}_{n_i}$  means an  $n_i \times n_i$  diagonal block.

### 3.2.3 Correlation terms

Hamlett et al. (2004) demonstrated how the between-subject and within subject variabilities can be expressed in terms of correlation terms.

$$\mathbf{D} = \begin{pmatrix} \sigma_A^2 \rho_A & \sigma_A \sigma_B \rho_{AB} \delta \\ \sigma_A \sigma_B \rho_{AB} \delta & \sigma_B^2 \rho_B \end{pmatrix}$$

$$\mathbf{\Lambda} = \begin{pmatrix} \sigma_A^2 (1 - \rho_A) & \sigma_{AB} (1 - \delta) \\ \sigma_{AB} (1 - \delta) & \sigma_B^2 (1 - \rho_B) \end{pmatrix}.$$

$\rho_A$  describe the correlations of measurements made by the method  $A$  at different times. Similarly  $\rho_B$  describe the correlation of measurements made by the method  $B$

at different times. Correlations among repeated measures within the same method are known as intra-class correlation coefficients.  $\rho_{AB}$  describes the correlation of measurements taken at the same same time by both methods. The coefficient  $\delta$  is added for when the measurements are taken at different times, and is a constant of less than 1 for linked replicates. This is based on the assumption that linked replicates measurements taken at the same time would have greater correlation than those taken at different times. For unlinked replicates  $\delta$  is simply 1. Hamlett et al. (2004) provides a useful graphical depiction of the role of each correlation coefficients.



### 3.3 Using LME for method comparison

Due to the prevalence of modern statistical software, Carstensen et al. (2008) advocates the adoption of computer based approaches, such as LME models, to method comparison studies. Carstensen et al. (2008) remarks upon ‘by-hand’ approaches advocated in Bland and Altman (1999) discouragingly, describing them as tedious, unnecessary and ‘outdated’. Rather than using the ‘by hand’ methods, estimates for required LME parameters can be read directly from program output. Furthermore, using computer approaches removes constraints associated with ‘by-hand’ approaches, such as the need for the design to be perfectly balanced.

#### 3.3.1 Roy’s Approach

For the purposes of comparing two methods of measurement, Roy (2009a) presents a framework that utilizes linear mixed effects model. This methodology provides for the formal testing of inter-method bias, between-subject variability and within-subject variability of two methods. The formulation contains a Kronecker product covariance structure in a doubly multivariate setup. By doubly multivariate set up, Roy means that the information on each patient or item is multivariate at two levels, the number of methods and number of replicated measurements. Further to Lam et al. (1999), it is assumed that the replicates are linked over time. However it is easy to modify to the unlinked case.

Roy (2009a) sets out three criteria for two methods to be considered in agreement. Firstly that there be no significant bias. Second that there is no difference in the between-subject variabilities, and lastly that there is no significant difference in the within-subject variabilities. Roy further proposes examination of the the overall variability by considering the second and third criteria be examined jointly. Should both the second and third criteria be fulfilled, then the overall variabilities of both methods would be equal.

A formal test for inter-method bias can be implemented by examining the fixed ef-

fects of the model. This is common to well known classical linear model methodologies. The null hypotheses, that both methods have the same mean, which is tested against the alternative hypothesis, that both methods have different means. The inter-method bias and necessary  $t$ -value and  $p$ -value are presented in computer output. A decision on whether the first of Roy's criteria is fulfilled can be based on these values.

Importantly Roy (2009a) further proposes a series of three tests on the variance components of an LME model, which allow decisions on the second and third of Roy's criteria. For these tests, four candidate LME models are constructed. The differences in the models are specifically in how the  $D$  and  $\Lambda$  matrices are constructed, using either an unstructured form or a compound symmetry form. To illustrate these differences, consider a generic matrix  $A$ ,

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}.$$

A symmetric matrix allows the diagonal terms  $a_{11}$  and  $a_{22}$  to differ. The compound symmetry structure requires that both of these terms be equal, i.e  $a_{11} = a_{22}$ .

The first model acts as an alternative hypothesis to be compared against each of three other models, acting as null hypothesis models, successively. The models are compared using the likelihood ratio test. Likelihood ratio tests are a class of tests based on the comparison of the values of the likelihood functions of two candidate models.

### 3.3.2 Likelihood Ratio Tests

The relationship between the respective models presented by Roy (2009a) is known as "nesting". A model A to be nested in the reference model, model B, if Model A is a special case of Model B, or with some specific constraint applied.

A general method for comparing models with a nesting relationship is the likelihood ratio test (LRTs). LRTs are a family of tests used to compare the value of likelihood functions for two models, whose respective formulations define a hypothesis to be tested

(i.e. the nested and reference model). The significance of the likelihood ratio test can be found by comparing the likelihood ratio to the  $\chi^2$  distribution, with the appropriate degrees of freedom.

When testing hypotheses around covariance parameters in an LME model, REML estimation for both models is recommended by West et al. REML estimation can be shown to reduce the bias inherent in ML estimates of covariance parameters (West et al., 2007). Conversely, ? advises that testing hypotheses on fixed-effect parameters should be based on ML estimation, and that using REML would not be appropriate in this context.

LRTs can be used to test hypotheses about covariance parameters or fixed effects parameters in the context of LMEs. The test statistic for the likelihood ratio test is the difference of the log-likelihood functions, multiplied by  $-2$ . The probability distribution of the test statistic is approximated by the  $\chi^2$  distribution with  $(\nu_1 - \nu_2)$  degrees of freedom, where  $\nu_1$  and  $\nu_2$  are the degrees of freedom of models 1 and 2 respectively. Each of these three test shall be examined in more detail shortly.

### 3.3.3 Testing Procedures

Roy's methodology requires the construction of four candidate models. The first candidate model is compared to each of the three other models successively. It is the alternative model in each of the three tests, with the other three models acting as the respective null models.

The probability distribution of the test statistic can be approximated by a chi-square distribution with  $(\nu_1 - \nu_2)$  degrees of freedom, where  $\nu_1$  and  $\nu_2$  are the degrees of freedom of models 1 and 2 respectively.

Likelihood ratio tests are very simple to implement in **R**, simply use the '`anova()`' commands. Sample output will be given for each variability test. The likelihood ratio test is the procedure used to compare the fit of two models. For each candidate model, the ' $-2 \log$  likelihood' ( $M2LL$ ) is computed. The test statistic for each of the three

hypothesis tests is the difference of the  $M2LL$  for each pair of models. If the  $p$ -value in each of the respective tests exceed as significance level chosen by the analyst, then the null model must be rejected.

$$-2 \ln \Lambda_d = [M2LL \text{ under } H_0 \text{ model}] - [M2LL \text{ under } H_A \text{ model}] \quad (3.7)$$

These test statistics follow a chi-square distribution with the degrees of freedom computed as the difference of the LRT degrees of freedom.

$$\nu = [\text{LRT df under } H_0 \text{ model}] - [\text{LRT df under } H_A \text{ model}] \quad (3.8)$$

```
> anova(MCS1,MCS2)
>
>
```

	Model	df	AIC	BIC	logLik	Test	L.Ratio	p-value
MCS1	1	8	4077.5	4111.3	-2030.7			
MCS2	2	7	4075.6	4105.3	-2030.8	1 vs 2	0.15291	0.6958

### 3.3.4 Correlation

In addition to the variability tests, Roy advises that it is preferable that a correlation of greater than 0.82 exist for two methods to be considered interchangeable. However if two methods fulfil all the other conditions for agreement, failure to comply with this one can be overlooked. Indeed Roy demonstrates that placing undue importance to it can lead to incorrect conclusions. Roy (2009a) remarks that current computer implementations only gives overall correlation coefficients, but not their variances. Consequently it is not possible to carry out inferences based on all overall correlation coefficients.

### 3.3.5 Variability test 1

The first test determines whether or not both methods  $A$  and  $B$  have the same between-subject variability, further to the second of Roy's criteria.

$$H_0 : d_A = d_B$$

$$H_A : d_A \neq d_B$$

This test is facilitated by constructing a model specifying a symmetric form for  $D$  (i.e. the alternative model) and comparing it with a model that has compound symmetric form for  $D$  (i.e. the null model). For this test  $\hat{\mathbf{A}}$  has a symmetric form for both models, and will be the same for both.

### 3.3.6 Variability test 2

This test determines whether or not both methods  $A$  and  $B$  have the same within-subject variability, thus enabling a decision on the third of Roy's criteria.

$$H_0 : \lambda_A = \lambda_B$$

$$H_A : \lambda_A \neq \lambda_B$$

This model is performed in the same manner as the first test, only reversing the roles of  $\hat{\mathbf{D}}$  and  $\hat{\mathbf{A}}$ . The null model is constructed a symmetric form for  $\hat{\mathbf{A}}$  while the alternative model uses a compound symmetry form. This time  $\hat{\mathbf{D}}$  has a symmetric form for both models, and will be the same for both.

As the within-subject variabilities are fundamental to the coefficient of repeatability, this variability test likelihood ratio test is equivalent to testing the equality of two coefficients of repeatability of two methods. In presenting the results of this test, Roy (2009a) includes the coefficients of repeatability for both methods.

### 3.3.7 Variability test 3

The last of the variability test examines whether or not methods  $A$  and  $B$  have the same overall variability. This enables the joint consideration of second and third criteria.

$$H_0 : \sigma_A = \sigma_B$$

$$H_A : \sigma_A = \sigma_B$$

The null model is constructed a symmetric form for both  $\hat{D}$  and  $\hat{\Lambda}$  while the alternative model uses a compound symmetry form for both.

### 3.3.8 Demonstration of Roy's testing

Roy provides three case studies, using data sets well known in method comparison studies, to demonstrate how the methodology should be used. The first two examples used are from the 'blood pressure' data set introduced by Bland and Altman (1999). The data set is a tabulation of simultaneous measurements of systolic blood pressure were made by each of two experienced observers (denoted 'J' and 'R') using a sphygmomanometer and by a semi-automatic blood pressure monitor (denoted 'S'). Three sets of readings were made in quick succession. Roy compares the 'J' and 'S' methods in the first of her examples.

The inter-method bias between the two method is found to be 15.62 , with a  $t$ -value of  $-7.64$ , with a  $p$ -value of less than 0.0001. Consequently there is a significant inter-method bias present between methods  $J$  and  $S$ , and the first of the Roy's three agreement criteria is unfulfilled.

Next, the first variability test is carried out, yielding maximum likelihood estimates of the between-subject variance covariance matrix, for both the null model, in compound symmetry (CS) form, and the alternative model in symmetric (symm) form. These matrices are determined to be as follows;

$$\hat{D}_{CS} = \begin{pmatrix} 946.50 & 784.32 \\ 784.32 & 946.50 \end{pmatrix}, \quad \hat{D}_{Symm} = \begin{pmatrix} 923.98 & 785.24 \\ 785.24 & 971.30 \end{pmatrix}.$$

A likelihood ratio test is performed to compare both candidate models. The log-likelihood of the null model is  $-2030.7$ , and for the alternative model  $-2030.8$ . The test statistic, presented with greater precision than the log-likelihoods, is  $0.1592$ . The  $p$ -value is  $0.6958$ . Consequently we fail to reject the null model, and by extension, conclude that the hypothesis that methods  $J$  and  $S$  have the same between-subject variability. Thus the second of the criteria is fulfilled.

The second variability test determines maximum likelihood estimates of the within-subject variance covariance matrix, for both the null model, in CS form, and the alternative model in symmetric form.

$$\hat{\Lambda}_{CS} = \begin{pmatrix} 60.27 & 16.06 \\ 16.06 & 60.27 \end{pmatrix}, \quad \hat{\Lambda}_{Symm} = \begin{pmatrix} 37.40 & 16.06 \\ 16.06 & 83.14 \end{pmatrix}.$$

Again, A likelihood ratio test is performed to compare both candidate models. The log-likelihood of the alternative model is  $-2045.0$ . As before, the null model has a log-likelihood of  $-2030.7$ . The test statistic is computed as  $28.617$ , again presented with greater precision. The  $p$ -value is less than  $0.0001$ . In this case we reject the null hypothesis of equal within-subject variability. Consequently the third of Roy's criteria is unfulfilled. The coefficient of repeatability for methods  $J$  and  $S$  are found to be  $16.95$  mmHg and  $25.28$  mmHg respectively.

The last of the three variability tests is carried out to compare the overall variabilities of both methods. With the null model the MLE of the within-subject variance covariance matrix is given below. The overall variabilities for the null and alternative models, respectively, are determined to be as follows;

$$\hat{\Sigma}_{CS} = \begin{pmatrix} 1007.92 & 801.65 \\ 801.65 & 1007.92 \end{pmatrix}, \quad \hat{\Sigma}_{Symm} = \begin{pmatrix} 961.38 & 801.40 \\ 801.40 & 1054.43 \end{pmatrix},$$

The log-likelihood of the alternative model is  $-2045.2$ , and again, the null model has a log-likelihood of  $-2030.7$ . The test statistic is  $28.884$ , and the  $p$ -value is less than  $0.0001$ . The null hypothesis, that both methods have equal overall variability,

is rejected. Further to the second variability test, it is known that this difference is specifically due to the difference of within-subject variabilities.

Lastly, Roy considers the overall correlation coefficient. The diagonal blocks  $\hat{\mathbf{r}}_{\Omega_{ii}}$  of the correlation matrix indicate an overall coefficient of 0.7959. This is less than the threshold of 0.82 that Roy recommends.

$$\hat{\mathbf{r}}_{\Omega_{ii}} = \begin{pmatrix} 1 & 0.7959 \\ 0.7959 & 1 \end{pmatrix}$$

The off-diagonal blocks of the overall correlation matrix  $\hat{\mathbf{r}}_{\Omega_{ii'}}$  present the correlation coefficients further to Hamlett et al. (2004).

$$\hat{\mathbf{r}}_{\Omega_{ii'}} = \begin{pmatrix} 0.9611 & 0.7799 \\ 0.7799 & 0.9212 \end{pmatrix}.$$

The overall conclusion of the procedure is that method  $J$  and  $S$  are not in agreement, specifically due to the within-subject variability, and the inter-method bias. The repeatability coefficients are substantially different, with the coefficient for method  $S$  being 49% larger than for method  $J$ . Additionally the overall correlation coefficient did not exceed the recommended threshold of 0.82.



### 3.4 Limits of agreement in LME models

Limits of agreement are used extensively for assessing agreement, because they are intuitive and easy to use. Necessarily their prevalence in literature has meant that they are now the best known measurement for agreement, and therefore any newer methodology would benefit by making reference to them.

Carstensen et al. (2008) uses LME models to determine the limits of agreement. Between-subject variation for method  $m$  is given by  $d_m^2$  and within-subject variation is given by  $\lambda_m^2$ . Carstensen et al. (2008) remarks that for two methods  $A$  and  $B$ , separate values of  $d_A^2$  and  $d_B^2$  cannot be estimated, only their average. Hence the assumption that  $d_x = d_y = d$  is necessary. The between-subject variability  $\mathbf{D}$  and within-subject variability  $\mathbf{\Lambda}$  can be presented in matrix form,

$$\mathbf{D} = \begin{pmatrix} d_A^2 & 0 \\ 0 & d_B^2 \end{pmatrix} = \begin{pmatrix} d^2 & 0 \\ 0 & d^2 \end{pmatrix}, \quad \mathbf{\Lambda} = \begin{pmatrix} \lambda_A^2 & 0 \\ 0 & \lambda_B^2 \end{pmatrix}.$$

The variance for method  $m$  is  $d_m^2 + \lambda_m^2$ . Limits of agreement are determined using the standard deviation of the case-wise differences between the sets of measurements by two methods  $A$  and  $B$ , given by

$$\text{var}(y_A - y_B) = 2d^2 + \lambda_A^2 + \lambda_B^2. \quad (3.9)$$

Importantly the covariance terms in both variability matrices are zero, and no covariance component is present.

Carstensen et al. (2008) presents a data set ‘fat’, which is a comparison of measurements of subcutaneous fat by two observers at the Steno Diabetes Center, Copenhagen. Measurements are in millimeters (mm). Each person is measured three times by each observer. The observations are considered to be ‘true’ replicates.

A linear mixed effects model is formulated, and implementation through several software packages is demonstrated. All of the necessary terms are presented in the computer output. The limits of agreement are therefore,

$$0.0449 \pm 1.96 \times \sqrt{2 \times 0.0596^2 + 0.0772^2 + 0.0724^2} = (-0.220, 0.309). \quad (3.10)$$

Roy (2009a) has demonstrated a methodology whereby  $d_A^2$  and  $d_B^2$  can be estimated separately. Also covariance terms are present in both  $\mathbf{D}$  and  $\mathbf{\Lambda}$ . Using Roy's methodology, the variance of the differences is

$$\text{var}(y_{iA} - y_{iB}) = d_A^2 + \lambda_B^2 + d_A^2 + \lambda_B^2 - 2(d_{AB} + \lambda_{AB}) \quad (3.11)$$

All of these terms are given or determinable in computer output. The limits of agreement can therefore be evaluated using

$$\bar{y}_A - \bar{y}_B \pm 1.96 \times \sqrt{\sigma_A^2 + \sigma_B^2 - 2(\sigma_{AB})}. \quad (3.12)$$

For Carstensen's 'fat' data, the limits of agreement computed using Roy's method are consistent with the estimates given by Carstensen et al. (2008);  $0.044884 \pm 1.96 \times 0.1373979 = (-0.224, 0.314)$ .

### 3.4.1 Linked replicates

Carstensen et al. (2008) proposes the addition of an random effects term to their model when the replicates are linked. This term is used to describe the 'item by replicate' interaction, which is independent of the methods. This interaction is a source of variability independent of the methods. Therefore failure to account for it will result in variability being wrongly attributed to the methods.

Carstensen et al. (2008) introduces a second data set; the oximetry study. This study done at the Royal Childrens Hospital in Melbourne to assess the agreement between co-oximetry and pulse oximetry in small babies.

In most cases, measurements were taken by both method at three different times. In some cases there are either one or two pairs of measurements, hence the data is unbalanced. Carstensen et al. (2008) describes many of the children as being very sick, and with very low oxygen saturations levels. Therefore it must be assumed that a biological change can occur in interim periods, and measurements are not true replicates.

Carstensen et al. (2008) demonstrate the necessity of accounting for linked replicated by comparing the limits of agreement from the 'oximetry' data set using a model

with the additional term, and one without. When the interaction is accounted for the limits of agreement are (-9.62,14.56). When the interaction is not accounted for, the limits of agreement are (-11.88,16.83). It is shown that the failure to include this additional term results in an over-estimation of the standard deviations of differences.

Limits of agreement are determined using Roy's methodology, without adding any additional terms, are found to be consistent with the 'interaction' model; (-9.562, 14.504). Roy's methodology assumes that replicates are linked. However, following Carstensen's example, an addition interaction term is added to the implementation of Roy's model to assess the effect, the limits of agreement estimates do not change. However there is a conspicuous difference in within-subject matrices of Roy's model and the modified model (denoted 1 and 2 respectively);

$$\hat{\mathbf{\Lambda}}_1 = \begin{pmatrix} 16.61 & 11.67 \\ 11.67 & 27.65 \end{pmatrix} \quad \hat{\mathbf{\Lambda}}_2 = \begin{pmatrix} 7.55 & 2.60 \\ 2.60 & 18.59 \end{pmatrix}. \quad (3.13)$$

(The variance of the additional random effect in model 2 is 3.01.)

Akaike (1974) introduces the Akaike information criterion (*AIC*), a model selection tool based on the likelihood function. Given a data set, candidate models are ranked according to their AIC values, with the model having the lowest AIC being considered the best fit. Two candidate models can said to be equally good if there is a difference of less than 2 in their AIC values.

The Akaike information criterion (AIC) for both models are  $AIC_1 = 2304.226$  and  $AIC_2 = 2306.226$ , indicating little difference in models. The AIC values for the Carstensen 'unlinked' and 'linked' models are 1994.66 and 1955.48 respectively, indicating an improvement by adding the interaction term.

The  $\hat{\mathbf{\Lambda}}$  matrices are informative as to the difference between Carstensen's unlinked and linked models. For the oximetry data, the covariance terms (given above as 11.67 and 2.6 respectively) are of similar magnitudes to the variance terms. Conversely for the 'fat' data the covariance term (-0.00032) is negligible. When the interaction term is added to the model, the covariance term remains negligible. (For the 'fat' data, the difference in AIC values is also approximately 2).

To conclude, Carstensen’s models provided a rigorous way to determine limits of agreement, but don’t provide for the computation of  $\hat{D}$  and  $\hat{\Lambda}$ . Therefore the test’s proposed by Roy (2009a) can not be implemented. Conversely, accurate limits of agreement as determined by Carstensen’s model may also be found using Roy’s method. Addition of the interaction term erodes the capability of Roy’s methodology to compare candidate models, and therefore shall not be adopted.

Finally, to complement the blood pressure (i.e. ‘J vs S’) method comparison from the previous section (i.e. ‘J vs S’), the limits of agreement are  $15.62 \pm 1.96 \times 20.33 = (-24.22, 55.46)$ .

### 3.5 Implementation in R

To implement an LME model in R, the `nlme` package is used. This package is loaded into the R environment using the `library` command, (i.e. `library(nlme)`). The `lme` command is used to fit LME models. The first two arguments to the `lme` function specify the fixed effect component of the model, and the data set to which the model is to be fitted. The first candidate model ('MCS1') fits an LME model on the data set 'dat'. The variable 'method' is assigned as the fixed effect, with the response variable 'BP' (i.e. blood pressure).

The third argument contains the random effects component of the formulation, describing the random effects, and their grouping structure. The `nlme` package provides a set of positive-definite matrices, the `pdMat` class, that can be used to specify a structure for the between-subject variance-covariance matrix for the random effects. For Roy's methodology, we will use the `pdSymm` and `pdCompSymm` to specify a symmetric structure and a compound symmetry structure respectively. A full discussion of these structures can be found in Pinheiro and Bates (1994, pg. 158).

Similarly a variety of structures for the within-subject variance-covariance matrix can be implemented using `nlme`. To implement a particular matrix structure, one must specify both a variance function and correlation structure accordingly. Variance functions are used to model the variance structure of the within-subject errors. `varIdent` is a variance function object used to allow different variances according to the levels of a classification factor in the data. A compound symmetry structure is implemented using the `corCompSymm` class, while the symmetric form is specified by `corSymm` class. Finally, the estimation method is specified as "ML" or "REML".

The first of Roy's candidate model can be implemented using the following code;

```
MCS1 = lme(BP ~ method-1, data = dat,  
random = list(subject=pdSymm(~ method-1)),  
weights=varIdent(form=~1|method),  
correlation = corSymm(form=~1 | subject/obs), method="ML")
```

For the blood pressure data used in Roy (2009a), all four candidate models are implemented by slight variations of this piece of code, specifying either `pdSymm` or `pdCompSymm` in the second line, and either `corSymm` or `corCompSymm` in the fourth line. For example, the second candidate model 'MCS2' is implemented with the same code as MCS1, except for the term `pdCompSymm` in the second line, rather than `pdSymm`.

```
MCS2 = lme(BP ~ method-1, data = dat,  
random = list(subject=pdCompSymm(~ method-1)),  
weights = varIdent(form=~1|method),  
correlation = corSymm(form=~1 | subject/obs), method="ML")
```

Using this R implementation for other data sets requires that the data set is structured appropriately (i.e. each case of observation records the index, response, method and replicate). Once formatted properly, implementation is simply a case of re-writing the first line of code, and computing the four candidate models accordingly.

To perform a likelihood ratio test for two candidate models, simply use the `anova` command with the names of the candidate models as arguments. The following piece of code implement the first of Roy's variability tests.

```
> anova(MCS1,MCS2)

      Model df      AIC      BIC  logLik  Test L.Ratio p-value
MCS1      1   8 4077.5 4111.3 -2030.7
MCS2      2   7 4075.6 4105.3 -2030.8 1 vs 2 0.15291 0.6958
>
```

The fixed effects estimates are the same for all four candidate models. The inter-method bias can be easily determined by inspecting a summary of any model. The summary presents estimates for all of the important parameters, but not the complete variance-covariance matrices (although some simple `R` functions can be written to overcome this). The variance estimates for the random effects for MCS2 is presented below.

```
Random effects:
Formula: ~method - 1 | subject
Structure: Compound Symmetry

      StdDev Corr
methodJ 30.765
methodS 30.765 0.829
Residual 6.115
```

Similarly, for computing the limits of agreement the standard deviation of the differences is not explicitly given. Again, A simple R function can be written to calculate the limits of agreement directly.



# Chapter 4

## Extending Current Methodologies

### 4.1 Extension of Roy's methodology

Roy's methodology is constructed to compare two methods in the presence of replicate measurements. Necessarily it is worth examining whether this methodology can be adapted for different circumstances.

An implementation of Roy's methodology, whereby three or more methods are used, is not feasible due to computational restrictions. Specifically there is a failure to reach convergence before the iteration limit is reached. This may be due to the presence of additional variables, causing the problem of non-identifiability. In the case of two variables, it is required to estimate two variance terms and four correlation terms, six in all. For the case of three variabilities, three variance terms must be estimated as well as nine correlation terms, twelve in all. In general for  $n$  methods has  $2 \times T_n$  variance terms, where  $T_n$  is the triangular number for  $n$ , i.e. the addition analogue of the factorial. Hence the computational complexity quite increases substantially for every increase in  $n$ .

Should an implementation be feasible, further difficulty arises when interpreting the results. The fundamental question is whether two methods have close agreement so as to be interchangeable. When three methods are present in the model, the null

hypothesis is that all three methods have the same variability relevant to the respective tests. The outcome of the analysis will either be that all three are interchangeable or that all three are not interchangeable.

The tests would not be informative as to whether any two of those three were interchangeable, or equivalently if one method in particular disagreed with the other two. Indeed it is easier to perform three pair-wise comparisons separately and then to combine the results.

Roy's methodology is not suitable for the case of single measurements because it follows from the decomposition for the covariance matrix of the response vector  $y_i$ , as presented in Hamlett et al. (2004). The decomposition depends on the estimation of correlation terms, which would be absent in the single measurement case. Indeed there can be no within-subject variability if there are no repeated terms for it to describe. There would only be the covariance matrix of the measurements by both methods, which doesn't require the use of LME models. To conclude, simpler existing methodologies, such as Deming regression, would be the correct approach where there only one measurements by each method.

## 4.2 Conclusion

Carstensen et al. (2008) and Roy (2009a) highlight the need for method comparison methodologies suitable for use in the presence of replicate measurements. Roy (2009a) presents a comprehensive methodology for assessing the agreement of two methods, for replicate measurements. This methodology has the added benefit of overcoming the problems of unbalanced data and unequal numbers of replicates. Implementation of the methodology, and interpretation of the results, is relatively easy for practitioners who have only basic statistical training. Furthermore, it can be shown that widely used existing methodologies, such as the limits of agreement, can be incorporated into Roy's methodology.

## 4.3 Outline of Thesis

In the first chapter the study of method comparison is introduced, while the second chapter provides a review of current methodologies. The intention of this thesis is to progress the study of method comparison studies, using a statistical method known as Linear mixed effects models. Chapter three shall describes linear mixed effects models, and how the use of the linear mixed effects models have so far extended to method comparison studies. Implementations of important existing work shall be presented, using the R programming language.

Model diagnostics are an integral component of a complete statistical analysis. In chapter three model diagnostics shall be described in depth, with particular emphasis on linear mixed effects models, further to chapter two.

For the fourth chapter, important linear mixed effects model diagnostic methods shall be extended to method comparison studies, and proposed methods shall be demonstrated on data sets that have become well known in literature on method comparison. The purpose is to both calibrate these methods and to demonstrate applications for them. The last chapter shall focus on robust measures of important parameters such as agreement.

# Bibliography

- ACR (2008). Acute Chest Pain ( suspected aortic dissection) - American College of Radiology Expert Group Report.
- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19(6), 716–723.
- Altman, D. (1978). Plotting probability ellipses. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 27(3), 347–349.
- Altman, D. and J. Bland (1983). Measurement in medicine: The analysis of method comparison studies. *Journal of the Royal Statistical Society. Series D (The Statistician)* 32(3), 307–317.
- Altman, D. G. (1991). *Practical Statistics for Medical Research*. Chapman and Hall.
- Barnhart, H., M. Haber, and L. Lin (2007). An overview of assessing agreement with continuous measurements. *Journal of Biopharmaceutical Statistics* 17, 529–569.
- Bartko, J. (1994). Measures of agreement: A single procedure. *Statistics in Medicine* 13, 737–745.
- Benjamini, Y. and Y. Hochberg (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. Roy. Statist. Soc. Ser. B* 57(1), 289–300.
- Bland, J. and D. Altman (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *The Lancet* i, 307–310.

- Bland, J. and D. Altman (1995). Comparing methods of measurement - why plotting difference against standard method is misleading. *The Lancet* 346, 1085–87.
- Bland, J. and D. Altman (1999). Measuring agreement in method comparison studies. *Statistical Methods in Medical Research* 8(2), 135–160.
- Bradley, E. L. and L. G. Blackwood (1989). Comparing paired data: A simultaneous test for means and variances. *The American Statistician* 43(4), 234–235.
- Carroll, R. and D. Ruppert (1996). The use and misuse of orthogonal regression in linear errors-in-variables models. *The American Statistician* 50(1), 1–6.
- Carstensen, B., J. Simpson, and L. C. Gurrin (2008). Statistical models for assessing agreement in method comparison studies with replicate measurements. *The International Journal of Biostatistics* 4(1).
- Cohen, J., P. Cohen, S. West, and L. Aiken. *Applied multiple regression / correlation analysis for the behavioral sciences* (Third ed.). Laurence Erlbaum Associates.
- Cornbleet, P. J. and D. Cochrane (1979). Regression methods for assessing agreement between two methods of clinical measurement. *Journal of Clinical Chemistry* 24(2), 342–345.
- Dempster, A., N. Laird, and D. Rubin (1977). Maximum likelihood from incomplete data via the em algorithm. *ournal of the Royal Statistical Society. Series B* 39(1), 1–38.
- Dewitte, K., C. Fierens, D. Stckl, and L. M. Thienpont (2002). Application of the Bland Altman plot for interpretation of method-comparison studies: A critical investigation of its practice. *Clinical Chemistry* 48, 799–801.
- Dodge, Y. (2003). *The Oxford Dictionary of Statistical Terms*. Oxford University Press.

- Dunn, G. (2002). *Statistical Evaluation of Measurement Error* (Second ed.). Stanford: American Mathematical Society and Digital Press.
- Eisenhart, C. (1947). The assumptions underlying the analysis of variance. *Biometrics* 3(1), 1–21.
- Eksborg, S. (1981). Evaluation of method-comparison data [letter]. *Clinical Chemistry* 27, 1311–1312.
- Fisher, R. (1918). The correlation between relatives on the supposition of mendelian inheritance. *Transactions of the Royal Society of Edinburgh* 2, 399–433.
- Green, P. J. and B. W. Silverman (1994). *Nonparametric Regression and Generalized Linear Models: a Roughness Penalty Approach*. Chapman & Hall Ltd.
- Grubbs, F. (1948). On estimating precision of measuring instruments and product variability. *Journal of the American Statistical Association* 43, 243–264.
- Grubbs, F. (1973). Errors of measurement, precision, accuracy and the statistical comparison of measuring instruments. *Technometrics* 15(1), 53–66.
- Hamlett, A., L. Ryan, and R. Wolfinger (2004). On the use of PROC MIXED to estimate correlation in the presence of repeated measures. *Proceedings of the Statistics and Data Analysis Section, SAS Users Group International 198-229*, 1–7.
- Hartley, H. and J. Rao (1967). Maximum-likelihood estimation for the mixed analysis of variance model. *Biometrika* 54(1/2), 93–108.
- Henderson, C. (1953). Estimation of variance and covariance components. *Biometrics* 9(2), 226–252.
- Henderson, C., O. Kempthorne, S. Searle, and C. von Krosigk (1959). The estimation of environmental and genetic trends from records subject to culling. *Biometrics* 15, 192–218.

- Henderson, C., O. Kempthorne, S. Searle, and C. von Krosigk (1963). The estimation of environmental and genetic trends from records subject to culling. *Biometrics* 15, 192–218.
- Henderson, C., O. Kempthorne, S. Searle, and C. von Krosigk (1973). The estimation of environmental and genetic trends from records subject to culling. *Biometrics* 15, 192–218.
- Henderson, C., O. Kempthorne, S. Searle, and C. von Krosigk (1984). The estimation of environmental and genetic trends from records subject to culling. *Biometrics* 15, 192–218.
- Henderson, C. R. (1950). Estimation of genetic parameters (abstract). *Annals of Mathematical Statistics* 21, 309–310.
- Henderson, C. R. (1973). Sire evaluation and genetic trends. In *Proceedings of the Animal Breeding and genetics Symposium in Honor of Dr Jay L. Lush*, pp. 10–41. Champaign, Illinois: American Society of Animal Science and American Dairy Science Association.
- Hollis, S. (1996). Analysis of method comparison studies. *Ann Clin Biochem* 33, 1–4.
- IUPAC (2009). IUPAC Compendium of Chemical Terminology - the Gold Book. <http://goldbook.iupac.org/R05293.html>.
- Kinsella, A. (1986). Estimating method precision. *The Statistician* 35, 421–427.
- Krummenauer, F., I. Genevriere, and U. Nixdorff (2000). The biometrical comparison of cardiac imaging methods. *Computer Methods and Programs in Biomedicine* 62, 21–34.
- Kummel, C. (1879). Reduction of observation equations which contain more than one observed quantity. *The Analyst* 6, 97–105.

- Laird, N. M. and J. H. Ware (1982). Random-effects models for longitudinal data. *Biometrics* 38(4), 963–974.
- Lam, M., K. Webb, and D. O'Donnell (1999). Correlation between two variables in repeated measurements. *American Statistical Association, Proceedings of the Biometric Session*, 213–218.
- Lee, Y. and J. A. Nelder (1996). Hierarchical generalized linear models (Disc: P656-678). *Journal of the Royal Statistical Society, Series B: Methodological* 58, 619–656.
- Lee, Y., J. A. Nelder, and Y. Pawitan (2006). *Generalized Linear Models with Random Effects: Unified Analysis via H-likelihood*. Chapman & Hall Ltd.
- Lewis, P., P. Jones, J. Polak, and H. Tillitson (1991). The problem of conversion in method comparison studies. *Applied Statistics* 40, 105–112.
- Lin, L. (2000). Total deviation index for measuring individual agreement with applications in laboratory performance and bioequivalence. *Statistics in medicine* 97, 255–270.
- Lin, L., A. Hedayat, B. Sinha, and M. Yang (2002). Statistical methods in assessing agreement: Models issues and tools. *Journal of American Statistical Association* 97, 257–270.
- Linnet, K. (1998). Performance of deming regression analysis in case of misspecified analytical error ratio in method comparison studies. *Clinical Chemistry* 44, 1024–1031.
- Linnet, K. (1999). Necessary sample size for method comparison studies based on regression analysis. *Clinical Chemistry* 45(6), 882–894.
- Ludbrook, J. (1997). Comparing methods of measurement. *Clinical and Experimental Pharmacology and Physiology* 24, 193–203.



- Luiz, R., A. Costa, P. Kale, and G. Werneck (2003). Assessment of agreement of a quantitative variable: a new graphical approach. *Journal of Clinical Epidemiology* 56, 963–967.
- Mantha, S., M. F. Roizen, L. A. Fleisher, R. Thisted, and J. Foss (2000). Comparing methods of clinical measurement: Reporting standards for bland and altman analysis. *Anaesthesia and Analgesia* 90, 593–602.
- Morgan, W. A. (1939). A test for the significance of the difference between two variances in a sample from a normal bivariate population. *Biometrika* 31, 13–19.
- Myles, P. (2007). Using the Bland Altman method to measure agreement with repeated measures. *British Journal of Anaesthesia* 99(3), 309–311.
- NIST (2009). Cesium fountain atomic clock : The primary time and frequency standard for the United States. <http://tf.nist.gov/timefreq/cesium/fountain.htm>.
- O’Brien, E., J. Petrie, W. Littler, M. de Swiet, P. L. Padfield, D. Altman, M. Bland, A. Coats, and N. Atkins (1990). The British Hypertension Society protocol for the evaluation of blood pressure measuring devices. *Journal of Hypertension* 8, 607–619.
- Patterson, H. and R. Thompson (1971). Recovery of inter-block information when block sizes are unequal. *Biometrika* 58(3), 545–554.
- Pawitan, Y. (2001). *In All Likelihood: Statistical Modelling and Inference Using Likelihood*. Oxford University Press.
- Phelps, C. and A. Hutson (1995). Estimating diagnostic test accuracy using a fuzzy gold standard. *Medical decision making* 15, 144–57.
- Pinheiro, J. and D. Bates (1994). *Mixed Effects Models in S and S plus* (2nd ed.). Reading, Massachusetts: Springer.
- Pitman, E. J. G. (1939). A note on normal correlation. *Biometrika* 31, 9–12.

- Pizzi, N. (1999). Fuzzy pre-processing of gold standards as applied to biomedical spectra classification. *Artificial Intelligence in Medicine* 16, 171–182.
- Robinson, G. K. (1991). That BLUP is a good thing: The estimation of random effects (Disc: P32-51). *Statistical Science* 6, 15–32.
- Roy, A. (2006). Estimating correlation coefficient between two variables with repeated observations using mixed effects models. *Biometric Journal* 2, 286–301.
- Roy, A. (2009a). An application of linear mixed effects model to assess the agreement between two methods with replicated observations. *Journal of Biopharmaceutical Statistics* 19, 150–173.
- Roy, A. (2009b). An application of the linear mixed effects model to ass the agreement between two methods with replicated observations. *Journal of Biopharmaceutical Statistics* 19, 150–173.
- Ryan, T. P. and W. H. Woodall (2005). The most-cited statistical papers. *Journal of Applied Statistics* 32(5), 461 – 474.
- Singer, J. D. (1998). Using sas proc mixed to fit multilevel models, hierarchical models, and individual growth models. *Journal of Educational and Behavioral Statistics* 24(4), 323–355.
- Thompson, W. (1963). Precision of simultaneous measurement procedures. *Journal of American Statistical Association* 58, 474–479.
- Tippett, L. (1931). *The Methods of Statistics* (1st ed.). London: Williams and Norgate.
- West, B., K. Welch, and A. Galecki (2007). *Linear Mixed Models: a Practical Guide Using Statistical Software*. Chapman and Hall CRC.
- Zhang, Y., S. Nitter-Hauge, H. Ihlen, K. Rootwelt, and E. Myhre (1986). Measurement of aortic regurgitation by doppler echocardiography. *British Heart Journal* 55, 32–38.