

One of the assumptions underlying ordinal logistic (and ordinal probit) regression is that the relationship between each pair of outcome groups is the same. In other words, ordinal logistic regression assumes that the coefficients that describe the relationship between, say, the lowest versus all higher categories of the response variable are the same as those that describe the relationship between the next lowest category and all higher categories, etc.

## Ordered logistic regression R

This is called the proportional odds assumption or the parallel regression assumption. Because the relationship between all pairs of groups is the same, there is only one set of coefficients. If this was not the case, we would need different sets of coefficients in the model to describe the relationship between each pair of outcome groups. Thus, in order to assess the appropriateness of our model, we need to evaluate whether the proportional odds assumption is tenable.

## Ordered logistic regression R

Statistical tests to do this are available in some software packages. However, these tests have been criticized for having a tendency to reject the null hypothesis (that the sets of coefficients are the same), and hence, indicate that there the parallel slopes assumption does not hold, in cases where the assumption does hold (see Harrell 2001 p. 335). We were unable to locate a facility in R to perform any of the tests commonly used to test the parallel slopes assumption.

## Ordered logistic regression R

However, Harrell does recommend a graphical method for assessing the parallel slopes assumption. The values displayed in this graph are essentially (linear) predictions from a logit model, used to model the probability that  $y$  is greater than or equal to a given value (for each level of  $y$ ), using one predictor ( $x$ ) variable at a time. In order create this graph, you will need the Hmisc library.

## Ordered logistic regression R

The code below contains two commands (the first command falls on multiple lines) and is used to create this graph to test the proportional odds assumption. Basically, we will graph predicted logits from individual logistic regressions with a single predictor where the outcome groups are defined by either apply  $i=2$  and apply  $i=3$ .

## Ordered logistic regression R

If the difference between predicted logits for varying levels of a predictor, say  $x$ , are the same whether the outcome is defined by  $y=2$  or  $y=3$ , then we can be confident that the proportional odds assumption holds. In other words, if the difference between logits for  $x=0$  and  $x=1$  is the same when the outcome is  $y=2$  as the difference when the outcome is  $y=3$ , then the proportional odds assumption likely holds.

## Ordered logistic regression R

The first command creates the function that estimates the values that will be graphed. The first line of this command tells R that `sf` is a function, and that this function takes one argument, which we label `y`. The `sf` function will calculate the log odds of being greater than or equal to each value of the target variable. For our purposes, we would like the log odds of apply being greater than or equal to 2, and then greater than or equal to 3. Depending on the number of categories in your dependent variable, and the coding of your variables, you may have to edit this function. Below the function is configured for a `y` variable with three levels, 1, 2, 3.