

# Introduction – systematic vs. random errors

Krzysztof Podgórski  
Department of Mathematics and Statistics  
University of Limerick

September 10, 2009

# Quantitative nature of analytical chemistry

# Quantitative nature of analytical chemistry

- Modern analytical chemistry is overwhelmingly a quantitative science.

# Quantitative nature of analytical chemistry

- Modern analytical chemistry is overwhelmingly a quantitative science.
- A quantitative answer is much more valuable than a qualitative one.

# Quantitative nature of analytical chemistry

- Modern analytical chemistry is overwhelmingly a quantitative science.
- A quantitative answer is much more valuable than a qualitative one. *It may be useful for an analyst to claim to have detected some boron in a distilled water sample, but it is much more useful to be able to say how much boron is present.*

# Quantitative nature of analytical chemistry

- Modern analytical chemistry is overwhelmingly a quantitative science.
- A quantitative answer is much more valuable than a qualitative one. *It may be useful for an analyst to claim to have detected some boron in a distilled water sample, but it is much more useful to be able to say how much boron is present.*
- Often it is only a quantitative result that has any value at all.

# Quantitative nature of analytical chemistry

- Modern analytical chemistry is overwhelmingly a quantitative science.
- A quantitative answer is much more valuable than a qualitative one. *It may be useful for an analyst to claim to have detected some boron in a distilled water sample, but it is much more useful to be able to say how much boron is present.*
- Often it is only a quantitative result that has any value at all. *for example, almost all samples of (human) blood serum contain albumin; the only question is, how much?*

# Quantitative nature of analytical chemistry

- Modern analytical chemistry is overwhelmingly a quantitative science.
- A quantitative answer is much more valuable than a qualitative one. *It may be useful for an analyst to claim to have detected some boron in a distilled water sample, but it is much more useful to be able to say how much boron is present.*
- Often it is only a quantitative result that has any value at all. *for example, almost all samples of (human) blood serum contain albumin; the only question is, how much?*
- Even where a qualitative answer is required, quantitative methods are used to obtain it.



# Quantitative nature of analytical chemistry

- Modern analytical chemistry is overwhelmingly a quantitative science.
- A quantitative answer is much more valuable than a qualitative one. *It may be useful for an analyst to claim to have detected some boron in a distilled water sample, but it is much more useful to be able to say how much boron is present.*
- Often it is only a quantitative result that has any value at all. *for example, almost all samples of (human) blood serum contain albumin; the only question is, how much?*
- Even where a qualitative answer is required, quantitative methods are used to obtain it. *Quantitative approaches might be used to compare two soil samples. For example, they might be subjected to a particle size analysis, in which the proportions of the soil particles falling within a number, say 10, of particle-size ranges are determined. Each sample would then be characterized by these 10 pieces of data, which could then be used to provide a quantitative assessment of their similarity.*

# Errors in quantitative analysis

# Errors in quantitative analysis

Since quantitative studies play a dominant role in any analytical laboratory, it must be accepted that the errors that occur in such studies are of supreme importance.

# Errors in quantitative analysis

Since quantitative studies play a dominant role in any analytical laboratory, it must be accepted that the errors that occur in such studies are of supreme importance. **No quantitative results are of any value unless they are accompanied by some estimate of the errors inherent in them!**

## Example 1 – detecting a new analytical reagent

## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.

## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.
- The compound is studied using a spectrometric method and gives a value of 104.

## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.
- The compound is studied using a spectrometric method and gives a value of 104.
- The chemist finds that no compound previously discovered has yielded a value of more than 100.



## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.
- The compound is studied using a spectrometric method and gives a value of 104.
- The chemist finds that no compound previously discovered has yielded a value of more than 100.
- Has the chemist really discovered a new compound?

## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.
- The compound is studied using a spectrometric method and gives a value of 104.
- The chemist finds that no compound previously discovered has yielded a value of more than 100.
- Has the chemist really discovered a new compound?
- The answer lies in the degree of reliance to experimental value of 104.

## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.
- The compound is studied using a spectrometric method and gives a value of 104.
- The chemist finds that no compound previously discovered has yielded a value of more than 100.
- Has the chemist really discovered a new compound?
- The answer lies in the degree of reliance to experimental value of 104.
- If the result is correct to within 2 (arbitrary) units, i.e. the true value probably lies in the range  $102 \pm 2$ , then a new material has probably been discovered.

## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.
- The compound is studied using a spectrometric method and gives a value of 104.
- The chemist finds that no compound previously discovered has yielded a value of more than 100.
- Has the chemist really discovered a new compound?
- The answer lies in the degree of reliance to experimental value of 104.
- If the result is correct to within 2 (arbitrary) units, i.e. the true value probably lies in the range  $102 \pm 2$ , then a new material has probably been discovered.
- If, however, investigations show that the error may amount to 10 units i.e.  $104 \pm 10$ , then it is quite likely that the true value is actually less than 100, in which case a new discovery is far from certain.

## Example 1 – detecting a new analytical reagent

- A chemist synthesizes an analytical reagent that is believed to be entirely new.
- The compound is studied using a spectrometric method and gives a value of 104.
- The chemist finds that no compound previously discovered has yielded a value of more than 100.
- Has the chemist really discovered a new compound?
- The answer lies in the degree of reliance to experimental value of 104.
- If the result is correct to within 2 (arbitrary) units, i.e. the true value probably lies in the range  $102 \pm 2$ , then a new material has probably been discovered.
- If, however, investigations show that the error may amount to 10 units i.e.  $104 \pm 10$ , then it is quite likely that the true value is actually less than 100, in which case a new discovery is far from certain.
- A knowledge of the experimental errors is crucial.

## Example 2 – replicates in a titrimetric experiment

## Example 2 – replicates in a titrimetric experiment

- Analysts commonly perform several replicate determinations in the course of a single experiment.

## Example 2 – replicates in a titrimetric experiment

- Analysts commonly perform several replicate determinations in the course of a single experiment.
- An analyst performs a titrimetric experiment four times and obtains values of 24.69, 24.73, 24.77 and 25.39 ml.



## Example 2 – replicates in a titrimetric experiment

- Analysts commonly perform several replicate determinations in the course of a single experiment.
- An analyst performs a titrimetric experiment four times and obtains values of 24.69, 24.73, 24.77 and 25.39 ml.
- All four values are different, because of the variations inherent in the measurements

## Example 2 – replicates in a titrimetric experiment

- Analysts commonly perform several replicate determinations in the course of a single experiment.
- An analyst performs a titrimetric experiment four times and obtains values of 24.69, 24.73, 24.77 and 25.39 ml.
- All four values are different, because of the variations inherent in the measurements
- The fourth value (25.39 ml) is substantially different from the other three.

## Example 2 – replicates in a titrimetric experiment

- Analysts commonly perform several replicate determinations in the course of a single experiment.
- An analyst performs a titrimetric experiment four times and obtains values of 24.69, 24.73, 24.77 and 25.39 ml.
- All four values are different, because of the variations inherent in the measurements
- The fourth value (25.39 ml) is substantially different from the other three.
- Can it be safely rejected, so that (for example) the mean titre is reported as 24.73 ml, the average of the other three readings?

# Systematic and random errors

# Systematic and random errors

Experimental scientists make a fundamental distinction between **random**, and **systematic errors**. To distinguish between random and systematic errors let us consider a real experiment.

# Systematic and random errors

Experimental scientists make a fundamental distinction between **random**, and **systematic errors**. To distinguish between random and systematic errors let us consider a real experiment.

- Four students (A-D) each perform an analysis in which exactly 10.00 ml of exactly 0.1 M sodium hydroxide is titrated with exactly 0.1 M hydrochloric acid.

# Systematic and random errors

Experimental scientists make a fundamental distinction between **random**, and **systematic errors**. To distinguish between random and systematic errors let us consider a real experiment.

- Four students (A-D) each perform an analysis in which exactly 10.00 ml of exactly 0.1 M sodium hydroxide is titrated with exactly 0.1 M hydrochloric acid.
- Each student performs five replicate titrations, with the results shown in Table 1.1.

**Table 1.1** Random and systematic errors


Student	Results (ml)					Comment
A	10.08	10.11	10.09	10.10	10.12	Precise, biased
B	9.88	10.14	10.02	9.80	10.21	Imprecise, unbiased
C	10.19	9.79	9.69	10.05	9.78	Imprecise, biased
D	10.04	9.98	10.02	9.97	10.04	Precise, unbiased

# Graphical illustration



# Graphical illustration

The results of experiment represented by dotplots



**The true value is 10.00**

# Systematic error and bias

# Systematic error and bias

- **Systematic error** is a deviation of all measurements in one direction from the true value.

# Systematic error and bias

- **Systematic error** is a deviation of all measurements in one direction from the true value.
- It is well represented by the difference between the average value of the determined values and the true value of the measured quantity.

# Systematic error and bias

- **Systematic error** is a deviation of all measurements in one direction from the true value.
- It is well represented by the difference between the average value of the determined values and the true value of the measured quantity.
- This difference is called the **bias** of measurements.

# Random error and precision

# Random error and precision

- **Random error** is a deviation of a measurement from the average of measured values.

# Random error and precision

- **Random error** is a deviation of a measurement from the average of measured values.
- It is well represented by the standard deviation of measurements.




# Random error and precision

- **Random error** is a deviation of a measurement from the average of measured values.
- It is well represented by the standard deviation of measurements.
- This value is often called **precision** of measurements.

# Combined error vs. accuracy

# Combined error vs. accuracy

- **Accuracy** is in inverse relation to the total deviation of a single measurement from the true value.



# VALIDATION

# VALIDATION

- A validation survey was conducted about these rules.

# VALIDATION

- A validation survey was conducted about these rules.
- Almost all the people surveyed about these rules were valid.

# VALIDATION

- A validation survey was conducted about these rules.
- Almost all the people surveyed about these rules were valid.
- A few were invalid, but they expected to recover soon.

# VALIDATION

- A validation survey was conducted about these rules.
- Almost all the people surveyed about these rules were valid.
- A few were invalid, but they expected to recover soon.
- Based on the survey, a statistical confidence level was determined.



# VALIDATION

- A validation survey was conducted about these rules.
- Almost all the people surveyed about these rules were valid.
- A few were invalid, but they expected to recover soon.
- Based on the survey, a statistical confidence level was determined.
- Ninety-five percent of the people surveyed have at least 67 percent confidence in statistics.

# VALIDATION

- A validation survey was conducted about these rules.
- Almost all the people surveyed about these rules were valid.
- A few were invalid, but they expected to recover soon.
- Based on the survey, a statistical confidence level was determined.
- Ninety-five percent of the people surveyed have at least 67 percent confidence in statistics.
- If you did not notice - **it is just a joke!**