

PRELIMINARY EXAM

PROGRAMME(S) : University of London Degree and Diploma Programmes
(Lead College: London School of Economics & Political Science)

SUBJECT : **04A Statistics 1 (half unit)**

DATE :

DURATION : 2 hours

INSTRUCTIONS :-

**DO NOT TURN OVER THIS QUESTION PAPER UNTIL YOU
ARE TOLD TO DO SO.**

Candidates should answer **THREE** of the following **FOUR** questions:
QUESTION 1 of Section A (50 marks) and **TWO** questions from Section B (25 marks each).

Graph paper is provided. When used, it should be fastened securely **inside** the answer book.

Statistical tables are provided.

A handheld calculator may be used when answering questions on this paper, but it must not be pre-programmed or able to display graphics, text, or algebraic equations. The make and type of the machine must be clearly stated on the front cover of the answer book.

Candidates are strongly advised to divide their time accordingly.

SECTION A

Answer all **eight** parts of question 1 (50 marks in total).

1. (a) Define each of the following briefly:

- i. quota sampling.
- ii. stratified sampling.
- iii. an alternative hypothesis. **(6 marks)**

(b) Which of the following is the odd one out?
Standard deviation, range, median, variance.
Explain briefly. **(2 marks)**

(c) The following tables show the scores of two groups of students in a test question.

i.	Mark	0	1	2	3	4
	Frequency	10	5	8	x	3

Determine the value of x if the median of these marks is 1.5

ii.	Mark	0	1	2	3	4
	Frequency	10	5	8	y	3


Determine the value of y if the mean of these marks is 1.5
(4 marks)

- (d) i. Two fair dice are thrown and you are told that the sum of their upturned faces is equal to 7. What is the probability that neither face is equal to 6?
- ii. Three balls are thrown at random into five bowls so that each ball has the same chance of going into any bowl independently of wherever the other 2 balls fall. Determine the probability distribution of the number of empty bowls. **(8 marks)**

(question continues on next page)

PLEASE TURN OVER

- (e) A scatter plot for two variables (x and y) is shown below.
- Do you consider these variables to be independent? Explain.
 - State an approximate value for their correlation coefficient.
 - If the data set were split into two at the centre of the x-range,
 - State an approximate value for the correlation of the smaller set.
 - State an approximate value for the correlation of the larger set.



- (f) In the past, 18% of shoppers have bought a particular brand of breakfast cereal. After an advertising campaign for this brand, a random sample of 220 shoppers is taken and 55 of these are found to have bought this brand of cereal. Carry out the appropriate significance test at the 5% level to test whether the advertising campaign has been successful. Comment briefly.
- (9 marks)

(question continues on next page)

(g) $x_1 = 1, x_2 = 3, x_3 = 4, x_4 = 2$

i. $\sum_{i=1}^{i=2} x_i^3$

ii. $\sum_{i=2}^{i=4} (x_i - 4)^2$

iii. $\sum_{i=3}^{i=4} 2x_i$

(6 marks)

(h) State whether the following are possible or not. Give a brief explanation. (*Note that no marks will be awarded for a simple possible/not possible reply*)

- i. Only John or Peter may get the end of year prize. The probability John gets it is .4 . That for Peter is .7 .
- ii. The correlation coefficient for the relationship between the number of years a woman spent at school and the number of children she has is minus 3 . 5 .
- iii. The variance of the results is minus 64.
- iv. The larger a sample, the smaller the variation in its sample mean.

(8 marks)

PLEASE TURN OVER

SECTION B

Answer **two** questions from this section (25 marks each).

2. (a) The following data show the lengths (inches) of fish caught in one day in a river.

15.2	15.6	14.6	10.5	13.7	10.6	13.9	14.7	13.9	10.6
10.3	10.6	12.1	12.8	14.9	15.2	13.8	14.5	15.0	11.7
10.9	18.4	14.1	14.9	10.2	13.9	11.6	13.9	12.5	11.1
15.1	14.9	15.0	12.6	10.9	13.7	12.9	12.1	14.0	13.5
15.3	13.8	14.9	10.6	15.1	11.6	11.0	14.8	13.8	10.3

- i. Construct a stem and leaf plot for these data. Label it carefully.
 - ii. Use this to determine the median and upper and lower data quartiles.
 - iii. Briefly describe the figures shown. **(12 marks)**
- (b) A machine filling cans supplies, to each can, a volume X of fruit and a volume Y of juice. It is known that X has a normal distribution with mean 260 and a standard deviation of 17 whereas Y has a normal distribution with mean 150 and a standard deviation of 10. These variables may be regarded as independent.
- i. Calculate the probability that the volume of juice supplied is greater than 290 units.
 - ii. Find the probability that the volume of fruit supplied is more than twice the amount of juice.
 - iii. If the capacity of a particular can is 400, what is the probability of the can being under-filled? **(13 marks)**

3. (a) The following data set shows the times (y) in seconds taken by international athletes to run 100 metres together with the corresponding wind speeds (x) at the time of running. A positive wind speed indicates the wind is in the direction of running and therefore considered to be helpful whereas a negative wind speed indicates the wind is against the runner.

Wind Speed (x)	-2.94	-1.63	-0.74	-0.38	-0.17
Time for 100 metres	10.51	10.44	10.39	10.22	10.44
Wind Speed (x)	0.21	0.60	1.08	2.25	2.81
Time for 100 metres	10.18	10.30	10.15	10.20	10.21

N.B. summary statistics for these data are,

Sum of x data = 1.09 Sum of y data = 103.04

Sum of squares of x data = 26.5505 Sum of squares of y data = 1061.8770

- i. Draw and label the scatter diagram of these data as described here. Label it carefully.
 - ii. Comment on your diagram.
 - iii. Calculate the correlation coefficient.
 - iv. Comment on your figures.
 - v. Would you think that it would be useful to calculate a regression line to use for predictive purposes? **(15 marks)**
- (b) You run a market research company and have been asked to give a tender showing how you could carry out a survey of a large international company's employees to find out about their holiday preferences. You have been told that you may use email (the company will make email addresses available for all by country)
- Outline your design making it clear whether it is random or not and giving appropriate design factors. **(10 marks)**

PLEASE TURN OVER

4. (a) The following table shows the classification by attitude and social class of a random group of 698 residents to a council proposal on traffic calming.

		Social Class		
		Lower	Middle	Upper
Attitude to proposal	Approved	34	45	61
	Indifferent	55	30	203
	Disapproved	28	22	220

- i. Carry out the appropriate formal test at two levels of significance.
 - ii. Comment on your results - are there any particular figures you find interesting here? **(12 marks)**
- (b) A simple random sample of 350 wives were asked whether they ever did 'Do it yourself' jobs in the home. Of these, 90 replied that they did. Calculate the 95% confidence interval for the proportion of housewives who do 'Do it yourself' jobs. **(4 marks)**
- (c) A study was carried out to see if an extended contact with their newborn babies in the first minute of life caused the mother to have a greater attachment to the baby. An attachment index was calculated for each mother based on the mother's attitude to the baby in the week after birth. For this study, two groups (control and treatment) of mothers were used. The *control group* of 62 mothers had the standard amount of contact with their babies after birth. The *treatment group* of 73 mothers were given extended contact with their babies in the first minute after birth. Summary statistics are given below. Note that the higher the index figure, the closer the mother's attachment.

	Control Group	Treatment Group
Number in group	62	73
Sample mean	6.29	8.53
Sample Standard Deviation	2.07	1.95

- i. Carry out a test, giving the null and alternative hypotheses, to see whether the extended contact has an effect.
- ii. Have you any reservations about these results in terms of the long term effect on family relationships? **(9 marks)**

END OF PAPER