$PETCT_Body_LM.MlAdult_PET5$

3. december 2012

Indhold

1	Top	Topogram														1							
	1.1	Routine																 					1
	1.2	Scan .						•															1
2	CT Body															1							
	2.1	Routine																 					1
	2.2	Scan .																					2
	2.3	Recons																 					2
		2.3.1	Recon 1															 					2
		2.3.2	Recon 2															 					3
		2.3.3	Recon 3																				4
3	Pau	ıse																					4
4	PE	ET Body LM															4						
	4.1	Routine																					4
	4.2	Scan .																 					5
	4.3	Recons																 					5
		4.3.1	Recon 1															 					5

1 Topogram

1.1 Routine

• mA: 35

• kV: 120

• Topogram length: 512 mm

• Tube position: Top

1.2 Scan

• mA: 35

• kV: 120

• Delay: 4s

• Topogram length: 512 mm

• Direction: Craniocaudal

• Tube position: Top

• API: None

• Kernel: 80s shar

• Window: Topogram Body

2 CT Body

2.1 Routine

• Eff. mAs: 170

• kV: 120

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

• CTDlvol: 11.4779mGy

• Scan time: 5.630 s

• Delay: 4.000 s

• Slice: 5.00 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

2.2 Scan

• Quality ref. mAs: 170

• Eff. mAs: 170

• kV: 120

• Scan time: 5.630 s

• Rotation time: 0.500 s

• Delay: 4.000 s

• Slice: 5.00 mm

• Pitch: 0.8000000

• Direction: Craniocaudal

2.3 Recons

2.3.1 Recon 1

• Series description: tmp CT Body RTD

• Slice: 5.00

• Kernel: B19f PET very smooth

• Window: Abdomen

• Extended FoV: On

• FoV: 700

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 11.500

• No. of images: 20

2.3.2 Recon 2

• Series description: AC CT

• Slice: 5.00

• Kernel: B19f PET very smooth

• Window: Abdomen

• Extended FoV: On

• FoV: 700

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 3.000

• No. of images: 74

2.3.3 Recon 3

• Series description: CT Body 3.0 B30f

• Slice: 3.00

• Kernel: B30f medium smooth

• Window: Abdomen

• Extended FoV: Off

• FoV: 500

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 2.000

• No. of images: 111

3 Pause

4 PET Body LM

4.1 Routine

• Isotope: F-18

• Pharm.: FDG

• Inj. Dose: 10 Millicuries

• Inj. date (date/month - year): 14/09 - 2006

• Inj. time: 11:24:00

- Scan mode: List mode
- Scan range: Match CT FOV
- No. of beds: 1
- Scan duration/bed: 60 Minutes

4.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: List mode
- Sinogram mode: Trues
- Input trigger signal: None
- LLD (keV): 435
- ULD (keV): 650

4.3 Recons

4.3.1 Recon 1

- Series description: PET WB
- Recon range (bed): Automatic/All/dunno
- Output image type: NoRecon
- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 168
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 5

• Offset X: 0 mm

• Offset Y: 0 mm

• Attenuation correction: On (1)

• Scatter correction: On

• Match CT slice location: On

• Save intermediate data: Off