# $PETCT\_WholeBody\_HD.MlAdult\_PET6$

30. maj 2013

## Indhold

1	Top	ogram																						
	1.1	Routine																						
	1.2	Scan			•		•			•	•						•			•				
<b>2</b>	CT	WB																						
	2.1	Routine																						
	2.2	Scan																						
	2.3	Recons .																						
		2.3.1 R	econ 1																					
		2.3.2 R	econ 2																					
		2.3.3 R	econ 3				•						•				•						•	
3	Pau	ıse																						
4	PET WB																							
	4.1	Routine																						
	4.2	Scan																						
	4.3	Recons .																						
		4.3.1 R	econ 1																					
		4.3.2 R	econ 2																					

## 1 Topogram

#### 1.1 Routine

mA: 35kV: 120

• Topogram length: 1024 mm

• Tube position: Top

#### 1.2 Scan

mA: 35kV: 120Delay: 4s

Topogram length: 1024 mmDirection: Craniocaudal

• Tube position: Top

API: NoneKernel: 80s shar

• Window: Topogram Body

## 2 CT WB

#### 2.1 Routine

• Eff. mAs: 170

• kV: 120

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

CTDlvol: 11.4779mGyScan time: 18.670 s

Delay: 4.000 sSlice: 5.00 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

#### 2.2 Scan

• Quality ref. mAs: 170

• Eff. mAs: 170

• kV: 120

Scan time: 18.670 sRotation time: 0.500 s

Delay: 4.000 sSlice: 5.00 mmPitch: 0.8000000

• Direction: Craniocaudal

#### 2.3 Recons

#### 2.3.1 Recon 1

- Series description: tmp CT WB RTD
- Slice: 5.00
- Kernel: B19f PET very smooth
- Window: AbdomenExtended FoV: On
- FoV: 700
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 11.500
- No. of images: 72

#### 2.3.2 Recon 2

- Series description: AC CT
- Slice: 5.00
- Kernel: B19f PET very smooth
- Window: AbdomenExtended FoV: On
- FoV: 700
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 3.000
- No. of images: 274

#### 2.3.3 Recon 3

- Series description: CT WB 3.0 B30f
- Slice: 3.00
- Kernel: B30f medium smooth
- Window: Abdomen
- Extended FoV: Off

- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 2.000
- No. of images: 411

### 3 Pause

## 4 PET WB

## 4.1 Routine

- $\bullet$  Isotope: F-18
- Pharm.: FDG
- Inj. Dose: 10 Millicuries
- Scan mode: Sinogram
- Scan range: Match CT FOV
- No. of beds: 5
- Scan duration/bed: 2 Minutes

#### 4.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: Trues
- 7
- Input trigger signal: None
- LLD (keV): 435
- ULD (keV): 650

#### 4.3 Recons

#### 4.3.1 Recon 1

- Series description: PET WB
- Recon range (bed): -1 to -1
- Output image type: Corrected
- Recon method: TrueX
- Iterations: 3
- Subsets: 21
- Image size: 168

- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

#### 4.3.2 Recon 2

- Series description: PET WB Uncorrected
- Recon range (bed): -1 to -1
- Output image type: Uncorrected
- Recon method: TrueX
- Iterations: 3
- Subsets: 21
- Image size: 168
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off