# $HUND\_64Cu\_Liposom\_Dag1.MlAdult\_PET5$

30. maj 2013

## Indhold

T	тор	ogram
	1.1	Routine
	1.2	Scan
<b>2</b>	$\mathbf{CT}$	1
	2.1	Routine
	2.2	Scan
	2.3	Recons
		2.3.1 Recon 1
		2.3.2 Recon 2
		2.0.2 1(0.001 2
3	Pau	se 2
4	PE	Γ LIVER
	4.1	Routine
	4.2	Scan
	4.3	Recons
	2.0	4.3.1 Recon 1
5	Pau	${f se}$
6	PET	Γ TUMOR
	6.1	Routine
	6.2	Scan
	6.3	Recons
		6.3.1 Recon 1

## 1 Topogram

## 1.1 Routine

mA: 35kV: 120

 $\bullet$  Topogram length: 0 mm

• Tube position: Top

## 1.2 Scan

mA: 35kV: 120Delay: 4s

Topogram length: 0 mmDirection: Craniocaudal

• Tube position: Top

API: NoneKernel: 80s shar

• Window: Topogram Body

## 2 CT

## 2.1 Routine

• Eff. mAs: 170

• kV: 120

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

CTDlvol: 11.4448mGyScan time: 12.680 s

Delay: 4.000 sSlice: 3.00 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

## 2.2 Scan

• Quality ref. mAs: 225

• Eff. mAs: 170

• kV: 120

Scan time: 12.680 sRotation time: 0.500 s

Delay: 4.000 sSlice: 3.00 mmPitch: 1.2000000

• Direction: Craniocaudal

#### 2.3 Recons

#### 2.3.1 Recon 1

- Series description: AC CT
- Slice: 3.00
- Kernel: B19f PET very smooth
- Window: AbdomenExtended FoV: On
- FoV: 700
- Center X: 0Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 3.000
- No. of images: 274

#### 2.3.2 Recon 2

- Series description: CT 3.0 B30f medium smooth
- Slice: 3.00
- Kernel: B30f medium smooth
- Window: AbdomenExtended FoV: Off
- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 3.000
- No. of images: 274

## 3 Pause

## 4 PET LIVER

## 4.1 Routine

• Isotope: Cu-64

- Pharm.: liposome
- Inj. Dose: 1 MegaBequerels
- Scan mode: List mode
- Scan range: Match CT FOV
- No. of beds: 1
- Scan duration/bed: 40 Minutes

## 4.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: List mode
- Sinogram mode: Trues
- Input trigger signal: None
- LLD (keV): 435
- ULD (keV): 650

## 4.3 Recons

#### 4.3.1 Recon 1

- Series description: PET WB Image check
- Recon range (bed): -1 to -1
- Output image type: NoRecon
- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 128
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

## 5 Pause

## 6 PET TUMOR

## 6.1 Routine

- Isotope: Cu-64
- Pharm.: liposome

- Inj. Dose: 1 MegaBequerels
- Scan mode: List mode
- Scan range: Match CT FOV
- No. of beds: 1
- Scan duration/bed: 20 Minutes

## 6.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: List mode
- Sinogram mode: Trues
- Input trigger signal: None
- LLD (keV): 435
- ULD (keV): 650

## 6.3 Recons

#### 6.3.1 Recon 1

- Series description: PET WB Image check
- Recon range (bed): -1 to -1
- Output image type: NoRecon
- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 128
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off