$4_PET_CT_WB_FDG_Terapi_XL_med_IV_Bolus.Adult_PET4$

3. december 2012

Indhold

T	Topogram			
	1.1	Routine		
	1.2	Scan		
2	Pre	Monitoring 1		
	2.1	Routine		
	2.2	Scan		
	2.3	Recons		
		2.3.1 Recon 1		
3	Mo	nitoring		
	3.1	Routine		
	3.2	Scan		
	3.3	Recons		
		3.3.1 Recon 1		
4	CT WB			
	4.1	Routine		
	4.2	Scan		
	4.3	Recons		
		4.3.1 Recon 1		
		4.3.2 Recon 2		
		4.3.3 Recon 3		
		4.3.4 Recon 4		
		4.3.5 Recon 5		
5	Pau	${f se}$		
6	PE	Γ WB		
	6.1	Routine		
	6.2	Scan		
	6.3	Recons		
		6.3.1 Recon 1		
		632 Recon 2		

1 Topogram

1.1 Routine

• mA: 35

• kV: 120

• Topogram length: 1024 mm

• Tube position: Top

1.2 Scan

• mA: 35

• kV: 120

• Delay: 4s

• Topogram length: 1024 mm

• Direction: Craniocaudal

• Tube position: Top

• API: API

• Kernel: 80

• Window: Topogram

2 PreMonitoring

2.1 Routine

• Eff. mAs: 20

• kV: 140

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

• CTDlvol: 1.782mGy

• Scan time: 0.500 s

• Delay: 2.000 s

• Slice: 10 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

2.2 Scan

• Quality ref. mAs: 225

• Eff. mAs: 20

• kV: 140

 \bullet Scan time: 0.500 s

• Rotation time: 0.500 s

• Delay: 2.000 s

• Slice: 10 mm

• Pitch: 0.00

• Direction: Craniocaudal

2.3 Recons

2.3.1 Recon 1

• Series description: PreMonitoring 10.0 eFoV

• Slice: 10

• Kernel: B19s

• Window: Abdomen

• Extended FoV: On

• FoV: 780

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 10.000

• No. of images: 1

3 Monitoring

3.1 Routine

• Eff. mAs: 20

• kV: 140

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

• CTDlvol: 53.46mGy

 \bullet Scan time: 0.500 s

• Delay: 10.000 s

• Slice: 10 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

3.2 Scan

• Quality ref. mAs: 225

• Eff. mAs: 20

• kV: 140

 \bullet Scan time: 0.500 s

• Rotation time: 0.500 s

• Delay: 10.000 s

• Slice: 10 mm

• Pitch: 0.00

• Direction: Craniocaudal

3.3 Recons

3.3.1 Recon 1

• Series description: Monitoring 10.0 eFoV

• Slice: 10

• Kernel: B19s

• Window: Abdomen

• Extended FoV: On

• FoV: 780

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 10.000

• No. of images: 30

4 CT WB

4.1 Routine

• Eff. mAs: 240

• kV: 140

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

 \bullet CTDlvol: 26.256mGy

• Scan time: 24.990 s

• Delay: 4.000 s

• Slice: 2 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

4.2 Scan

• Quality ref. mAs: 240

• Eff. mAs: 240

• kV: 140

• Scan time: 24.990 s

• Rotation time: 0.500 s

• Delay: 4.000 s

• Slice: 2 mm

• Pitch: 1.05

• Direction: Craniocaudal

4.3 Recons

4.3.1 Recon 1

- Series description: tmp CT WB RTD
- Slice: 4
- Kernel: B19f
- Window: Abdomen
- Extended FoV: Off
- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 4.000
- No. of images: 245

4.3.2 Recon 2

- Series description: AC CT 2 mm
- Slice: 2
- Kernel: B19f
- Window: Abdomen
- Extended FoV: On
- FoV: 780
- Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 2.000

• No. of images: 489

4.3.3 Recon 3

• Series description: CT WB 2.0 B30f

• Slice: 2

• Kernel: B30f

• Window: Abdomen

• Extended FoV: Off

• FoV: 500

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 2.000

• No. of images: 489

4.3.4 Recon 4

 $\bullet\,$ Series description: CT WB 2.0 SPO cor

• Slice: 2

• Kernel: B30f

• Window: Abdomen

• Extended FoV: Off

• FoV: 977

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: 3D

• Recon Axis: Coronal

• Image order: AP

• Recon increment: 2.000

• No. of images: 251

4.3.5 Recon 5

• Series description: CT WB 2.0 SPO sag

• Slice: 2

• Kernel: B30f

• Window: Abdomen

• Extended FoV: Off

• FoV: 977

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: 3D

• Recon Axis: Sagittal

• Image order: LeftToRight

• Recon increment: 2.000

• No. of images: 489

5 Pause

6 PET WB

6.1 Routine

• Isotope: F-18

• Pharm.: Fluorodeoxyglucose

 $\bullet\,$ Inj. Dose: 400000000.000000 Bequerels

 $\bullet\,$ Inj. date (date/month - year): 14/05 - 2012

 $\bullet\,$ Inj. time: 14:20:43

• Scan mode: Sinogram

• Scan range: Match CT FOV

 $\bullet\,$ No. of beds: Not given. Check recon range

• Scan duration/bed: 2.000000 Minutes

6.2 Scan

• Autoload: On

• Rebinner LUT: N/A

• Scan output: Sinogram

- Sinogram mode: false
- Input trigger signal: None
- LLD (keV): N/A
- ULD (keV): N/A

6.3 Recons

6.3.1 Recon 1

- Series description: PET WB Image check HD
- Recon range (bed): 1 to 7
- Output image type: Corrected
- Recon method: TrueX
- Iterations: 2
- Subsets: 12
- Image size: 200
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

6.3.2 Recon 2

• Series description: PET WB Corr TrueX+TOF 2 mm

• Recon range (bed): 1 to 7

• Output image type: Corrected

• Recon method: TrueXTOF

• Iterations: 2

• Subsets: 21

• Image size: 400

• Zoom: 1

• Filter: Gaussian

• FWHM (mm): 2.000000

• Offset X: 0 mm

• Offset Y: 0 mm

• Attenuation correction: Off (1)

• Scatter correction: On

• Match CT slice location: On

• Save intermediate data: Off