$4_PET_CT_WB_FDG_Terapi_IV_Breathold.Adult_PET4$

7. marts 2013

Indhold

1	Top	ogram				Topogram 1.1 Routine																											1
	1.1	Routin	ne																														1
	1.2	Scan										•		•									•								•		1
2	3																		1														
	2.1	Routin	ne																														1
	2.2	Scan																															1
	2.3	Recon	s																														2
		2.3.1	Recon 1																					•									2
3	Monitoring															2																	
	3.1		ne																														2
	3.2	Scan																															2
	3.3		S																														3
		3.3.1	Recon 1																														3
4	WP	вст																															3
-	4.1	Routir	ne																														3
	4.2																																3
	4.3		 s																														4
	4.0	4.3.1	Recon 1																														4
		4.3.2	Recon 2																														4
		4.3.3	Recon 3																														4
		4.3.4	Recon 4																														4
		4.3.4 $4.3.5$																															
		4.5.5	Recon 5		•	•	•	•		•	•	•	•	•	•	•	•	•		 •	•	•	•	•	•	•	•	•	•	•	•	•	5
5	Pau	ise																															6
6																6																	
	6.1	Routin	ne																														6
	6.2	Scan																															6
	6.3	Recon	s																														6
		6.3.1	Recon 1																														6
		6.3.2	Recon 2																														7
7	Pau	ıse																															7
8	\mathbf{CT}	Lunge																															7
	8.1	Routin	ne																														7
	8.2	Scan																															7
	83	Recon	C																														\$

		8.3.1	Recon 1																8
		8.3.2	Recon 2																8
		8.3.3	Recon 3																8
9	Pau	se																	9
10	PET	Γ Lung	ge :																9
	10.1	Routin	ie																9
	10.2	Scan																	g
	10.3	Recons	3																g
		10.3.1	Recon 1																g
		10.3.2	Recon 2																10
		10.3.3	Recon 3																10
11	Pau	\mathbf{se}																	11
12	PET	Γ Lung	ge :																11
	12.1	Routin	ie																11
	12.2	Scan																	11
	12.3	Recons	5																11
		12.3.1	Recon 1																11
		12.3.2	Recon 2																12
		1999	Dogge 9																10

1 Topogram

1.1 Routine

mA: 35kV: 120

• Topogram length: 1024 mm

• Tube position: Top

1.2 Scan

mA: 35kV: 120Delay: 4s

Topogram length: 1024 mmDirection: Craniocaudal

• Tube position: Top

API: APIKernel: 80

• Window: Topogram

2 PreMonitoring

2.1 Routine

• Eff. mAs: 20

• kV: 120

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

CTDlvol: 1.18627mGyScan time: 0.500 s

• Delay: 2.000 s

• Slice: 10 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

2.2 Scan

• Quality ref. mAs: 225

• Eff. mAs: 20

• kV: 120

Scan time: 0.500 sRotation time: 0.500 s

Delay: 2.000 sSlice: 10 mmPitch: 0.00

• Direction: Craniocaudal

2.3 Recons

2.3.1 Recon 1

• Series description: PreMonitoring 10.0 eFoV

Slice: 10Kernel: B19s

Window: AbdomenExtended FoV: On

FoV: 780Center X: 0Center Y: 0Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: CraniocaudalRecon increment: 10.000

• No. of images: 1

3 Monitoring

3.1 Routine

• Eff. mAs: 20

• kV: 120

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

CTDlvol: 35.5881mGy
 Scan time: 0.500 s

Delay: 10.000 sSlice: 10 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

3.2 Scan

• Quality ref. mAs: 225

Eff. mAs: 20kV: 120

Scan time: 0.500 sRotation time: 0.500 s

Delay: 10.000 sSlice: 10 mm

• Pitch: 0.00

• Direction: Craniocaudal

3.3 Recons

3.3.1 Recon 1

• Series description: Monitoring 10.0 eFoV

Slice: 10Kernel: B19s

Window: AbdomenExtended FoV: On

FoV: 780Center X: 0Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: CraniocaudalRecon increment: 10.000

• No. of images: 30

4 WB CT

4.1 Routine

• Eff. mAs: 200

• kV: 120

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

CTDlvol: 14.6179mGyScan time: 32.745 s

• Delay: 4.000 s

• Slice: 2 mm

 $\bullet\,$ No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

4.2 Scan

• Quality ref. mAs: 200

• Eff. mAs: 200

• kV: 120

Scan time: 32.745 sRotation time: 0.500 s

• Delay: 4.000 s

Slice: 2 mmPitch: 0.80

• Direction: Craniocaudal

4.3 Recons

4.3.1 Recon 1

• Series description: WB CT RTD

• Slice: 4

• Kernel: B19f

Window: AbdomenExtended FoV: Off

FoV: 500Center X: 0Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: CraniocaudalRecon increment: 4.000

• No. of images: 245

4.3.2 Recon 2

• Series description: AC CT 2 mm

• Slice: 2

• Kernel: B19f

Window: AbdomenExtended FoV: On

FoV: 780Center X: 0Center Y: 0Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

• Image order: Craniocaudal

 \bullet Recon increment: 2.000

• No. of images: 489

4.3.3 Recon 3

• Series description: CT WB 2.0 B30f

• Slice: 2

• Kernel: B30f

- Window: AbdomenExtended FoV: Off
- FoV: 500Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: AxialRecon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 2.000
- No. of images: 489

4.3.4 Recon 4

- Series description: CT WB 2.0 cor B30f
- Slice: 2
- Kernel: B30f
- Window: Abdomen
 Extended FoV: Off
- Extended FoV: Off
- FoV: 977
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: 3D
- Recon Axis: Coronal
- Image order: AP
- Recon increment: 2.000
- No. of images: 251

4.3.5 Recon 5

- Series description: CT WB 2.0 sag B30f
- Slice: 2
- Kernel: B30f
- Window: Abdomen
- Extended FoV: Off
- FoV: 977
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: 3D
- Recon Axis: Sagittal

- Image order: LeftToRight
 Recon increment: 2.000
- No. of images: 489

5 Pause

6 PET WB

6.1 Routine

- Isotope: F-18
- Pharm.: FluorodeoxyglucoseInj. Dose: 265000000 Bequerels
- Scan mode: Sinogram
- Scan range: Do not match CT FOV
- No. of beds: Not given. Check recon range
- Scan duration/bed: 2.000000 Minutes

6.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: false
- Input trigger signal: None
- LLD (keV): n/a
- ULD (keV): n/a

6.3 Recons

6.3.1 Recon 1

- Series description: PET WB Image check HD
- Recon range (bed): 1 to 7
- Output image type: Corrected
- Recon method: TrueX
- Iterations: 2
- Subsets: 12
- Image size: 200
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On

- Match CT slice location: On
- Save intermediate data: Off

6.3.2 Recon 2

- Series description: PET WB Corr TrueX+TOF 2 mm
- Recon range (bed): 1 to 7
- Output image type: Corrected
- Recon method: TrueXTOF
- Iterations: 2
- Subsets: 21
- Image size: 400
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 2.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

7 Pause

8 CT Lunge

8.1 Routine

- Eff. mAs: 225
- kV: 120
- CARE Dose4D: On
- CareDoseType: CareDoseAEC
- CTDlvol: 17.1537mGy
- Scan time: 0.933 s
- Delay: 10.000 s
- Slice: 3 mm
- No. of images: Samme som i foerste recon, slet?(y/n)
- Tilt: 0.0 grader

8.2 Scan

- Quality ref. mAs: 225
- Eff. mAs: 225
- kV: 120
- Scan time: 0.933 s
- Rotation time: 0.500 s

- Delay: 10.000 sSlice: 3 mmPitch: 1.40
- Direction: Caudocranial

8.3 Recons

8.3.1 Recon 1

- Series description: CT Lunge RTD
- Slice: 5
- Kernel: B18f
- Window: MediastinumExtended FoV: Off
- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Caudocranial
- Recon increment: 5.000
- No. of images: 4

8.3.2 Recon 2

- Series description: CT Lunge 3.0 eFoV
- Slice: 3
- Kernel: B18f
- Window: Mediastinum
- Extended FoV: On
- FoV: 780
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 3.000
- No. of images: 1

8.3.3 Recon 3

- Series description: CT CT Lunge 3.0 B70f
- Slice: 3

Kernel: B70fWindow: LungExtended FoV: Off

FoV: 331Center X: 0Center Y: 0Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: CraniocaudalRecon increment: 3.000

• No. of images: 1

9 Pause

10 PET Lunge

10.1 Routine

• Isotope: F-18

Pharm.: FluorodeoxyglucoseInj. Dose: 300000000 Bequerels

• Scan mode: Sinogram

• Scan range: Do not match CT FOV

• No. of beds: Not given. Check recon range

• Scan duration/bed: 20.000000 Seconds

10.2 Scan

• Autoload: Off

Rebinner LUT: OffScan output: SinogramSinogram mode: false

• Input trigger signal: No

• Input trigger signal: None

LLD (keV): n/aULD (keV): n/a

10.3 Recons

10.3.1 Recon 1

• Series description: PET Lunge Iterativ

• Recon range (bed): 1 to 1

• Output image type: Corrected

• Recon method: Iterative3D

- Iterations: 4
- Subsets: 8
- Image size: 200
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

10.3.2 Recon 2

- Series description: PET Lunge TrueX+TOF
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 3
- Subsets: 24
- Image size: 400
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

10.3.3 Recon 3

- Series description: PET Lunge OSEM
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 4
- Subsets: 8
- Image size: 256
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm

- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

11 Pause

12 PET Lunge

12.1 Routine

- Isotope: F-18
- Pharm.: Fluorodeoxyglucose
- Inj. Dose: 300000000 Bequerels
- Scan mode: Sinogram
- Scan range: Do not match CT FOV
- No. of beds: Not given. Check recon range
- Scan duration/bed: 20.000000 Seconds

12.2 Scan

- Autoload: Off
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: false
- Input trigger signal: None
- LLD (keV): n/a
- ULD (keV): n/a

12.3 Recons

12.3.1 Recon 1

- Series description: PET Lunge Iterativ
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 4
- Subsets: 8
- Image size: 200
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm

- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

12.3.2 Recon 2

- Series description: PET Lunge TrueX+TOF
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 3
- Subsets: 24
- Image size: 400
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

12.3.3 Recon 3

- Series description: PET Lunge OSEM
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 4
- Subsets: 8
- Image size: 256
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off