# $2\_PET\_AC\_WB\_FDG.Adult\_PET4$

7. marts 2013

# Indhold

| 1 | Topogram      |               |   |  |  |
|---|---------------|---------------|---|--|--|
|   | 1.1           | Routine       | 1 |  |  |
|   | 1.2           | Scan          | 1 |  |  |
| 2 | LavdoseCTWB 1 |               |   |  |  |
|   | 2.1           | Routine       | 1 |  |  |
|   | 2.2           | Scan          | 1 |  |  |
|   | 2.3           | Recons        | 2 |  |  |
|   |               | 2.3.1 Recon 1 | 2 |  |  |
|   |               | 2.3.2 Recon 2 | 2 |  |  |
| 3 | Pau           | se            | 2 |  |  |
| 4 | PET           | $^{\circ}$ WB | 2 |  |  |
|   | 4.1           | Routine       | 2 |  |  |
|   | 4.2           | Scan          | 3 |  |  |
|   | 4.3           | Recons        | 3 |  |  |
|   |               | 4.3.1 Recon 1 | 3 |  |  |
|   |               | 4.3.2 Recon 2 | 3 |  |  |

# 1 Topogram

## 1.1 Routine

mA: 35kV: 120

• Topogram length: 1024 mm

• Tube position: Top

## 1.2 Scan

mA: 35kV: 120Delay: 4s

Topogram length: 1024 mmDirection: Craniocaudal

• Tube position: Top

API: APIKernel: 80

• Window: Topogram

# 2 LavdoseCTWB

#### 2.1 Routine

• Eff. mAs: 50

• kV: 120

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

CTDlvol: 3.65446mGyScan time: 32.745 s

Delay: 2.000 sSlice: 2 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

## 2.2 Scan

• Quality ref. mAs: 50

• Eff. mAs: 50

• kV: 120

Scan time: 32.745 sRotation time: 0.500 s

Delay: 2.000 sSlice: 2 mmPitch: 0.80

• Direction: Craniocaudal

## 2.3 Recons

#### 2.3.1 Recon 1

- Series description: LavdoseCTWB RTD
- Slice: 4
- Kernel: B19f
- Window: AbdomenExtended FoV: Off
- FoV: 500
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 4.000
- No. of images: 245

#### 2.3.2 Recon 2

- Series description: AC lavCT 2 mm
- Slice: 2
- Kernel: B19f
- Window: Abdomen
- Extended FoV: On
- FoV: 780
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: Axial
- Recon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 2.000
- No. of images: 489

# 3 Pause

## 4 PET WB

# 4.1 Routine

• Isotope: F-18

- Pharm.: Fluorodeoxyglucose
- Inj. Dose: 300000000 Bequerels
- Scan mode: Sinogram
- Scan range: Do not match CT FOV
- No. of beds: Not given. Check recon range
- Scan duration/bed: 2.000000 Minutes

#### 4.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: false
- Input trigger signal: None
- LLD (keV): n/a
- ULD (keV): n/a

#### 4.3 Recons

#### 4.3.1 Recon 1

- Series description: PET WB Image check HD
- Recon range (bed): 1 to 7
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 4
- Subsets: 8
- Image size: 200
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

#### 4.3.2 Recon 2

- Series description: PET WB Corr TrueX+TOF 2 mm
- Recon range (bed): 1 to 7
- Output image type: Corrected
- Recon method: TrueXTOF
- Iterations: 2
- Subsets: 21

• Image size: 400

• Zoom: 1

• Filter: Gaussian

• FWHM (mm): 2.000000

Offset X: 0 mmOffset Y: 0 mm

• Attenuation correction: Off ()

• Scatter correction: On

Match CT slice location: On Save intermediate data: Off