$PROJ_Tweak_FDG_PET_5.MlAdult_PET5$

30. maj 2013

Indhold

| 1 | 1 Topogram | Topogram | | | | | | | | | | | | | | 1 | |
|----------|--------------------|----------|--|--|--|--|--|--|--|--|--|--|--|---|--|---|---|
| | 1.1 Routine | | | | | | | | | | | | | | | |] |
| | 1.2 Scan | | | | | | | | | | | | | | | • |] |
| 2 | 2 Lavdos CT WB | | | | | | | | | | | | | | | | 1 |
| | 2.1 Routine | | | | | | | | | | | | | | | |] |
| | 2.2 Scan \dots | | | | | | | | | | | | | | | | - |
| | 2.3 Recons | | | | | | | | | | | | | | | | 4 |
| | 2.3.1 Recon 1 | L | | | | | | | | | | | | | | | 4 |
| 3 | 3 Pause | | | | | | | | | | | | | | | | 6 |
| 4 | 4 PET WB | PET WB | | | | | | | | | | | | 2 | | | |
| | 4.1 Routine | | | | | | | | | | | | | | | | 4 |
| | 4.2 Scan \dots | | | | | | | | | | | | | | | | 4 |
| | 4.3 Recons | | | | | | | | | | | | | | | | • |
| | 4.3.1 Recon 1 | l | | | | | | | | | | | | | | | • |
| | 4.3.2 Recon 2 | 2 | | | | | | | | | | | | | | | • |
| | 4.3.3 Recon 3 | 3 | | | | | | | | | | | | | | | ٠ |

1 Topogram

1.1 Routine

mA: 35kV: 120

• Topogram length: 1024 mm

• Tube position: Top

1.2 Scan

mA: 35kV: 120Delay: 4s

Topogram length: 1024 mmDirection: Craniocaudal

• Tube position: Top

API: NoneKernel: 80s shar

• Window: Topogram Body

2 Lavdos CT WB

2.1 Routine

• Eff. mAs: 40

• kV: 120

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

CTDlvol: 2.70069mGyScan time: 18.670 s

Delay: 4.000 sSlice: 3.00 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

2.2 Scan

• Quality ref. mAs: 50

• Eff. mAs: 40

• kV: 120

Scan time: 18.670 sRotation time: 0.500 s

Delay: 4.000 sSlice: 3.00 mmPitch: 0.8000000

• Direction: Craniocaudal

2.3 Recons

2.3.1 Recon 1

• Series description: AC CT

• Slice: 3.00

• Kernel: B19f PET very smooth

Window: AbdomenExtended FoV: On

FoV: 700Center X: 0Center Y: 0Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: Craniocaudal
Recon increment: 3.000
No. of images: 274

3 Pause

4 PET WB

4.1 Routine

Isotope: F-18 Pharm.: FDG

• Inj. Dose: 400 MegaBequerels

• Scan mode: Sinogram

• Scan range: Match CT FOV

• No. of beds: 1

• Scan duration/bed: 150 Seconds

4.2 Scan

• Autoload: On

Rebinner LUT: OffScan output: SinogramSinogram mode: Trues

• Input trigger signal: None

LLD (keV): 435ULD (keV): 650

4.3 Recons

4.3.1 Recon 1

- Series description: PET WB Corrected
- Recon range (bed): -1 to -1
- Output image type: Corrected
- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 168
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 4
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

4.3.2 Recon 2

- Series description: PET WB Corr. True X 2 mm
- Recon range (bed): -1 to -1
- Output image type: Corrected
- Recon method: TrueX
- Iterations: 3
- Subsets: 21
- Image size: 336
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 2
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

4.3.3 Recon 3

- Series description: PET NAC
- Recon range (bed): -1 to -1
- Output image type: Uncorrected
- Recon method: Iterative

Iterations: 3 Subsets: 21 Image size: 336

• Zoom: 1

Filter: GaussianFWHM (mm): 2Offset X: 0 mmOffset Y: 0 mm

• Attenuation correction: On (1)

• Scatter correction: On

 $\bullet\,$ Match CT slice location: On

• Save intermediate data: Off