$4_PET_CT_FDG_Brain.Adult_PET4$

30. maj 2013

Indhold

1	Top	ogram																					
	1.1	Routin	ne																				
	1.2	Scan				•																	
2	CT	Brain																					
	2.1	Routin	ne																				
	2.2	Scan																 					
	2.3	Recons	5																				
		2.3.1	Recon 1																				
		2.3.2	Recon 2															 					
		2.3.3	Recon 3																	•			
3	Pau	ıse																					
4	PET Brain																						
	4.1	Routin	ne															 					
	4.2	Scan																 					
	4.3		3																				
		4.3.1	Recon 1																				
		4.3.2	Recon 2															 					

1 Topogram

1.1 Routine

mA: 35kV: 120

Topogram length: 256 mmTube position: Lateral

1.2 Scan

mA: 35kV: 120Delay: 4s

Topogram length: 256 mmDirection: CraniocaudalTube position: Lateral

API: APIKernel: 20

• Window: Topogram

2 CT Brain

2.1 Routine

• Eff. mAs: 380

• kV: 120

• CARE Dose4D: On

• CareDoseType: CareDoseAEC

CTDlvol: 64.9045mGyScan time: 69.697 s

Delay: 2.000 sSlice: 2 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

2.2 Scan

• Quality ref. mAs: 380

• Eff. mAs: 380

• kV: 120

Scan time: 69.697 sRotation time: 1.000 s

Delay: 2.000 sSlice: 2 mmPitch: 0.55

• Direction: Craniocaudal

2.3 Recons

2.3.1 Recon 1

• Series description: CT Brain RTD

Slice: 1.5Kernel: H19s

Window: CerebrumExtended FoV: Off

FoV: 300Center X: 0Center Y: 0Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: Craniocaudal
Recon increment: 1.500

• No. of images: 149

2.3.2 Recon 2

• Series description: AC CT

• Slice: 2

• Kernel: H19s

Window: CerebrumExtended FoV: Off

FoV: 300Center X: 0Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: CraniocaudalRecon increment: 2.000

• No. of images: 112

2.3.3 Recon 3

• Series description: Cranie VIR 2.0 H30S

• Slice: 2

• Kernel: H30s

Window: CerebrumExtended FoV: Off

- FoV: 300
- Center X: 0
- Center Y: 0
- Mirroring: None
- Extended CT scale: Standard
- Recon job: AxialRecon Axis: Axial
- Image order: Craniocaudal
- Recon increment: 2.000
- No. of images: 112

3 Pause

4 PET Brain

4.1 Routine

- Isotope: F-18
- Pharm.: Fluorodeoxyglucose
- Inj. Dose: 200000000 Bequerels
- Scan mode: Sinogram
- Scan range: Do not match CT FOV
- No. of beds: Not given. Check recon range
- Scan duration/bed: 10.000000 Minutes

4.2 Scan

- Autoload: On
- Rebinner LUT: Off
- Scan output: Sinogram
- Sinogram mode: false
- Input trigger signal: None
- LLD (keV): n/a
- ULD (keV): n/a

4.3 Recons

4.3.1 Recon 1

- Series description: PET Brain Corrected 3 mm zoom 3
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- Iterations: 4
- Subsets: 24
- Image size: 400

- Zoom: 3.000000Filter: Gaussian
- FWHM (mm): 3.000000
- Offset X: 0 mmOffset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

4.3.2 Recon 2

- Series description: Neurostat
- Recon range (bed): 1 to 1
- Output image type: Corrected
- Recon method: Iterative3D
- \bullet Iterations: 4
- Subsets: 8
- Image size: 128
- Zoom: 3.000000
- Filter: Gaussian
- FWHM (mm): 8.000000
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: Off ()
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off