TUMOR_HYPOXI_HUND_LM1.MlAdult_PET5

7. marts 2013

Indhold

1	Topogram			
	1.1	Routine	1	
	1.2	Scan	1	
2	\mathbf{CT}		1	
	2.1	Routine	1	
	2.2	Scan	1	
	2.3	Recons	2	
		2.3.1 Recon 1	2	
3	Pau	ise	2	
4	PET	Γ LM	2	
	4.1	Routine	2	
	4.2	Scan	2	
	4.3		3	
		4.3.1 Recon 1	3	

1 Topogram

1.1 Routine

mA: 35kV: 120

• Topogram length: 1024 mm

• Tube position: Top

1.2 Scan

mA: 35kV: 120Delay: 4s

Topogram length: 1024 mmDirection: Craniocaudal

• Tube position: Top

API: NoneKernel: 80s shar

• Window: Topogram Body

2 CT WB

2.1 Routine

• Eff. mAs: 11

• kV: 120

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

CTDlvol: 0.7414mGy
 Scan time: 3.360 s

Delay: 4.000 sSlice: 3.00 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 0.0 grader

2.2 Scan

• Quality ref. mAs: 170

• Eff. mAs: 11

• kV: 120

Scan time: 3.360 sRotation time: 0.500 s

Delay: 4.000 sSlice: 3.00 mmPitch: 1.5000000

• Direction: Craniocaudal

2.3 Recons

2.3.1 Recon 1

• Series description: AC CT

• Slice: 3.00

• Kernel: B18f very smooth

Window: AbdomenExtended FoV: On

FoV: 700Center X: 0Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

Recon job: AxialRecon Axis: Axial

Image order: Craniocaudal
Recon increment: 2.000
No. of images: 111

• No. of images: 111

3 Pause

4 PET LM

4.1 Routine

Isotope: Cu-64Pharm.: ATSM

• Inj. Dose: 10 MegaBequerels

• Scan mode: List mode

• Scan range: Match CT FOV

• No. of beds: 1

• Scan duration/bed: 60 Minutes

4.2 Scan

• Autoload: On

• Rebinner LUT: Off

Scan output: List modeSinogram mode: Trues

• Input trigger signal: ECG

LLD (keV): 435ULD (keV): 650

4.3 Recons

4.3.1 Recon 1

• Series description: PET Cardiac

• Recon range (bed): -1 to -1

• Output image type: NoRecon

• Recon method: Iterative

 \bullet Iterations: 4

• Subsets: 8

• Image size: 168

• Zoom: 2

• Filter: Gaussian

• FWHM (mm): 2

• Offset X: 0 mm

• Offset Y: 0 mm

• Attenuation correction: On (1)

• Scatter correction: On

• Match CT slice location: On

• Save intermediate data: Off