# 6\_TUMOR\_HYPOXI\_HUND\_DAG\_3.MlAdult

22. oktober 2012

# Indhold

1	Topogram		
	1.1	Routine	
	1.2	Scan	
2	$\mathbf{CT}$	TUMOR	
	2.1	Routine	
	2.2	Scan	
	2.3	Recons	
		2.3.1 Recon 1	
		2.3.2 Recon 2	
3	Pau	se 4	
4	PE	Γ STATIC	
	4.1	Routine	
	4.2	Scan	
	4.3	Recons	
		4.3.1 Recon 1	
		4.3.2 Recon 2	
5	Pau	se	
6	Dyr	m nMulti	
	6.1	Routine	
	6.2	Scan	
	6.3	Recons	
		6.3.1 Recon 1	

## 1 Topogram

## 1.1 Routine

• mA: 35

• kV: 120

• Topogram length: 512 mm

• Tube position: Top

#### 1.2 Scan

• mA: 35

• kV: 120

• Delay: 4s

• Topogram length: 270 mm

• Direction: Craniocaudal

• Tube position: Top

• API: None

• Kernel: 80s shar

• Window: Topogram Body

## 2 CT TUMOR

### 2.1 Routine

• Eff. mAs: 170

• kV: 120

• CARE Dose4D: Off

• CareDoseType: CareDoseAEC

• CTDlvol: 11.4448mGy

• Scan time: 4.000 s

• Delay: 4.000 s

• Slice: 3.00 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 170 grader

## 2.2 Scan

• Quality ref. mAs: 170

• Eff. mAs: 170

• kV: 120

 $\bullet$  Scan time: 4.000 s

• Rotation time: 0.500 s

• Delay: 4.000 s

• Slice: 3.00 mm

• Pitch: 1.2000000

• Direction: Craniocaudal

### 2.3 Recons

#### 2.3.1 Recon 1

• Series description: AC CT

• Slice: 3.00

• Kernel: B19f PET very smooth

• Window: Abdomen

• Extended FoV: On

• FoV: 700

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 3.000

• No. of images: 74

#### 2.3.2 Recon 2

• Series description: CT TUMOR 3.0 B30f

• Slice: 3.00

• Kernel: B30f medium smooth

• Window: Abdomen

• Extended FoV: Off

• FoV: 500

• Center X: 0

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 3.000

• No. of images: 74

## 3 Pause

## 4 PET STATIC

### 4.1 Routine

• Isotope: Cu-64

• Pharm.: ATSM

• Inj. Dose: 1e-006 MegaBequerels

 $\bullet$  Inj. date (date/month - year): 13/01 - 2009

• Inj. time: 16:49:00

• Scan mode: Sinogram

• Scan range: Match CT FOV

• No. of beds: 1

• Scan duration/bed: 20 Minutes

#### 4.2 Scan

• Autoload: On

• Rebinner LUT: Off

• Scan output: Sinogram

• Sinogram mode: Trues

• Input trigger signal: None

• LLD (keV): 435

• ULD (keV): 650

#### 4.3 Recons

#### 4.3.1 Recon 1

- Series description: PET TUMOR Corrected 24 hr
- Recon range (bed): Automatic/All/dunno
- Output image type: Corrected
- Recon method: Iterative
- Iterations: 4
- Subsets: 8
- Image size: 256
- Zoom: 1
- Filter: Gaussian
- FWHM (mm): 3
- Offset X: 0 mm
- Offset Y: 0 mm
- Attenuation correction: On (1)
- Scatter correction: On
- Match CT slice location: On
- Save intermediate data: Off

#### 4.3.2 Recon 2

- Series description: PET TUMOR Uncorrected 24 hr
- Recon range (bed): Automatic/All/dunno
- Output image type: Uncorrected
- Recon method: Iterative
- Iterations: 4
- Subsets: 8

• Image size: 256

• Zoom: 1

• Filter: Gaussian

• FWHM (mm): 3

• Offset X: 0 mm

• Offset Y: 0 mm

• Attenuation correction: On (1)

• Scatter correction: Off

• Match CT slice location: On

• Save intermediate data: Off

## 5 Pause

## 6 DynMulti

### 6.1 Routine

• Eff. mAs: 170

• kV: 120

• CARE Dose4D: Off

• CareDoseType: CareDose

• CTDlvol: 1032.37mGy

• Scan time: 90.100 s

• Delay: 4.000 s

• Slice: 7.20 mm

• No. of images: Samme som i foerste recon, slet?(y/n)

• Tilt: 170 grader

## 6.2 Scan

• Quality ref. mAs: 100

• Eff. mAs: 170

• kV: 120

• Scan time: 90.100 s

• Rotation time: 1.000 s

• Delay: 4.000 s

• Slice: 7.20 mm

• Pitch: 1.0000000

• Direction: Craniocaudal

#### 6.3 Recons

#### 6.3.1 Recon 1

• Series description: DynMulti 7.2 B30s

• Slice: 7.20

• Kernel: B30s medium smooth

• Window: Abdomen

• Extended FoV: Off

• FoV: 0

• Center X: -3

• Center Y: 0

• Mirroring: None

• Extended CT scale: Standard

• Recon job: Axial

• Recon Axis: Axial

• Image order: Craniocaudal

• Recon increment: 1.000

• No. of images: 360